



EpiList 1.0: a global checklist of vascular epiphytes

Zotz, Gerhard ; Weigelt, Patrick ; Kessler, Michael ; Kreft, Holger ; Taylor, Amanda

Abstract: Epiphytes make up roughly 10% of all vascular plant species globally and play important functional roles, especially in tropical forests. However, to date, there is no comprehensive list of vascular epiphyte species. Here, we present EpiList 1.0, the first global list of vascular epiphytes based on standardized definitions and taxonomy. We include obligate epiphytes, facultative epiphytes, and hemiepiphytes, as the latter share the vulnerable epiphytic stage as juveniles. Based on 978 references, the checklist includes >31,000 species of 79 plant families. Species names were standardized against World Flora Online for seed plants and against the World Ferns database for lycophytes and ferns. In cases of species missing from these databases, we used other databases (mostly World Checklist of Selected Plant Families). For all species, author names and IDs for World Flora Online entries are provided to facilitate the alignment with other plant databases, and to avoid ambiguities. EpiList 1.0 will be a rich source for synthetic studies in ecology, biogeography, and evolutionary biology as it offers, for the first time, a species-level overview over all currently known vascular epiphytes. At the same time, the list represents work in progress: species descriptions of epiphytic taxa are ongoing and published life form information in floristic inventories and trait and distribution databases is often incomplete and sometimes even wrong. Since the epiphytic growth blends into soil-rooted growth and vice versa, the inclusion or exclusion of particular species in the current list will sometimes be contentious. Thus, initiating a well-founded discussion was one of the motivations for compiling this database; our list represents 31,311 hypotheses on the life form of plant species, and we welcome feedback on possible omission or erroneous inclusions. We release these data into the public domain under a Creative Commons Zero license waiver. When you use the data in your publication, we request that you cite this data paper. If EpiList 1.0 is a major part of the data analyzed in your study, you may consider inviting the EpiList 1.0 core team as collaborators.

DOI: <https://doi.org/10.1002/ecy.3326>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-202729>

Journal Article

Accepted Version



The following work is licensed under a Creative Commons: Public Domain Dedication: CC0 1.0 Universal (CC0 1.0) License.

Originally published at:

Zotz, Gerhard; Weigelt, Patrick; Kessler, Michael; Kreft, Holger; Taylor, Amanda (2021). EpiList 1.0: a global checklist of vascular epiphytes. *Ecology*, 102(6):e03326.

DOI: <https://doi.org/10.1002/ecy.3326>

EpiList 1.0: a global checklist of vascular epiphytes

Authors:

Gerhard Zotz^{1,2}, Patrick Weigelt³, Michael Kessler⁴, Holger Kreft^{3,5}, & Amanda Taylor³

¹ Functional Ecology of Plants, Institute of Biology and Environmental Sciences, Carl von Ossietzky University, Ammerländer Heerstr. 114-118, 26129 Oldenburg, Germany;

² Smithsonian Tropical Research Institute, Apdo 2072 Balboa, Panama

³ Biodiversity, Macroecology & Biogeography, University of Göttingen, Büsgenweg 1, 37077 Göttingen, Germany;

⁴ Institute of Systematic and Evolutionary Botany, University of Zürich, Zollikerstrasse 107, 8008 Zürich, Switzerland

⁵ Centre of Biodiversity and Sustainable Land Use (CBL), University of Göttingen, Büsgenweg 1, 37077 Göttingen, Germany

Corresponding author:

Gerhard Zotz, University Oldenburg, Institute for Biology and Environmental Sciences, Functional Ecology Group, Box 2503, D-26111 Oldenburg, Germany

Tel.: +49-(0)441-798-3794

Fax: +49-(0)441-798-3331

e-mail: gerhard.zotz@uol.de

ORCID IDs

Gerhard Zotz: <https://orcid.org/0000-0002-6823-2268>

Patrick Weigelt: <https://orcid.org/0000-0002-2485-3708>

Michael Kessler: <https://orcid.org/0000-0003-4612-9937>

Holger Kreft: <https://orcid.org/0000-0003-4471-8236>

Amanda Taylor: <https://orcid.org/0000-0002-0420-2203>

Introduction

Epiphytes are defined as “plants that germinate and root non-parasitically on other plants at all stages of life” (Zotz 2016). Vascular epiphytes are most conspicuous in the tropics, particularly in montane tropical areas, where their species numbers may locally exceed those of all other life forms, like trees, shrubs or understory herbs, combined (Kelly et al. 1994), although epiphyte-rich vegetation is also found in some temperate regions (Hofstede et al. 2001, Díaz et al. 2010). When reaching such high diversity and abundance, vascular epiphytes play an important role in ecosystem processes like in forest hydrology (Van Stan et al. 2015, Hargis et al. 2019), nutrient fluxes (Nadkarni and Matelson 1991, Nadkarni and Matelson 1992, Richardson et al. 2000), and by provisioning food and shelter for canopy fauna (Melnychuk and Srivastava 2002, Stuntz et al. 2002, McCracken and Forstner 2014). This ecological importance has motivated a large number of studies and a rich body of literature on vascular epiphytes has accumulated: a recent monograph (Zotz 2016) summarizes the knowledge of all aspects of the biology of this plant group.

With the onset of the scientific study of vascular epiphytes (Schimper 1888), researchers have started to enumerate epiphytic taxa. For a given location, this is necessary for, e.g., quantitative comparisons of the local or regional diversity of life forms (Whitmore et al. 1985, Balslev et al. 1998, Linares-Palomino et al. 2009), while more comprehensive lists for entire plant families or for different geographical entities up to the global scale are an indispensable resource for researchers who want to study evolutionary and / or (macro)ecological questions, regardless of whether the focus is clearly on vascular epiphytes (Cascante-Marín and Nivia-Ruiz 2013, Ramos et al. 2019) or whether a comparative approach with different life forms is used (Crayn et al. 2004, Givnish et al. 2014, Givnish et al. 2015). We note that such usage of epiphyte lists can only yield reproducible results if definitions are compatible. Unfortunately, this is frequently not the case. Some studies lump epiphytes with (hemiparasitic) mistletoes (e.g. Cascante-Marín and Estrada-Chavarría 2012, Fall and Drezner 2020), soil-rooted climbers (Nieder et al. 2000, Irupe et al. 2013, Engemann et al. 2016), or accidental epiphytes (e.g. Sugden and Robins 1979, Prescott et al. 2015), i.e. rare arboreal individuals of an otherwise terrestrial plant species. Such ambiguities are highly problematic because, e.g., differences in the vertical distribution patterns of “epiphyte” communities may simply be an artefact of the inclusion of climbers (compare, e.g., Nieder et al. 2000 vs. Zotz and Schultz 2008).

Thus, it is fundamental to address the problem of the delimitation of structurally dependent life forms. In the following, we provide detailed definitions of these different life forms, but

note from the start that an entirely unambiguous list of epiphytes is impossible for practical purposes: similar to many other categories in biology, e.g. “rheophytes” (Costa et al. 2020), “CAM-plants” (Zotz 2002), “halophytes” (Grigore et al. 2014), or “succulents” (Ogburn and Edwards 2010) there are numerous cases among epiphytes that defy a clear categorisation, e.g. species with individuals growing as both epiphytes and terrestrials in about equal numbers (“facultative epiphytes”).

There have been a number of previous attempts to compile global lists of epiphytes, most noteworthy Madison (1977), Gentry and Dodson (1987), Kress (1986), and Zotz (2013). All previous compilations gave only total numbers per family and genus, but lacked documentation for the individual species. Here, we go beyond these earlier efforts and present a list of more than 31,000 species with at least one reference from the literature or online databases in >99.9% of all cases to justify their inclusion. Unfortunately, these sources did not necessarily use the same implicit or explicit definition. This ambiguity will inevitably lead to some erroneous inclusions or omissions of species. Even though information from more than 7000 original papers, books, floras, gray literature, online platforms, webpages, and expert comments was used and critically reviewed (for details see: **Sampling Design and Research Methods**), a compilation like the current one depends on the quality and unambiguity of the original sources.

METADATA

Class I. Data set descriptors

I.A. Data set identity: The database presented here lists 31,311 vascular plant species. Each data entry includes family, genus and species names, taxonomic authority, and with very few exceptions at least one published source for each species. Species names were standardized as follows: The primary reference for seed plants was the taxonomic backbone of World Flora Online (wfo) version 1.2 from 10 October, 2019. The taxonomic backbone of wfo was initially based on the now discontinued Plant List (theplantlist.org), and is actively being curated by taxonomic specialists of particular plant groups. Since the original working list of epiphytes did not include authorities and most of the names had been manually standardized based on the Plant List when incorporated into the epiphyte list, the taxonomic names had to be matched to wfo names and their respective authorities. Genus names that could not be matched were checked and corrected manually. Species epithets were compared to all species epithets within a genus and best matching epithets were chosen based on orthographic distances. If several names matched equally well (either because of similar spelling or the

occurrence of several names with different authorities and/or infraspecific information) we combined all synonyms that would lead to the same accepted name and prioritized taxonomic names without infraspecific information over names at the species level, checked names over unchecked names and accepted names over synonyms. The results of this automated name assignment were checked for consistency by GZ. Names that matched poorly or still matched several names (e.g. synonyms of two different, accepted species names) were checked manually based on other global resources like Plants of the World Online (powo) or the World Checklist of Selected Plant Families (wmsp) or the original resource reporting the epiphyte species. Cases that could not be matched to any name in wfo typically were recent additions to the EpiList coming from wmsp. Those cases apparently have not yet been added to wfo and were adopted from wmsp. After name matching, synonyms were replaced by their corresponding accepted names, and infraspecific taxa were replaced by their parent species name and authority in wfo. Although we had initially planned to use wfo for ferns and lycophytes as well, many inconsistencies emerged and a very large proportion of species could not be matched. Therefore, we standardized fern and lycophyte names against WorldFerns 8.30 (Hassler and Schmitt 2019), which follows the family and genus taxonomy of the Pteridophyte Phylogeny Group (PPG I, Schuettpelz et al. 2016) with some recent updates.

I.B. Data set identification code: N/A

I.C. Data set description: N/A

I.C.1. Originators: Gerhard Zotz

I.C.2. Abstract: Epiphytes make up roughly 10% of all vascular plant species globally and play important functional roles, especially in tropical forests. However, to date there is no comprehensive list of vascular epiphyte species. Here, we present **EpiList 1.0**, the first global list of vascular epiphytes based on standardized definitions and taxonomy. We include obligate epiphytes, facultative epiphytes and hemiepiphytes, as the latter share the vulnerable epiphytic stage as juveniles. Based on 978 references, the checklist includes of > 31,000 species of 79 plant families. Species names were standardized against World Flora Online for seed plants and against the World Ferns database for lycophytes and ferns. In cases of species missing from these databases, we used other databases (mostly World Checklist of Selected Plant Families). For all species, author names and IDs for World Flora Online entries are provided to facilitate the alignment with other plant databases, and to avoid ambiguities.

EpiList 1.0 will be a rich source for synthetic studies in ecology, biogeography, and evolutionary biology as it offers - for the first time - a species-level overview over all

currently known vascular epiphytes. At the same time, the list represents work in progress: species descriptions of epiphytic taxa are ongoing, published life form information in floristic inventories and trait and distribution databases is often incomplete and sometimes even wrong. Since the epiphytic growth blends into soil-rooted growth, the inclusion or exclusion of particular species in the current list will sometimes be contentious. Thus, initiating a discussion was one of the motivations for compiling this database – our list represents 31,311 hypotheses on the life form of plant species and we welcome feedback on possible omission or erroneous inclusions. We release these data into the public domain under a Creative Commons Zero license waiver. When you use the data in your publication, we request that you cite this data paper. If **EpiList 1.0** is a major part of the data analyzed in your study, you may consider inviting the **EpiList 1.0** core team as collaborators. We release these data into the public domain under a Creative Commons Zero license waiver (<https://creativecommons.org/share-your-work/public-domain/cc0/>). When you use the data in your publication, we request that you cite this data paper. Should **EpiList 1.0** be a major part of the data analyzed in your study, you should consider inviting the **EpiList 1.0** core team (see Originators in the Overall project description) as collaborators. If you plan to use the **EpiList 1.0** dataset, we request that you contact the **EpiList 1.0** core team to learn whether updates of the dataset are available, and whether similar analyses are already ongoing.

I.D. Key words: *Araceae, biodiversity, Bromeliaceae, global distribution, hemiepiphyte, life form, nomadic vine, Orchidaceae, structural dependence, tropical, vascular plants*

Class II. Research origin descriptors

II.A. Overall project description: A list of all plant species that typically grow epiphytically (holoepiphytes and facultative epiphytes) and species that start epiphytically and later establish root contact with the soil (hemiepiphytes).

II.A.1. Identity: 31,311 vascular plant species.

II.A.2. Originators: see I.C.1.

II.A.3. Period of study: Literature and online data sources from 1851 to 2020.

II.A.4. Objectives: The purpose of the EpiList database is to provide a standardized, fully referenced species list of vascular epiphytes following up-to-date taxonomy to be used in ecological and evolutionary studies.

II.A.5. Abstract: see I.C.2.

II.A.6. Sources of funding: none

II.B. “Specific” subproject description: N/A

II.B.1. Site description: planet earth

II.B.2. Experimental or sampling design:

II.B.2.a. Design characteristics: See below

II.B.2.b. Permanent plots: N/A

II.B.2.c. Data collection period, frequency, etc.: 1851-2020

II.B.3. Research methods: See below

II.B.3.a. Field/laboratory:

Sampling Design and Research Methods

Information on the epiphytic occurrence of a species was reviewed in more than 7,000 sources. These include papers, books, book chapters, theses, reports, webpages, digital herbaria, and online databases such as wesp (2019), Flora of China (2008), Brazilian Flora 2020 in construction (continuously updated), and the Global Inventory of Floras and Traits (GIFT) database (Weigelt et al. 2020). In addition, we also contacted experts of many taxa and included their feedback (see acknowledgements and list in Zotz 2013). There was much overlap of species reported in these diverse sources – in most cases, we only provide one key source in **EpiList 1.0**, so that the actual number of sources appearing in the database is only 978.

Definitions

Taxonomic Identification: Family, genus and species names and authority are given. Because of taxonomic revisions or the use of synonyms, many of the names are different from those used in the original sources.

EpiList only includes species, whereas infraspecific taxa, hybrids (e.g. *Laelia* x *tlaxiacoensis* Solano et al. 2019) and variants were excluded. In two exceptional cases, we included unidentified species in **EpiList 1.0**. The first case represents the only report of an epiphytic Poaceae in the genus *Tripogon*. The second case is a *Loxostigma* species (Gesneriaceae), again the only report for this genus. Future studies will have to evaluate whether there really are epiphytic *Tripogon* and *Loxostigma* species or whether accidental epiphytic occurrences of terrestrial species were reported. However, even if it eventually turned out that these were indeed accidentals, reporting accidental epiphytic occurrences is still valuable, because the

study of accidental epiphytes may allow us to understand the first evolutionary steps towards true epiphytism (Hoeber et al. 2020).

Definition and Assignment of Mechanically Dependent Life Forms

A summary of five different types of mechanically dependent life forms is given in Fig. 1.

Since hemiepiphytes, nomadic vines, and mistletoes are sometimes lumped with true epiphytes (holoepiphytes), it is important to provide unambiguous definitions of these groups (Sperotto et al. 2020).




Figure 1: Graphical definition and assumed evolutionary connections (arrows) between different forms of structurally dependent plants. For each life form, (1) the site of germination, (2) the site of attachment of feeding roots during vegetative growth and (3) during reproduction are given (T = terrestrial, H = host tree). Each life form has a unique combination of (1) – (3), with the exception of mistletoes (grey arrows) and epiphytes. The former differ from epiphytes by their parasitic life style.

We define epiphytes as “plants that germinate and root non-parasitically on other plants at all stages of life” (Zotz 2016). While this definition can be unambiguously applied in the case of an individual plant, variation may exist within a species. Many species that typically grow on trees may also occur on rocks or soil as long as competition by terrestrial plants is low

(“facultative epiphytes”), while a considerable number of terrestrial plant species may sporadically grow on living substrate as “accidental epiphytes” (Hoeber et al. 2019). There are numerous reports in the literature in which such terrestrial species are included in “epiphyte” lists without qualifying comments (e.g. Sugden and Robins 1979, Prescott et al. 2015). While facultative epiphytes are included in **EpiList1.0**, the inclusion of accidentals would lead to an excessive and ecologically rather meaningless species inflation, the magnitude of which becomes apparent with the following number: more than 600 vascular plant species have been recorded as accidental epiphytes in Europe alone (V. Hoeber, M. Klinghardt, G. Zotz, unpubl.). Resolving this issue is relatively easy in reported cases of epiphytic bananas, epiphytic dandelions, or epiphytic red mangroves (G. Zotz, unpubl. obs.), but is much more challenging in less obvious cases: careful investigation of the typical growing site of a species is necessary. However, although there are schemes how to quantify the “arboreality” of a species (Ibisch 1996, Burns 2010), a lack of relevant observations and possible regional differences make an absolutely watertight list of epiphytes impossible.

The inclusion of facultative epiphytes in **EpiList 1.0** has important implications for its use. Most importantly, the inclusion of a species should not be taken as an indication that the species can always be treated as an epiphyte in a binary classification (epiphytic vs. non-epiphytic), because a species may be epiphytic or not in different parts of its range, which must be taken into account in a regional classification by life form. For example, while almost exclusively terrestrial in Europe, *Asplenium scolopendrium* is mainly epiphytic in much more humid forests in Myanmar (Khine et al. 2017).

Hemiepiphytes are plants that begin their life with an epiphytic stage, but later establish root contact with the soil (Fig. 1). They can be found in many families with epiphytic members (e.g. Rubiaceae, Urticaceae), but others are in families with few or no true epiphytes (e.g. Araliaceae, Moraceae). When studies lump “epiphytic” and “hemiepiphytic” species, it is hardly possible to make that distinction afterwards. We thus opted for the following approach: since all hemiepiphytes, by definition, share an epiphytic juvenile stage with true epiphytes, which is arguably the most crucial life history stage (Zotz et al. 2001), we included them in **EpiList 1.0**, but marked them as hemiepiphytes. This is also true for facultative hemiepiphytes. Similar to the situation in epiphytes, hemiepiphytism is frequently facultative, e.g. a large proportion of *Schefflera* species occur both as hemiepiphytes or free-standing trees (Plunkett et al. 2020).

Hemiparasitic mistletoes and epiphytes share tree crowns as habitat, but central aspects of their biology, especially in regard to plant water relations, are fundamentally different (e.g.

Poltz and Zotz 2011). Mistletoes are found in five families in the order Santalales (Vidal-Russell and Nickrent 2008); epiphytic members are missing. Thus, exclusion from **EpiList 1.0** was straightforward even when included as “epiphyte” in several of the publications on which **EpiList 1.0** is based (e.g. Fall and Drezner 2020). However, there is the interesting case of two species of root parasites that are typically found in tree crowns, where both *Gaiadendron punctatum* (Loranthaceae) (Kuijt 1963) and *Pedicularis dendrothauma* (Orobanchaceae) (Allard et al. 2005) attach themselves to roots or rhizomes of epiphytes. As noted by Kuijt (1963), one could describe both species as true epiphytes in regard to the supporting tree, but as parasites to its co-occurring epiphytes. We opted not to include the two species in **EpiList 1.0** since being non-parasitic is an essential part of our definition of an epiphyte (Fig. 1).

A third group that is sometimes lumped with epiphytes are **nomadic vines** (Fig. 1). This is probably the most contentious group. Previously dubbed “secondary” hemiepiphytes, these plants are thought to germinate on the ground, and after growing up a support “sever all connections” (Kress 1986), thus becoming epiphytic. Recent field observations on aroids (Zotz et al. 2020) are incompatible with this concept. None of the studied root-climbing Araceae became epiphytic, but all individuals maintained continuous contact with the soil via adventitious roots. Consequently, species labelled as “secondary hemiepiphyte” or “nomadic vine” in publications were not included in **EpiList 1.0** either. Unfortunately, since such plants have been frequently treated as “epiphytes” in original publications, it is likely that **EpiList 1.0** still contains some of these climbers. We also note that the issue whether plants with a truly epiphytic stage later in life do exist is unresolved: there are observations that some of the cacti included in **EpiList 1.0** as epiphytes (e.g. *Selenicereus* spp. or *Weberocereus* spp.) may actually belong in that category (Martínez-Quezada et al. 2020), although the necessary studies covering the entire ontogeny are still lacking. If corroborated, such species would also be included but marked – analogous to hemiepiphytes.

Even with the clear definitions given above, problems in the assignment of species to these categories arise 1) because of variation within a species (e.g. facultative epiphytism) and 2) because of conceptual and terminological ambiguities in the sources. The first problem is related to real biological variation, common to many biological categories (e.g. “succulence is not a binary trait”, Ogburn and Edwards 2010) and thus - by its very nature - cannot be entirely resolved. This differs from the second problem: treating, e.g., “mistletoes” or climbing “secondary hemiepiphytes” as “epiphytes” ignores fundamental biological differences in the ecology and resource acquisition and should be avoided, whereas

comparisons of different groups of structurally dependent plants or the study of their possible interaction should be highly informative and are therefore encouraged.

There is yet another terminological issue. Most sources treat epiphyte as a life form (for a discussion of “life form”, “growth form”, “habit” see Zotz 2016). However, the Brazilian Flora (2020) diverges from this practice and distinguishes “life form” and “substrate”. For example, a species like *Scaphyglottis livida* will be assigned to life form “herb” and to substrate “epiphyte”. A problem arises when researchers only report the life form as defined in the Flora do Brasil (Mota et al. 2018, da Fonseca-da-Silva et al. 2020). In these studies, *Scaphyglottis livida*, *Aechmea bromeliifolia*, and *Tillandsia streptocarpa* are reported as herbs, while no information is given on “substrate”. Both *Aechmea bromeliifolia* and *Tillandsia streptocarpa* are known as facultative epiphytes (Brazilian Flora 2020 in construction; continuously updated), but this practice makes it impossible to judge unambiguously whether these populations are epiphytic, terrestrial, or both. Similarly, possible terrestrial occurrences of the typically arboreal *Scaphyglottis livida* will be missed as well. We therefore urge researchers following the scheme of the Flora do Brasil to report both life form and substrate.

Due to the lack of data on the relative proportion of terrestrial and epiphytic individuals of a species in all but the exceptional case (e.g. Burns 2008, Burns 2010), we could not distinguish true epiphytes from facultative epiphytes, and both were included in **EpiList 1.0**, while accidental epiphytes were not. In the following, we describe our procedures with a few typical examples.

Example 1: Scrutinized data base entries

Life form information in data bases (WCSP 2019) or large treatises (Pridgeon et al. 1999-2014) was accepted unless there was convincing evidence against the categorization as epiphyte. As a concrete example, Munn-Estrada (2017) list two “epiphytic” *Agave* species. Similar records of epiphytic occurrences of a number of other congeners were found in a monograph (Gentry 2004) or in WCSP (2019) (e.g. *Agave seemannii*). Although there is no doubt that such rather spectacular-looking epiphytic growth of large *Agave* individuals may occur, they clearly represent exceptions within each *Agave* species which do not justify the inclusion of any *Agave* species in **EpiList 1.0**. We note, however, that important insights into the epiphytic life style could be gained from studying such accidental epiphytism.

Example 2: Contradictions within a publication or between publications

Munn-Estrada (2017) describes *Marcgravia stonei* as "epiphyte" in the main text, but as "shrub" in the species list. The entire family Marcgraviaceae is variously described as "epiphytic/hemiepiphytic" (Giraldo-Cañas 2018) or as lianas, climbing shrubs and treelets (Lens et al. 2005). It is currently unclear how much of these differences reflect biological plasticity or terminological inconsistency: our selection of included members of this family is mostly based on expert advice (D. Giraldo-Cañas, pers. comm.). Clearly, Marcgraviaceae is one of several families that deserve much more field work to understand their life forms as a basic attribute of their biology (a good example being the study by Massa 1996). We hope that **EpiList 1.0** will provide the traceable starting point to resolve contradictory life form assignments in this and numerous other cases in the future.

Example 3: Doubtful species, rare and unconfirmed reports

The genus *Elaeocarpus* (Elaeocarpaceae) represents c. 500 species of woody plants, mostly trees (WCVP 2020). The single existing specimen of *Elaeocarpus crenulatus* is labelled "epiphyte" (Coode 2005). Only known from the type, which was collected 70 years ago, *E. crenulatus* represents the only "epiphytic" species within this terrestrial genus. Its taxonomic status is doubtful (Coode 2005), so we opted for excluding the species from the list. Epiphytic species within otherwise terrestrial taxa are particularly interesting both ecologically and evolutionarily, but also deserve particular scrutiny. Nguyen et al. (2013) described an epiphytic *Alocasia* species (Araceae). However, the photo in their publication clearly shows feeder roots that connect *Alocasia vietnamensis* to the ground. Thus, the species is most likely a nomadic vine and was not included in **EpiList** either.

Example 4: Epiphytes vs. hemiepiphytes

Since hemiepiphytes have an epiphytic stage by definition, it is unsurprising that ambiguity exists whether a species is categorized as epiphyte or hemiepiphyte. For example, Penneys and Judd (2011) note that "general collectors often erroneously describe *Blakea* and *Topobea* species as being epiphytic, although, in fact, most are hemiepiphytes". Incorrect labelling has certainly occurred in other taxa as well. Moreover, similar to epiphytes, there are numerous terrestrial tree species that sometimes grow as facultative hemiepiphytes. With few exceptions (e.g. *Weinmannia tinctoria*, Derroire et al. 2007), facultative hemiepiphytism is even less documented than facultative epiphytism. A major problem is the need for long-term observations to establish the defining ontogenetic trajectory of a hemiepiphyte (Watts et al. 2019). As a case in point, the presence of feeder roots in *Alocasia vietnamensis* (Example 3)

does not rule out that this species is actually a hemiepiphyte, i.e. the species typically establishes epiphytically.

Species without reference

In > 99% of all cases, we supported each entry in **EpiList** with at least one reference from the literature, a freely accessible online source, or the personal observation of an expert. The 11 species without references are ferns in genera like *Grammitis* or *Pecluma* that are known to be mainly epiphytic and where morphology further suggests an epiphytic life style.

Known unknowns

For some genera (e.g., *Clusia*, *Schefflera*) in which a large proportion of all species regularly occur as an epiphyte or hemiepiphyte we certainly underestimate the number of species in **EpiList 1.0**. However, without documentation for individual species there was no objective way to include additional species. Another known unknown refers to taxa in underexplored geographical regions. For example, we currently know of 35 epiphytic *Impatiens* species, mostly from Africa. More than half of these species (19) have been described in the last 20 years, and more botanical explorations should lead to the discovery of additional species. Similarly, over 80 new species of epiphytic grammitid ferns (Polypodiaceae), especially in the genus *Oreogrammitis*, are in the process of being described (B. Parris, pers. comm.).

II.B.3.b. Instrumentation: N/A

II.B.3.c. Taxonomy and systematics:

EpiList 1.0 encompasses 31,311 epiphytic and hemiepiphytic species (28227 angiosperms, 2 gymnosperms, 2942 Polypodiopsida, 142 Lycopodiopsida). Table 1 summarizes species numbers per genus and per family for seed plants and Table 2 for Polypodiopsida and Lycopodiopsida. Although epiphytes are found in 79 families of vascular plants in 918 genera, their taxonomic distribution is very uneven. Among the 59 families of seed plants, Orchidaceae clearly stands out by accounting for almost 75% of the total epiphyte species count (21,169 of 28,227 species) and 63% of all genera with epiphytes (559 of 892 genera). This dominance would look even more impressive if we excluded all families and genera that have mostly or exclusively hemiepiphytic members. These sum up to 14 families and 42

genera (Table 1). Among Polypodiopsida, epiphytes are particularly common in Polypodiaceae with almost 50% of the total epiphyte species count (1450 of 2942 species) and 55% of all genera (68 of 123 genera). Hemiepiphytism in ferns has only recently been discovered (Nitta and Epps 2009), and subsequently confirmed cases of hemiepiphytic ferns are still confined to a few lineages. Many of the ferns labelled as hemiepiphytic in **EpiList 1.0** are presented as hypotheses to be confirmed or rejected as new data become available.

With 31,311 species, **EpiList 1.0** documents a substantial increase in the number of epiphytes and hemiepiphytes compared to previous estimates of 23,456 species (Kress 1986) and 27,614 species (Zotz 2013). Compared to the last global revision (Zotz 2013), there is also an, albeit somewhat smaller, increase in the number of genera (918 vs. 913, + 1%) and families (79 vs. 73, + 7%).

A synopsis of the taxonomic distribution of vascular epiphytes within families and genera is given separately for seed plants (Table 1), and ferns and lycopods (Table 2). Species listed in wfo are either “accepted”, “synonyms” or in revision, i.e. labelled “ambiguous”. A considerable number of species in **EpiList 1.0** are in the last category. Therefore, Table 1 details separate numbers for “accepted” and all (“accepted” + “ambiguous”) epiphytic species, as well as for the total number of “accepted” and the total number of all (“accepted” + “ambiguous”) species in each family and genus. It also gives the proportion of epiphytes among accepted and all species per genus and family (in %). These proportions are highly correlated ($r = 0.96$). Some species in **EpiList 1.0** belong to genera that are not found in wfo or treated as synonymous, only accepted in wesp or powo, e.g. *Trachoma* spp. in Orchidaceae or *Wittmackia* spp. in Bromeliaceae. These obviously unresolved cases could not be matched with any species in wfo, and we did not calculate relative proportions of epiphytic members.

II.B.3.d. Permit history: N/A

II.B.3.e. Legal/organizational requirements: N/A

II.B. 4. Project personnel: See I.C.1.

Class III. Data set status and accessibility

III.A. Status

III.A.1. Latest update: 2020

III.A.2. Latest archive date: 2020

III.A.3. Metadata status: Last updated June 2020, this document.

III.A.4. Data verification:

III.B. Accessibility

III.B.1. Storage location and medium: The primary data are stored on a server at the University of Göttingen, Germany. The published data are available online through the current publication as a csv.file.

III.B.2. Contact person(s):

Please direct any feedback and questions to:

Gerhard Zotz, Functional Ecology of Plants, Institute of Biology and Environmental Sciences, Ammerländer Heerstr. 114-118, D-26129 Oldenburg, Germany, email: gerhard.zotz@uol.de

III.B.3. Copyright restrictions: None

III.B.4. Proprietary restrictions: We release these data into the public domain under a Creative Commons Zero license waiver (<https://creativecommons.org/share-your-work/public-domain/cc0/>). When you use the data in your publication, we request that you cite this data paper. Should **EpiList 1.0** be a major part of the data analyzed in your study, you should consider inviting the **EpiList 1.0** core team (see Originators in the Overall project description) as collaborators. If you plan to use the **EpiList 1.0** dataset, we request that you contact the **EpiList 1.0** core team to learn whether updates of the dataset are available, and whether similar analyses are already ongoing.

III.B.4.a. Release date: 13 March 2021

III.B.4.b. Citation: Zotz, G., P. Weigelt, M. Kessler, H. Kreft, and A. Taylor. 2021. EpiList 1.0: a global checklist of vascular epiphytes. Ecology e03326.

<https://doi.org/10.1002/ecy.3326>

III.B.4.c. Disclaimer(s): none

III.B.5. Costs: none

Class IV. Data structural descriptors

IV.A. Data set file

IV.A.1. Identity: EpiList1.0.csv

EpiList1.0 (31,311 entries standardized with wfo and WorldFerns. Up to three references that report epiphytic growth are given. For full list of citations see **Class V.F. Publications and results**)

IV.A.2. Size: EpiList1.0.csv (3.5 MB)

IV.A.3. Format and storage mode: Data table available as comma-separated values (*.csv).

IV.A.4. Header information: See column descriptions in **IV.B.2.**

IV.A.5. Alphanumeric attributes: N/A

IV.A.6. Special characters/fields: N/A

IV.A.7. Authentication procedures: N/A

IV.B. Variable information:

IV.B.1. Variable identity: N/A

IV.B.2. Variable definition:

The following variables are found in EpiList1.0.csv:

Family = plant family

Species = species names following the Standardization reference (= tax source)

Genus = genus name of each species

Epithet = species epithet of each species

Author(s) = Author(s) of species name

Status = Status of species name in standardization reference (= tax source), either “accepted” or “ambiguous” when tax source is wfo or PPG-I, “empty” when name is not found in these sources and standardization was done with other sources

Usage ID = Usage ID of wfo

Tax source = reference used for taxonomic standardization (“Hassler” = WorldFerns Version 8.30, “powo” = plants of the world online; “reference I = Reference I; “wcsp” = world checklist of selected plant families; “wfo” = World Flora Online

Reference I – III = reference stating that the species is an epiphyte / hemiepiphyte

Hemi = “H” indicates that the species is a hemiepiphyte, “H?” if doubtful

IV.B.3. Units of measurement: N/A

IV.B.4. Data type

IV.B.4.a. Storage type: csv.files

IV.B.4.b. List and definition of variable codes: See Definitions in “Sampling Design and Research Methods.”

IV.B.4.c. Range for numeric values: N/A

IV.B.4.d. Missing value codes: N/A

IV.B.4.e. Precision: N/A

IV.B.5. Data format

IV.B.5.a. Fixed, variable length: N/A

IV.B.5.b. Columns: See IV.B.2.

IV.B.5.c. Optional number of decimal places: N/A

IV.C. Data anomalies: N/A

Class V. Supplemental descriptors

V.A. Data acquisition

V.A. 1. Data forms or acquisition methods: available online through the current publication.

V.A. 2. Location of completed data forms: N/A

V.A. 3. Data entry verification procedures: N/A

V.B. Quality assurance/quality control procedures: N/A

V.C. Related materials: N/A

V.D. Computer programs and data-processing algorithms: N/A

V.E. Archiving

V.E.1. Archival procedures: available online through the current publication.

V.E.2. Redundant archival sites: N/A

V.F. Publications and results: The following list of sources represents the web resources and print publications reporting epiphytic growth of each species in EpiList1.0.csv. We used information from c. 7000 sources, but only a fraction of these is actually shown in the following list. Priority was given to large online data bases or floras, followed by peer-reviewed publications, but annotated species lists and other internet resources were also used. Our goal was to have at least one citation for each species in **EpiList1.0**.

A world of Pinguicula. 2010.

http://www.pinguicula.org/pages/pages_principales/content.html.

A.G.T. 1918. A contribution to the phytogeography and flora of the Arfak Mountains, etc. *Journal of Ecology* 6:157-159.

Acebey, A. R., T. Krömer, and M. Kessler. 2017. Species richness and vertical distribution of ferns and lycophytes along an elevational gradient in Los Tuxtlas, Veracruz, Mexico. *Flora* 235:83-91.

Acevedo-Rodríguez, P., and M. T. Strong. 2005. Monocotyledons and gymnosperms of Puerto Rico and the Virgin Islands. Department of Botany, National Museum of Natural History, Smithsonian Institution, Washington, DC.

Achoundong, G. 1996. Mountain forests in the Cameroon: Flora of the hills of Yaounde. *Bois et Forêts des Tropiques* 247:37-52.

Adams, C. D. 1992. Eleven new species, a new variety, and a new varietal combination in the fern genera *Asplenium* and *Diplazium* in Central America. *Novon* 2:290-298.

- African_Plant_Database. 2012. <http://www.village.ch/musinfo/bd/cjb/africa/details.php?langue=en&id=48569>.
- Aguiar, S., A. Herrero, and L. G. Quintanilla. 2006. Confirmación citológica de la presencia de *Hymenophyllum wilsonii* en España. *Lazaroa* 27:129.
- Aguirre, A., R. Guevara, M. García, and J. C. López. 2010. Fate of epiphytes on phorophytes with different architectural characteristics along the perturbation gradient of *Sabal mexicana* forests in Veracruz, Mexico. *Journal of Vegetation Science* 21:6-15.
- Aguirre-Leon, E. 1992. Vascular epiphytes of Mexico: a preliminary inventory. *Selbyana* 13:72-76.
- Airy Shaw, H. K. 1935. Studies in the Ericales: I. New and less-known species of Agapetes. *Bulletin of Miscellaneous Information (Royal Gardens, Kew)* 1935:24-53.
- Airy Shaw, H. K. 1968. Studies in the Ericales: XV: New or noteworthy Agapetes from Assam and Burma. *Kew Bulletin* 21:471-476.
- Alemayehu, T. 2006. Diversity and ecology of vascular epiphytes in Harennia afro-montane forest, Bale, Ethiopia. MSc-Arbeit. Addis Ababa University, Addis Ababa.
- Almeda, F. 1990. New species and new combinations in *Blakea* and *Topobea* (Melastomataceae), with an historical perspective on generic limits in the tribe Blakeeae. *Proceedings of the California Academy of Sciences* 46:299-326.
- Almeda, F. 2000. The hexandrous species of *Topobea* (Melastomataceae). *Proceedings of the California Academy of Sciences* 52:97-109.
- Almeda, F., and D. E. Breedlove. 1992. A new *Vaccinium* (Ericaceae, Vaccinieae) from Guerrero, Mexico. *Brittonia* 44:50-53.
- Almeida, T. E., A. Salino, J.-Y. Dubuisson, and S. Hennequin. 2017. *Adetogramma* (Polypodiaceae), a new monotypic fern genus segregated from *Polypodium*. *PhytoKeys*:109-131.
- Alvaro Vindas, E. 2003. Plantas comunes del Parque Nacional Chirripó. 2nd edition. Instituto Nacional de Biodiversidad, INBio, Santo Domingo, Heredia.
- Alzate, F., F. Cardona, and R. Callejas. 2001. Diversidad y composición de epífitas vasculares en robledales de Antioquia (Colombia). *Actualidades Biológicas Medellín* 23:25-31.
- Amoroso, V. B., F. P. Coritico, and P. W. Fritsch. 2016. Species richness and conservation status of ferns and lycophytes in Mt. Hamiguitan Range Wildlife Sanctuary, Davao Oriental, Philippines. *Philippine Journal of Science* 145:127-137.
- Amoroso, V. B., F. P. Coritico, and P. W. Fritsch. 2020. *Actinostachys minuta*, a new species of grass fern from Mindanao, Philippines. *PhytoKeys* 151:59-66.
- Anjum, S., S. Pant, H. C. Pande, and T. Rinchen. 2014. Ecological observations on the fern flora of Thannamandi District Rajouri, Jammu & Kashmir, India. *Indian Forester* 140:1235-1244.
- Annaselvam, J., and N. Parthasarathy. 2001. Diversity and distribution of herbaceous vascular epiphytes in a tropical evergreen forest at Varagalaia, Western Ghats, India. *Biodiversity and Conservation* 10:317-329.
- Anonymous. Group D: Epiphytes (other than ferns). *Mangrove guidebook for Southeast Asia*.
- Arévalo, R., and J. Betancur. 2004. Diversidad de epífitas vasculares en cuatro bosques del sector suroriental de la Serranía de Chiribiquete, Guayana colombiana. *Caldasia* 26:359-380.
- Argent, G. 1988. *Vireya* taxonomy in field and laboratory. In Clyde Smith, J., editor. *Proceedings of the Fourth International Rhododendron Conference*, Wollongong, New South Wales, Australia.
- Argent, G. C. G. 1982. *Notulae et novitates Muluenses*. New taxa and combinations and comments on Ericaceae. *Botanical Journal of the Linnean Society* 85:2-17.
- Argent, G. C. G. 2002. New taxa and combinations in the genus *Diplycosia* (Ericaceae) of Borneo and Peninsular Malaysia. *Gardens' Bulletin, Singapore* 54:217-238.

- Argent, G. C. G. 2014. A contribution to the study of the genus *Diplycosia* (Ericaceae) in Sulawesi, Indonesia. *Edinburgh Journal of Botany* 71:83-115.
- Australian_National_Botanic_Gardens. <http://www.anbg.gov.au/>
- Australian_Orchid_Council_Inc. <http://orchidsaustralia.com.au/>
- Australian_tropical_rainforest-orchids. 2010. <http://www.anbg.gov.au/cpbr/cd-keys/orfk/index.html>.
- Auvray, G., and M. F. Newman. 2010. A revision of *Cautleya* (Zingiberaceae) *Edinburgh Journal of Botany* 67:451-465.
- Bach, T. T., J.-H. Kim, D.-K. Kim, J. Lee, B. T. Ha, N. S. Juhonewe, and M. Rodda. 2011. *Hoya ignorata* (Apocynaceae, Asclepiadoideae): An overlooked species widely distributed across Asia. *Novon* 21:508-514.
- Badré, F., and M. Hoff. 1995. Les ptéridophytes des Iles Wallis et Futuna (Pacifique Sud) écologie et répartition. *Feddes Repertorium* 106:271-290.
- Baksh-Comeau, Y. S., Maharaj, S.S., Adams, C.D., Harris, S.A., Filer, D.L., Hawthorne, W.D. 2016. An annotated checklist of the vascular plants of Trinidad and Tobago with analysis of vegetation types and botanical 'hotspots'. *Phytotaxa* 250:1-431.
- Bamps, P. 1974. Flore d'Afrique Centrale: Spermatophytes. Araliaceae. Jardin Botanique National de Belgique Meise.
- Banik, D., and M. Sanjappa. 2007. Three new species of *Agapetes* (Ericaceae) from India. *Edinburgh Journal of Botany* 64:335-342.
- Banik, D., and M. Sanjappa. 2008. Notes on identity of some Indian species of *Agapetes* (Ericaceae). *Nordic Journal of Botany* 26:4-9.
- Barahona Carvajal, M. E. 1977. Estudio morfológico comparativo de las inflorescencias de dos especies de Araceae: *Anthurium denudatum* Engler y *Philodendron radiatum* Schott. *Revista de Biología Tropical* 25:301-334.
- Baranow, P. 2015. Taxonomic notes on *Sobralia* section *Abbreviatae* (Orchidaceae) in Colombia, with description of a new species. *Plant Systematics and Evolution* 301:41-60.
- Barbhuiya, H. A., B. K. Dutta, A. K. Das, and A. K. Baishya. 2014. The family Rubiaceae in southern Assam with special reference to endemic and rediscovered plant taxa. *Journal of Threatened Taxa* 6:5649-5659.
- Barcelona, J., and M. Price. 1999. New Philippine *Goniophlebium* (Polypodiaceae: Pteridophyta). *Fern Gazette* 15:261-264.
- Barkworth, M. E., and R. P. Adams. 1993. *Flora of North America: Volume 2: Pteridophytes and Gymnosperms*. Oxford University Press.
- Barringer, K. 2004. A revision of *Gibsoniothamnus* L. O. Williams (Schlegeliaceae). *Brittonia* 56:213-237.
- Barroso, G. M. 1957. *Araceae novae*. *Arquivos do Jardim Botânico do Rio de Janeiro* 15:96-97.
- Barthlott, W. pers. comm.
- Barthlott, W., V. Schmit-Neuerburg, J. Nieder, and S. Engwald. 2001. Diversity and abundance of vascular epiphytes: a comparison of secondary vegetation and primary montane rain forest in the Venezuelan Andes. *Plant Ecology* 152:145-156.
- Bellefroid, E., S. K. Rambe, O. Leroux, and R. L. L. Viane. 2010. The base number of 'loxoscaphoid' *Asplenium* species and its implication for cytoevolution in Aspleniaceae. *Annals of Botany* 106:157-171.
- Benavides, A. M., A. J. Duque, J. F. Duivenvoorden, G. A. Vasco, and R. Callejas. 2005. A first quantitative census of vascular epiphytes in rain forests of Colombian Amazonia. *Biodiversity and Conservation* 14:739-758.

- Benavides, A. M., A. Vasco, A. J. Duque, and J. F. Duivenvoorden. 2011. Association of vascular epiphytes with landscape units and phorophytes in humid lowland forests of Colombian Amazonia. *Journal of Tropical Ecology* 27:223-237.
- Benavides, A. M., J. H. D. Wolf, and J. F. Duivenvoorden. 2006. Recovery and succession of epiphytes in upper Amazonian fallows. *Journal of Tropical Ecology* 22:705-717.
- Benl, G. 1991. Family Aspleniaceae. *Acta Botanica Barcinonensia* 40:3-104.
- Benzing, D. H. 1990. Vascular epiphytes. General biology and related biota. Cambridge University Press, Cambridge.
- Berg, C. C. 1990. New species of *Coussapoa* and *Pourouma* (Cecropiaceae). *Brittonia* 42:59-65.
- Berg, C. C., R. W. A. P. Akkermans, and E. C. H. v. Heusden. 1990. Cecropiaceae: *Coussapoa* and *Pourouma*, with an introduction to the family. *Flora Neotropica* 51:1-208.
- Bergamini Scheer, M. 2010. Ambientes altomontanos no Paraná: Florística vascular, estrutura arbórea, relações pedológicas e datações por ¹⁴C. PhD. Universidade Federal do Paraná, Curitiba.
- Bernacci, L. C. 2009 onwards. Neotropical Myrsinaceae. In W. Milliken, B. Klitgård, and A. Barakat, editors. Neotropikey - Interactive key and information resources for flowering plants of the Neotropics. www.kew.org/neotropikey
- Bernal, R., S. R. Gradstein, and M. Celis. 2019. Catálogo de plantas y líquenes de Colombia. Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Bogotá. <http://catalogoplantasdecolombia.unal.edu.co>.
- Bhaskar, V., and B. Razi. 1982. Two more new species of *Impatiens* L. from South India. *Journal of the Bombay Natural History Society* 79:382-384.
- Bidin, A. A., and R. Jaman. 1999. The pteridophytes of Tawau Hills Park, Sabah. *ASEAN Review of Biodiversity and Environmental Conservation (ARBEC)* 1:1-11.
- Biedinger, N., and E. Fischer. 1996. Epiphytic vegetation and ecology in central African forests (Rwanda, Zaire). *Ecotropica* 2:121-142.
- Binns, D. L., and S. F. o. N. S. W. R. Division. 1995. Flora survey, Dorrigo three-year Environmental Impact Statement Area, Northern Region, New South Wales. Research Division, State Forests of New South Wales, Sydney.
- Bishop, L. E. 1974. Revision of genus *Adenophorus* (Grammitidaceae). *Brittonia* 26:217-240.
- Bishop, L. E. 1978. Revision of genus *Cochlidium* (Grammitidaceae). *American Fern Journal* 68:76-94.
- Bishop, L. E. 1989. New species of *Ceradenia* subg. *Ceradenia*. *American Fern Journal* 79:14-25.
- Bishop, L. E., and A. R. Smith. 1992. Revision of the fern genus *Enterosora* (Grammitidaceae) in the New World. *Systematic Botany* 17:345-362.
- Blum, C. T., C. V. Roderjan, and F. Galvao. 2011. Composição florística e distribuição altitudinal de epífitas vasculares da Floresta Ombrófila Densa na Serra da Prata, Morretes, Paraná, Brasil. *Biota Neotropica* 11:141-159.
- Bodegom, S., and J. F. Veldkamp. 2001. Revision of the pseudo-stipular species of *Medinilla* (Melastomataceae). *Blumea* 46:527-567.
- Boelter, C. R., C. E. Zartman, and C. R. Fonseca. 2011. Exotic tree monocultures play a limited role in the conservation of Atlantic Forest epiphytes. *Biodiversity and Conservation* 20:1255-1272.
- Bongers, F. 2001. *Nouragues: Dynamics and plant-animal interactions in a neotropical rainforest*. Kluwer Academic Publishers, Dordrecht.
- Bosman, M. T. M. 1991. A monograph of the fern genus *Microsorium* (Polypodiaceae): including an attempt towards a reconstruction of the phylogenetic history of the microsorioids. *Leiden Botanical Series* 14: 1-161.

- Bosser, J. 1971. Contribution a l'étude des Orchidaceae de Madagascar. XVII. *Adansonia* sér. 2 11:519-543.
- Box, E. O., C.-H. Chou, and K. Fujiwara. 1998. Richness, climatic position, and biogeographic potential of East Asian laurophyll forests, with particular reference to examples from Taiwan Bulletin, Institute of Environmental Science and Technology, Yokohama National University 24:61-95.
- Boyce, P. pers. comm.
- Brass, L. J. 1953. Ferns of a New Guinea gully. *American Fern Journal* 43:150-158.
- Brazilian Flora 2020 in construction. continuously updated. - available in <http://floradobrasil.jbrj.gov.br/>, Rio de Janeiro Botanical Garden.
- Breier, T. B. 2005. O epifitismo vascular em florestas do sudeste do Brasil. Ph.D. Universidade Estadual de Campinas, Campinas.
- Bremer, B. 1989. The genus *Argostemma* (Rubiaceae-Argostemmataceae) in Borneo. *Annals of the Missouri Botanical Garden* 76:7-49.
- Bridgewater, S. G. M., D. J. Harris, C. Whiteford, A. K. Monro, M. G. Penn, D. A. Sutton, B. Sayer, B. Adams, M. J. Balick, D. H. Atha, J. Solomon, and B. K. Holst. 2006. A preliminary checklist of the vascular plants of the Chiquibul Forest, Belize. *Edinburgh Journal of Botany* 63:269-321.
- Brownlie, G. 1960. Studies on Pacific ferns, Part I: Nomenclature changes and distributions of some species of *Hymenophyllum*, *Arthropteris*, *Microlepia*, *Oleandra*, and *Adiantum*. *Pacific Science* 14:242-245.
- Brownlie, G. 1969. Fasc. 3, Ptéridophytes. Muséum National d'Histoire Naturelle, Paris, Paris.
- Brownlie, G., and W. Philipson. 1971. Pteridophyta of the Southern Cook Group. *Pacific Science* 25:502-511.
- Brownsey, P., and L. Perrie. 2014. Flora of New Zealand: ferns and lycophytes. Fascicle 1. Polypodiaceae. Manaaki Whenua Press, Lincoln, NZ.
- Brownsey, P., and L. Perrie. 2016. Flora of New Zealand: ferns and lycophytes. Fascicle 16. Hymenophyllaceae. Manaaki Whenua Press, Lincoln, NZ.
- Burns, K. C. 2007. Network properties of an epiphyte metacommunity. *Journal of Ecology* 95:1142-1151.
- Burns, K. C. 2008. Meta-community structure of vascular epiphytes in a temperate rainforest. *Botany* 86:1252-1259.
- Burrows, J., Burrows, S. 2003. Figs of Southern and South-Central Africa. Umdaus Press, Hatfield.
- Burt, B. L. 1948. Studies in the Ericales: IX. The taxonomic position of *Wittsteinia*. *Kew Bulletin* 3:493-495.
- Bussmann, R. W. 2001. Epiphyte diversity in a tropical Andean forest - Reserva Biológica San Francisco, Zamora-Chinchipe, Ecuador. *Ecotropica* 7:43-60.
- Buzatto, C. R., K. L. Davies, R. B. Singer, R. Pires dos Santos, and C. van den Berg. 2012. A comparative survey of floral characters in *Capanemia* Barb. Rodr. (Orchidaceae: Oncidiinae). *Annals of Botany* 109:135-144.
- C.V. Starr Virtual Herbarium. <http://sweetgum.nybg.org/science/vh/>.
- Cadet, T. 1977. La végétation de l'Île de la Réunion. Etude phytocéologique et phytosociologique. PhD Thesis. Université d'Aix-Marseille, St. Denis, La Réunion.
- Callmander, M. W. pers. comm.
- Callmander, M. W., P. P. Lowry, II, F. Forest, D. S. Devey, H. Beentje, and S. Buerki. 2012. *Benstonea* Callm. & Buerki (Pandanaceae): characterization, circumscription, and distribution of a new genus of screw-pines, with a synopsis of accepted species. *Candollea* 67:323-345.

- Callmander, M. W., T. J. Booth, H. Beentje, and S. Buerki. 2013. Update on the systematics of *Benstonea* (Pandanaeae): When a visionary taxonomist foresees phylogenetic relationships. *Phytotaxa* 112:57-60.
- Camelo, M. C., M. A. N. Coelho, and L. G. Temponi. 2018. Two new species of *Anthurium* (Araceae) from Caparao National Park, southeastern Brazil. *Phytotaxa* 373:121-130.
- Cardozo, A. P., L. Godinho Temponi, I. Moreira de Andrade, S. J. Mayo, and E. de Camargo Smidt. 2014. A morphometric and taxonomic study of *Anthurium augustinum* complex (Araceae), endemic to the Brazilian Atlantic Forest. *Feddes Repertorium* 125:43-58.
- Carlsen, M. M., and T. B. Croat. 2004. New species of *Anthurium* Sect. *Semaephyllium* (Araceae) from Central and South America. *Novon* 14:401-412.
- Carlsen, M. M., and T. B. Croat. 2007. Taxonomic revision of *Anthurium* section *Semaephyllium* Schott (Araceae). *Harvard Papers in Botany* 12:173-234.
- Cascante, A., and A. Estrada. 1999. Lista con anotaciones de la flora vascular de la Zona Protectora El Rodeo, Costa Rica (un Bosque Húmedo Premontano del Valle Central). *Brenesia* 51:1-44.
- Cascante-Marín, A., and A. Estrada-Chavarría. 2012. Las plantas vasculares de El Rodeo, Costa Rica. *Brenesia* 77:71-128.
- Casper, S. J. 1987. On *Pinguicula lignicola*, an epiphytic heterophyllic member of the Lentibulariaceae in Cuba. *Plant Systematics and Evolution* 155:349-354.
- Castañeda G., G. 2001. Aves asociadas a plantas epifitas de un bosque nublado en la reserva de bosque integral Otonga, noroccidente de Ecuador. Pages 327-334 in J. Nieder and W. Barthlott, editors. *Epiphytes and canopy fauna of the Otonga rain forest (Ecuador). Results of the Bonn - Quito epiphyte project, funded by the Volkswagen Foundation.* Books on Demand, Bonn.
- Catarino, L., Martins, E.S., Basto, M.F., Diniz, M.A., 2008. An annotated checklist of the vascular flora of Guinea-Bissau (West Africa). *Blumea* 53:1-222.
- Catling, P. M., and L. P. Lefkovitch. 1989. Association of vascular epiphytes in a Guatemalan cloud forest. *Biotropica* 21:35-40.
- Ceja-Romero, J., A. Mendoza-Ruiz, A. R. López-Ferrari, A. Espejo-Serna, B. Perez-Garcia, and J. Garcia-Cruz. 2010. The vascular epiphytic flora of the State of Hidalgo, Mexico diversity and distribution. *Acta Botánica Mexicana* 93:1-39.
- Cerneaux, L., T. Jaffré, M. Lescot, P. Morat, and J. M. Veillon. 1990. Flore de Nouvelle Calédonie. Familles revues (Nomenclature et écologie). *Sciences de la Vie Botanique* 4, Orstom, Nouméa.
- Cetzal-Ix, W., E. Noguera-Savelli, M. Martinez-Ico, and N. Ramirez-Marcial. 2013. Diversity of ferns and lycophytes in parts of semievergreen forest in southern Tabasco, Mexico. *Botanical Sciences* 91:261-271.
- Chaerle, P., and R. Viane. 2007. A new species of *Asplenium* (Asplenaceae, Pteridophyta) from Ethiopia. *Novon* 17:148-153.
- Champluvier, D., and B. Senterre. 2010. New taxa in *Crossandrella*, *Dischistocalyx* and *Ascotheca* (Acanthaceae) from Equatorial Guinea. *Plant Ecology and Evolution* 143:181-190.
- Chandra, S. 1983. Morphology of *Dictymia brownii* (Wikstr.) Copel. (Polypodiaceae). *Australian Journal of Botany* 31:557-563.
- Chang, M. 2005. The vines and epiphytes of a warm-temperate evergreen broad-leaved forest in Chungtzekuan area. National University of Tainan.
- Chao, C.-T., Y.-H. Tseng, and H.-Y. Tzeng. 2013. *Heteropolygonatum altelobatum* (Asparagaceae), comb. nova. *Annales Botanici Fennici* 50:91-94.
- Chapman, V. J. 1976. Mangrove vegetation. Vaduz, Cramer. 447 pp.
- Cheek, M. 2004. *Impatiens letouzeyi*. IUCN 2011. IUCN Red List of Threatened Species. Version 2011.2.

- Cheek, M. 2009. *Mussaenda epiphytica* sp nov (Rubiaceae), an epiphytic shrub from cloud forest of the Bakossi Mts, western Cameroon. *Nordic Journal of Botany* 27:456-459.
- Cheek, M., and E. Fischer. 1999. A tuberous and epiphytic new species of *Impatiens* (Balsaminaceae) from southwest Cameroon. *Kew Bulletin* 54:471-475.
- Cheek, M., and L. Csiba. 2002. A new epiphytic species of *Impatiens* (Balsaminaceae) from western Cameroon. *Kew Bulletin* 57:669-674.
- Chen, C. W. pers. comm.
- Chen, C. W., A. Ebihara, W. L. Chiou, and C. W. Li. 2014. *Haplopteris yakushimensis* (Pteridaceae, Vittarioideae), a new species from Yakushima Island, Japan. *Phytotaxa* 156:229-234.
- Chen, C. W., J. H. Nitta, M. Fanerii, T. Y. A. Yang, F. Pitisopa, C. W. Li, and W. L. Chiou. 2015. *Antrophyum solomonense* (Pteridaceae), a new species from the Solomon Islands, and its systematic position based on phylogenetic analysis. *Systematic Botany* 40:645-651.
- Chen, C. W., Y. M. Huang, L. Y. Kuo, N. Quoc Dat, L. Hong Truong, J. R. Callado, D. R. Farrar, and W. L. Chiou. 2013b. trnL-F is a powerful marker for DNA identification of field vittarioid gametophytes (Pteridaceae). *Annals of Botany* 111:663-673.
- Chen, C. W., Y. M. Huang, L. Y. Kuo, Y. H. Chang, Y. C. Liu, and W. L. Chiou. 2013a. A new vittarioid fern species, *Haplopteris heterophylla* (Pteridaceae). *Systematic Botany* 38:901-909.
- Chen, C., S. Lindsay, C. Mynssen, L. Kuo, and V. Amoroso. 2018. A new *Haplopteris* species from the Philippines and clarification of the status of *H. amboinensis* in China and Indochina. *Gardens Bulletin (Singapore)* 70:177-190.
- Chettri, R., and B. Rai. 1996. A survey on the distribution of vascular epiphytes of East Sikkim forest with reference to host specificity. *Advances in Plant Sciences* 9:199-206.
- Ching, R. C. 1978. *Tricholepidium* Ching, a new genus of the Polypodiaceae in Asia. *Acta Phytotaxonomica et Geobotanica* 29:41-46.
- Chinnock, R. J. 1975. The New Zealand species of *Tmesipteris* (Psilotaceae). *New Zealand Journal of Botany* 13:743-768.
- Chomicki, G., and S. S. Renner. 2016. Evolutionary relationships and biogeography of the ant-epiphytic genus *Squamellaria* (Rubiaceae: Psychotrieae) and their taxonomic Implications. *PLoS ONE* 11:e0151317-e0151317.
- Chong, K. Y., H. T. W. Tan, and R. T. Corlett. 2009. A checklist of the total vascular plant flora of Singapore: Native, naturalised and cultivated species. Raffles Museum of Biodiversity Research, National University of Singapore, Singapore.
- Christenhusz, M. J. 2009. Index Pteridophytorum Guadalupensium or a revised checklist to the ferns and club mosses of Guadeloupe (French West Indies). *Botanical Journal of the Linnean Society* 161:213-277.
- Christenhusz, M. J., P. B. Schwartsburd, and P. H. Labiak. 2009. *Hymenophyllum filmenofilicum* (Hymenophyllaceae, Pteridophyta): a new epipetric filmy fern from Paraná, southern Brazil. *Kew Bulletin* 64:175-178.
- Christensen, C. 1912. XIX. - On the ferns of the Seychelles and the Aldabra Group. *Transactions of the Linnean Society of London -Second Series Botany VII*: 409-424.
- Christensen, C. 1937. Botanical results of the Archbold Expedition No. 8: New and noteworthy Papuan ferns. *Brittonia* 2:265-317.
- Christensen, C. 1943. A revision of the Pteridophyta of Samoa. *Bernice P. Bishop Museum Bulletin* 177:1-138.
- Clark, J. L., and J. R. Clark. 2008. *Drymonia collegarum* (Gesneriaceae), a new species from Ecuador. *Selbyana* 29:152-156.

- Clark, J. L., and L. E. Skog. 2008. Novae Gesneriaceae neotropicarum xv: *Kohleria hypertrichosa*, a new species of Gesneriaceae from northwestern Ecuador. *Journal of the Botanical Research Institute of Texas* 2:19-23.
- Clark, J. L., P. S. Herendeen, L. E. Skog, and E. A. Zimmer. 2006. Phylogenetic relationships and generic boundaries in the Episcieae (Gesneriaceae) inferred from nuclear, chloroplast, and morphological data. *Taxon* 55:313-336.
- Clarkson, B. D. 1985. The vegetation of the Kaitake Range Egmont National Park, New Zealand. *New Zealand Journal of Botany* 23:15-31.
- Clausen, R. T. 1954. Ophioglossaceae of the Hawaiian Islands. *American Journal of Botany* 41:493-500.
- Clausing, G. 2000. Revision of *Pachycentria* (Melastomataceae). *Blumea* 45:341-375.
- Co's_digital_flora_of_the_Philippines. 2019. <https://www.philippineplants.org/Families/Ericaceae.html>.
- Coelho, M. A. N., and E. L. M. Catharino. 2008. Duas espécies novas de *Anthurium* (Araceae) endêmicas do litoral de São Paulo, Brasil *Rodriguezia* 59:829-833.
- Conlon, T. 2012. *Dimorphanthera* (F.Muell. Ex Drude) F.Muell. – ericaceous gems from New Guinea at the Royal Botanic Garden Edinburgh. *Sibbaldia* 10:133.
- Copeland, E. 1929a. Papuan pteridophytes collected for the Arnold Arboretum by LJ Brass. *Journal of the Arnold Arboretum* 10:174-182.
- Copeland, E. B. 1908. New genera and species of Bornean ferns. *The Philippine Journal of Science* 3:343-352.
- Copeland, E. B. 1910. The ferns of Mount Apo. *Leaflets of Philippine Botany* 3:791-851.
- Copeland, E. B. 1929b. Ferns of Fiji. *Bernice P. Bishop Museum Bulletin* 59:1-105.
- Copeland, E. B. 1931. Pteridophytes collected for the Arnold Arboretum on Vanikoro, Santa Cruz Islands, by SF Kajewski. *Journal of the Arnold Arboretum* 12:46-49.
- Copeland, E. B. 1941. Notes on Hymenophyllaceae. *Philippine Journal of Science* 73:457-469.
- Cornejo, X., and C. H. Dodson. 2011. *Sobralia rhizophorae*: A new species of Orchidaceae from the mangroves in northwestern Ecuador. *Harvard Papers in Botany* 16:53-56.
- Cornell_University_Vascular_Plant_Herbarium. 2019. <http://phytoimages.siu.edu/>.
- Correa A., M. D., C. Galdames, and M. S. Stapf de. 2004. Catálogo de las plantas vasculares de Panamá. Universidad de Panamá & Instituto Smithsonian de Investigaciones Tropicales, Panama City.
- Corso, M., and M. Fontanier. 2008. A photographic field guide to epiphytic vascular plants on Dominica Dominica Study Abroad 2008 Texas A&M University.
- Couto Rodrigues, D., H. Machado Dias, M. C. Alvarez Pereira, C. Nicoletti de Fraga, and J. E. Macedo Pezzopane. 2016. Vascular epiphytes on *Pseudobombax* (Malvaceae) in rocky outcrops (inselbergs) in Brazilian Atlantic Rainforest: basis for conservation of a threatened ecosystem. *Rodriguezia* 67:583-601.
- Crane, E. H. 1997. A revised circumscription of the genera of the fern family Vittariaceae. *Systematic Botany* 22:509-517.
- Croat, T. B. 1978. *Flora of Barro Colorado Island*. Stanford University Press, Stanford.
- Croat, T. B. 1988. Ecology and life forms of Araceae. *Aroideana* 11:4-55.
- Croat, T. B. 1998. *Anthurium*. Retrieved from <http://www.aroid.org/genera/anthurium/anthintro.php> checked on 17 May 2019.
- Croat, T. B. 2017. New species of *Anthurium* sect. *Calomystrium* (Araceae) from Mexico and Panama. *Aroideana* 40:97-116.
- Croat, T. B. 2019. *Anthurium rosselianum* (Araceae), a new species of *Anthurium* section *Belolonchium* from Colombia. *Phytotaxa* 391:101-107.
- Croat, T. B. pers. comm.

- Croat, T. B., and A. Acebey. 2005. New species of Araceae from Bolivia and the tropical Andes. *Novon* 15:80-103.
- Croat, T. B., and M. M. Carlsen. 2013. A reassessment of *Anthurium* species with palmately divided leaves, and a reinterpretation of *Anthurium* section *Dactylophyllum* (Araceae). *PhytoKeys* 23:41-54.
- Croat, T. B., M. Joyce, and C. V. Kostelac. 2010. New species of Araceae from Ecuador. *Willdenowia* 40:331-343.
- Crouch, N. R. 2012. *Ferns of Southern Africa: a comprehensive guide*. Penguin Random House South Africa, Cape Town.
- Croxall, J. 1975. The Hymenophyllaceae of Queensland. *Australian Journal of Botany* 23:509-547.
- Croxall, J. P. 1986. *Microgonium* (Hymenophyllaceae) in Malesia, with special reference to Peninsular Malaysia. *Kew Bulletin* 41:519-531.
- Crutwell, N. E. G. 1988. Plant hunting in Papua New Guinea with special reference to *Rhododendrons*. In *Proceedings of the Fourth International Rhododendron Conference*, Wollongong, New South Wales, Australia.
- Cuello, N. L. 2002. Altitudinal changes of forest diversity and composition in the ramal de guamacal in the Venezuelan Andes. *Ecotrópicos* 15:160-176.
- da Costa, A. F., T. Fontoura, and A. M. Amorim. 2012. Novelties in Bromeliaceae from the northeastern Brazilian Atlantic Rainforest. *Journal of the Torrey Botanical Society* 139:34-45.
- Dahal, S. 2015-16. *Sikkim Himalayan Rhododendron*. *Panda* 8:15-27.
- Dalström, S., and S. Ruíz Pérez. 2015. Three new species of *Masdevallia* (Orchidaceae: Pleurothallidinae) from the ayacucho and puno regions in Peru. *Lankesteriana* 15:141-149.
- Dalström, S., G. Deburghgraeve, and S. R. Pérez. 2012. Three new showy but endangered *Cyrtorchilum* species (Orchidaceae: Oncidiinae) from Peru. *Lankesteriana* 12:93-99.
- Darwin, S. P. 1994. Systematics of *Timonius* subgenus *Abbottia* (Rubiaceae-Guettardeae). *Systematic Botany Monographs* 42:1-86.
- Dawson, J. W. 1988. *Forest vines to snow tussocks: the story of New Zealand plants*. Victoria University Press, Wellington.
- de Araujo Goes-Neto, L. A., and M. R. Pietrobon. 2012. Aspleniaceae (Polypodiopsida) from the Northern Para Biodiversity Corridor, Brazil: a fragment of the Guiana Centre of Endemism. *Acta Botanica Brasílica* 26:456-463.
- de Gandolle, C. 1919. Beiträge zur Kenntnis der Piperaceen von Papuasien *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie*:204-220.
- De Gasper, A. L., V. A. D. O. Dittrich, A. R. SMITH, and A. Salino. 2016. A classification for Blechnaceae (Polypodiales: Polypodiopsida): New genera, resurrected names, and combinations. *Phytotaxa* 275:191-227.
- De la Sota, E. R. 1965. Las especies escamosas del genero *Polypodium* L. (s. str.) en Brasil. *Revista del Museo de La Plata* 9:243-271.
- De la Sota, E. R. 1966. Revisión de las especies americanas del grupo *Polypodium squamatum* L. *Polypodiaceae* (s. str.). *Revista del Museo de La Plata* 10:69-186.
- De la Sota, E. R. 1968. Acerca del género *Synammia* Presl (*Polypodiaceae* s.str.). *Revista del Museo de La Plata* 11:129-132.
- De la Sota, E. R., and O. G. Martinez. 2005. The pteridophytic diversity in the cloud montane forests from Salta, NW Argentina. *Fern Gazette* 17:138.
- de Matos, F. B. 2015. Systematic studies of *Elaphoglossum* section *Polytrichia* (Dryopteridaceae). City University of New York, New York.
- de Roon, A. C. 2005. A new species of *Ruyschia* (Marcgraviaceae) from the South American Andes. *Novon* 15:414-417.

- de Roon, A. C., and D. Giraldo-Cañas. 2006. Contributions toward a monograph of *Norantea* complex (Marcgraviaceae, Ericales): Validation of four new species for *Marcgraviastrum*. *Caldasia* 28:15-22.
- De Wilde, J.J.F.E., 2002. *Begonia* section *Tetraphila* A.DC, a taxonomic revision. Wageningen Agricultural University Papers: 5-258.
- Deepa, J., T. R. Parashurama, M. Krishnappa, and S. Nataraja. 2013. Pteridophytic flora of Kemmangundi Forest, Karnataka, South India. *Annals of Plant Sciences* 2:484-488.
- Derroire, G., L. Schmitt, J. N. Riviere, J. M. Sarrailh, and J. Tassin. 2007. The essential role of tree-fern trunks in the regeneration of *Weinmannia tinctoria* in rain forest on Reunion, Mascarene Archipelago. *Journal of Tropical Ecology* 23:487-492.
- Dessai, J. R. N., L. Joseph, and M. K. Janarthanam. 2009. A new species of epiphytic *Impatiens* (Balsaminaceae) from the Western Ghats, India. *Taiwania* 54:149-151.
- Díaz, I. A., K. E. Sieving, M. E. Peña-Foxon, J. Larraín, and J. J. Armesto. 2010. Epiphyte diversity and biomass loads of canopy emergent trees in Chilean temperate rain forests: A neglected functional component. *Forest Ecology and Management* 259:1490-1501.
- Diels, L. 1938. Neue Arten aus Ecuador. *Notizblatt des Königl. botanischen Gartens und Museums zu Berlin* 14:25-44.
- Diels, L. 1942. Neue Arten aus Ecuador V. *Notizblatt des Königl. botanischen Gartens und Museums zu Berlin* 15:783-787.
- Diem, J., and J. S. D. Lichtenstein. 1959. Las Himenofiláceas del área argentino-chilena del sur. *Darwiniana* 11:611-760.
- Dittrich, V. A. d. O., J. L. Waechter, and A. Salino. 2005. Species richness of pteridophytes in a montane Atlantic rain forest plot of Southern Brazil. *Acta Botanica Brasílica* 19:519-525.
- Dodero, G. U., and F. Almeda. 1991. *Blakea costaricensis* (Melastomataceae): A remarkable new epiphyte from southeastern Costa Rica. *Selbyana* 12:1-4.
- Donadio, S. 2011. A valid name for the taxon known as *Tillandsia bryoides* auct. (Bromeliaceae). *Darwiniana* 49:131-138.
- Dong, S., L. Wei, and Y. Chao. 2012. A new species of *Asplenium* section *Thamnopteris* (Aspleniaceae) from Indonesia. *Blumea* 57:190-194.
- Dressler, R. L. 2005. New orchid species from Costa Rica. *Novon* 5:141-145.
- Dressler, R. L., and N. H. Williams. 1970. The genus *Systeloglossum*. *American Orchid Society Bulletin* 39:323-328.
- Droissart, V., B. Sonke, O. J. Hardy, M. Simo, H. Taedoumg, C. K. Nguembou, and T. Stevart. 2011. Do plant families with contrasting functional traits show similar patterns of endemism? A case study with Central African Orchidaceae and Rubiaceae. *Biodiversity and Conservation* 20:1507-1531.
- Dubuisson, J. Y., G. Rouhan, A. Grall, S. Hennequin, B. Senterre, K. Pynee, and A. Ebihara. 2013. New insights into the systematics and evolution of the filmy fern genus *Crepidomanes* (Hymenophyllaceae) in the Mascarene Archipelago with a focus on dwarf species. *Acta Botanica Gallica* 160:173-194.
- Dubuisson, J. Y., L. Bauret, A. Grall, B. Senterre, A. H. Said, K. Pynee, A. Ebihara, and S. Hennequin. 2017. Taxonomic study of the fern genera *Abrodictyum* C.Presl and *Trichomanes* L. (Hymenophyllaceae, Polypodiidae) in the western Indian Ocean, and description of a new *Abrodictyum* species for Madagascar. *Phytotaxa* 327:201-222.
- Dubuisson, J. Y., S. Hennequin, and V. Droissart. 2016. *Hymenophyllum senterreanum* Dubuisson & Deblauwe, sp. nov. (Hymenophyllaceae) and its relatives in western Central Africa. *Phytotaxa* 257:287-294.
- Dubuisson, J. Y., S. Hennequin, F. Rakotondrainibe, and H. Schneider. 2003. Ecological diversity and adaptive tendencies in the tropical fern *Trichomanes* L.

- (Hymenophyllaceae) with special reference to climbing and epiphytic habits. *Botanical Journal of the Linnean Society* 142:41-63.
- Dubuisson, J.-Y., S. Hennequin, S. Bary, A. Ebihara, and E. Boucheron-Dubuisson. 2011. Anatomical diversity and regressive evolution in trichomanoid filmy ferns (Hymenophyllaceae): A phylogenetic approach. *Comptes Rendus Biologies* 334:880-895.
- Earnshaw, M. J., K. Winter, H. Ziegler, W. Stichler, N. E. G. Cruttwell, K. Kerenga, P. J. Cribb, J. Wood, J. R. Croft, K. A. Carver, and T. C. Gunn. 1987. Altitudinal changes in the incidence of crassulacean acid metabolism in vascular epiphytes and related life forms in Papua New Guinea. *Oecologia* 73:566-572.
- Ebihara, A. 2008. Hymenophyllaceae collected in the Mt. Kinabalu Area in 2000 and 2007. *Memoirs National Museum Natural Sciences* 45:105-110.
- Ebihara, A., and K. Iwatsuki. 2007. The Hymenophyllaceae of the Pacific Area. 1. Hymenophyllum subgenus Hymenophyllum. *Bulletin of the National Museum of Natural Sciences—B* 33:55-68.
- Ebihara, A., C. R. Fraser-Jenkins, B. S. Parris, X.-C. Zhang, Y.-H. Yang, W.-L. Chiou, H.-M. Chang, S. Lindsay, D. Middleton, and M. Kato. 2012. Rare and threatened pteridophytes of Asia 1. An enumeration of narrowly distributed taxa. *Bulletin of the National Museum of Nature and Science, Series B* 38:93-119.
- Ebihara, A., J. H. Nitta, and K. Iwatsuki. 2010. The Hymenophyllaceae of the Pacific Area. 2. Hymenophyllum (excluding subgen. Hymenophyllum). *Bulletin of the National Museum of Nature and Science Series B, Botany* 36:43-59.
- Ebihara, A., J.-Y. Dubuisson, K. Iwatsuki, S. Hennequin, and M. Ito. 2006. A taxonomic revision of Hymenophyllaceae. *Blumea* 51:221-280.
- Ebihara, A., K. Iwatsuki, T. A. Ohsawa, and M. Ito. 2003. Hymenophyllum paniense (Hymenophyllaceae), a new species of filmy fern from New Caledonia. *Systematic Botany* 28:228-235.
- Ebihara, A., S. Matsumoto, and M. Ito. 2009. Taxonomy of the reticulate Vandemboschia radicans complex (Hymenophyllaceae) in Japan. *Acta Phytotaxonomica et Geobotanica* 60:26-40.
- Edinburgh Royal Botanical Garden Herbarium Catalogue.
<https://data.rbge.org.uk/search/herbarium/>.
- Eggli, U. 2005. *Illustrated handbook of succulent plants: Crassulaceae*. Springer, Berlin.
- Elmer, A. D. E. 1910a. The Ericaceae of Mount Apo. *Leaflets of Philippine Botany* 3:1089-1107.
- Elmer, A. D. E. 1910b. New and interesting Gesneriaceae. *Leaflets of Philippine Botany* 3:947-970.
- Elmer, A. D. E. 1910c. New and noteworthy Rubiaceae. *Leaflets of Philippine Botany* 3:971-1046.
- Endres Junior, D., M. H. Sasamori, and A. Droste. 2014. In vitro propagation of Anathallis adenochila (Loefgr.) F. Barros (Orchidaceae), a species endemic to southern and southeastern Brazil. *Acta Botanica Brasílica* 28:489-494.
- Engemann, K., Sandel, B., Boyle, B., Enquist, B.J., Jørgensen, P.M., Kattge, J., McGill, B.J., Morueta-Holme, N., Peet, R.K., Spencer, N.J., Violle, C., Wiser, S.K., Svenning, J.-C., 2016. A plant growth form dataset for the New World. *Ecology* 97:3243-3243.
- Engwald, S. 1999. *Diversität und Ökologie der vaskulären Epiphyten in einem Berg- und einem Tieflandregenwald in Venezuela*. Books on Demand, Bonn.
- Enright, N. J., and E. K. Cameron. 1988. The soil seed bank of a kauri (Agathis australis) forest remnant near Auckland, New Zealand. *New Zealand Journal of Botany* 26: 223-236.
- Eskov, A. K. pers. comm.

- Espejo-Serna, A., A. R. López-Ferrari, and J. Ceja Romero. 1996. *Maianthemum comaltepecense* (Convallariaceae), una nueva especie del estado de Oaxaca, Mexico. *Acta Botánica Mexicana* 36:21-28.
- Espinosa-Jimenez, A. J., A. M. Perez-Farrera, and R. Martinez-Camilo. 2011. Floristic inventory of Sumidero Canyon National Park, Chiapas, Mexico. *Boletín de la Sociedad Botánica de México* 89:37-82.
- Esser, H.-J., and M. H. P. Jebb. 2010. A new *Schefflera* and taxonomic notes on Araliaceae from Thailand. *Thai Forest Bulletin (Botany)* 38:160-166.
- Etl, F., A. Franschitz, A. J. C. Aguiar, J. Schonenberger, and S. Dotterl. 2017. A perfume-collecting male oil bee? Evidences of a novel pollination system involving *Anthurium acutifolium* (Araceae) and *Paratetrapedia chocoensis* (Apidae, Tapinotaspidini). *Flora* 232:7-15.
- Evans, A. M., and J. T. Mickel. 1969. A re-evaluation of *Polypodium bradeorum* and *P. colysoides*. *Brittonia* 21:255-260.
- Fall, P. L., and T. D. Drezner. 2020. Vascular plant species of the Kingdom of Tonga by vegetation type, species origin, growth form, and dispersal mechanism. *Ecology* 101.
- Fedorov, A. A. 1959. Woody epiphytes and strangler figs in the tropical [rain]forests of China. *Botanicheskii Zhurnal St Petersburg* 44:1409-1424.
- Ferreira de Lima, R. A., V. A. de Oliveira Dittrich, V. C. de Souza, A. Salino, T. B. Breier, and O. T. de Aguiar. 2011. Flora vascular do Parque Estadual Carlos Botelho, São Paulo, Brasil. *Biota Neotropica* 11:173-214.
- Ferreira, R. D., A. C. C. Borges, J. A. R. dos Campos, A. M. L. Medeiros, C. M. Sakuragui, R. C. Vieira, and V. Tenorio. 2020. Anatomy of the adventitious roots of *Philodendron* (Araceae) and its importance for the systematics of the genus. *Australian Systematic Botany* 33:207-219.
- Ferreras, U. F., and G. C. G. Argent. 2011. *Diplycosia bartolomei* (Ericaceae), a new species from the Philippines. *Edinburgh Journal of Botany* 68:39-42.
- Field, D. V., and J. B. Hall. 1982. *Epistemma*, a new genus of Periplocaceae from West Africa. *Kew Bulletin* 37:117-120.
- Fieldmuseum. without year. Fieldguides; <https://fieldguides.fieldmuseum.org/>.
- Figueiredo, E. 2001. A revision of *Calvoa* Hook. f. (Melastomataceae). *Botanical Journal of the Linnean Society* 136:179-205.
- Figueiredo, E. 2005. The Rubiaceae of Sao Tome e Principe (Gulf of Guinea): taxonomy and conservation. *Botanical Journal of the Linnean Society* 149:85-114.
- Figueiredo, E., and A. Gascoigne. 2001. Conservation of pteridophytes in São Tomé e Príncipe (Gulf of Guinea). *Biodiversity & Conservation* 10:45-68.
- Fischer, E. 1996. *Die Vegetation des Parc National de Kahuzi-Biéga, Sud-Kivu, Zaïre*. Franz Steiner, Stuttgart.
- Fischer, E. pers. comm.
- Fischer, E., and M. E. Rahelivololona. 2002. New taxa of *Impatiens* (Balsaminaceae) from Madagascar. I. *Adansonia* 24:271-294.
- Fischer, E., and M. E. Rahelivololona. 2004a. A new epiphytic species of *Impatiens* (Balsaminaceae) from the Comoro Islands. *Adansonia* 26:93-95.
- Fischer, E., and M. E. Rahelivololona. 2004b. New taxa of *Impatiens* (Balsaminaceae) from Madagascar. III. *Adansonia* 26:37-52.
- Fischer, E., and M. E. Rahelivololona. 2007. New taxa of *Impatiens* (Balsaminaceae) from Madagascar. IV. *Adansonia* 29:269-315.
- Fischer, E., H. J. T. Venter, D. Killmann, U. Meve, and A. M. Venter. 2011. *Epistemma neuerburgii* (Apocynaceae, Periplocoideae): A new epiphytic species from Nyungwe National Park, Rwanda, Africa. *South African Journal of Botany* 77:680-684.

- Fischer, E., K. Rembold, A. Althof, J. Obholzer, I. Malombe, G. Mwachala, J. C. Onyango, B. Dumbo, and I. Theisen. 2010. Annotated checklist of the vascular plants of Kakamega Forest, Western Province, Kenya. *Journal of East African Natural History* 99:129-226.
- Fischer, E., S. Wohlhauser, and M. E. Rahelivololona. 2003. New taxa of Impatiens (Balsaminaceae) from Madagascar. II. A collection from Masoala Peninsula. *Adansonia* 25:17-31.
- Flakus, P., and S. Beck. 2007. Distribution of an endemic local species on montane forest of Apa Apa – Chulumani: the case of *Coussapoa davidsmithii* (Cecropiaceae). *Revista Estudiantil de Biología* 5:44-54.
- Flora of Australia online. without year. <http://www.environment.gov.au/>.
- Flora of China. 2008. Published on the Internet <http://www.efloras.org> [last accessed January 2020] Missouri Botanical Garden, St. Louis, MO & Harvard University Herbaria, Cambridge, MA. .
- Flora_Mesoamericana. in progress. <http://www.tropicos.org/Project/FM>.
- Flora_of_Australia. Australian Biological Resources Study, Canberra. <http://www.ausflora.org.au>.
- Flora_of_Guatemala. http://www.archive.org/stream/floraofguatemala241034stan/floraofguatemala241034stan_djvu.txt.
- Flora_of_North_America. http://www.efloras.org/flora_page.aspx?flora_id=1.
- Florence, J., S. Waldren, and A. J. Chepstow-Lusty. 1995. The flora of the Pitcairn Islands: a review. *Biological Journal of the Linnean Society* 56:79-119.
- Fontoura, T., M. A. Rocca, A. C. Schilling, and F. Reinert. 2009. Epífitas da floresta seca da Reserva Ecológica Estadual de Jacarepiá, sudeste do Brasil: relações com a comunidade arbórea. *Rodriguésia* 60:171-185.
- Forbes, H. O. 1888. A new fern from New Guinea. *Journal of Botany, British and Foreign* 26:33.
- Forest_Flora. <http://www.forestflora.co.nz/>.
- Forney, C. F., W. Kalt, and S. P. Vander Kloet. 2012. Comparison of berry composition of selected *Vaccinium* species (Ericaceae) with *Gaylussacia dumosa*. *Botany-Botanique* 90:355-363.
- Fowler, R. L. 1940. Annotated list of ferns of the Kilauea-Mauna Loa Section of Hawaii National Park. *American Fern Journal* 30:9-18.
- Freitas, J., and A. M. de Assis. 2013. Structure of vascular epiphyte component in a fragment of the Atlantic Forest in the mountain region of Espírito Santo. *Revista Arvore* 37:815-823.
- Freitas, L., A. Salino, L. Menini Neto, T. Almeida, S. Mortara, J. Stehmann, A. M. Amorim, E. Guimaraes, M. A. Nadruz Coelho, A. Zanin, and R. Forzza. 2016. A comprehensive checklist of vascular epiphytes of the Atlantic Forest reveals outstanding endemic rates. *PhytoKeys* 58:65-79.
- Friis, I. 1989. A revision of *Pilea* (Urticaceae) in Africa. *Kew Bulletin* 44:557-600.
- Friis, I., and K. Vollesen. 1998. Flora of the Sudan-Uganda border area east of the Nile, I: Catalogue of vascular plants, 1st part. The Royal Danish Academy of Sciences and Letters, Copenhagen.
- Fuentes, A. F. 2013. Nuevos registros de la región Madidi (Dpto. La Paz), para la flora de plantas vasculares de Bolivia. *Kempffiana* 9:3-20.
- Fujiwara, T., S. Serizawa, and Y. Watano. 2018. Phylogenetic analysis reveals the origins of tetraploid and hexaploid species in the Japanese *Lepisorus thunbergianus* (Polypodiaceae) complex. *Journal of Plant Research* 131:945-959.

- Gabanchó, L. R. 2015. La colección de helechos y licófitos cubanos del Hno. Clemente en el herbario HAC/Collection of Cuban ferns and lycophytes by Bro. Clement, conserved in HAC herbarium. *Revista del Jardín Botánico Nacional* 36:201-218.
- Garay, L. A. 1972. On the systematics of the monopodial orchids I. *Botanical Museum Leaflets Harvard University* 23:149-212.
- Gay, H., E. Hennipman, C. R. Huxley, and F. J. E. Parrott. 1993. The taxonomy, distribution and ecology of the epiphytic Malesian ant-fern *Lecanopteris Reinw.* (Polypodiaceae). *Gardens' Bulletin, Singapore* 45:293-335.
- Gemma, J. N., and R. E. Koske. 1995. Mycorrhizae in Hawaiian epiphytes. *Pacific Science* 49:175-180.
- Gentry, A. 1987. Bookreview: The botany of mangroves by P. B. Tomlinson. *Annals of the Missouri Botanical Garden* 74:460-462.
- Gentry, A. H., and C. H. Dodson. 1987. Diversity and biogeography of Neotropical vascular epiphytes. *Annals of the Missouri Botanical Garden* 74:205-233.
- Gerlach, G. pers. comm.
- Gill, L. S., and H. I. Onyibe. 1986. Phytosociological studies of epiphytic flora of oil palm (*Elaeis guineensis* Jacq.) in Benin City, Nigeria. *Feddes Repertorium* 97:691-695.
- Gilli, A. 1978. Beiträge zur Flora von Papua-New Guinea. I. Bryophyta, Pteridophyta und Coniferae. *Annalen des Naturhistorischen Museums in Wien* 81:19-29.
- Gilli, A. 1980. Beiträge zur Flora von Papua-New Guinea II. Dicotylédones. *Annalen des Naturhistorischen Museums in Wien* 83:417-474.
- Giongo, C., and J. L. Waechter. 2004. Composição florística e estrutura comunitária de epífitos vasculares em uma floresta de galeria na Depressão Central do Rio Grande do Sul. *Revista Brasileira de Botânica* 27:563-572.
- Giraldo-Cañas, D. 2011. Estudios morfológicos y taxonómicos en el complejo *Norantea* (Marcegraviaceae): Revisión de *Norantea*, *Pseudosarcopera* y *Schwartzia*. *Biblioteca José Jerónimo Triana (Bogotá D. C.: Instituto de Ciencias Naturales, Universidad Nacional de Colombia)* 22:1-176.
- Giraldo-Cañas, D. pers. obs.
- Global_Plants. 2019. <https://plants.jstor.org/>.
- Goebel, K. 1889. *Pflanzenbiologische Schilderungen*. 1. Teil. Elwert'sche Verlagsbuchhandlung, Marburg.
- Gómez-Pompa, A., and J. Valdés. 1962. Una especie epífita de *Yucca* de la selva lacandona. *Boletín de la Sociedad Botánica de México* 27:43-46.
- Gómez-Pompa, A., T. Krömer, and R. Castro-Cortés. 2010. Atlas de la flora de Veracruz.
- Gonzatti, F., and P. G. Windisch. 2018. Flora of Espírito Santo: *Hymenophyllum* (Hymenophyllaceae). *Rodriguésia* 69:611-629.
- Grant, J. R., and L. Struwe. 2001. De *Macrocarpaea* Grisebach (ex gentianacezs) *Speciebus novis* I: An introduction to the genus *Macrocarpaea* and three new species from Colombia, Ecuador, and Guyana. *Harvard Papers in Botany* 5:489-498.
- Grayum, M. H., and H. W. Churchill. 1989. The vascular flora of La Selva Biological Station, Costa Rica - Polypodiophyta. *Selbyana* 11:66-118.
- Green, P. S. 1986. Notes relating to the floras of Norfolk and Lord Howe Islands, II. *Journal of the Arnold Arboretum* 67:109-122.
- Grey-Wilson, C. 1980. *Impatiens of Africa*. A.A.Balkema, Rotterdam.
- Gruezo, W. S. 2012. *Tmesipteris zamorae*: A new species of *Tmesipteris* Bernh. (Psilotaceae) from the Philippines. *Asia Life Sciences* 21:653-670.
- Gustafsson, M.H.G., Borchsenius, F. 2016. *Clusia nubium* (Clusiaceae): a new species from cloud-forests of southwestern Ecuador. *Phytotaxa* 253:219-222.

- Guyane: flore et de la végétation. without year.
<https://www.flickr.com/photos/stationalpinejosephfourier/collections/72157627913811661/>.
- Guzman, V. pers. comm.
- Haantjens, H. A., J. R. McAlpine, E. Reiner, R. G. Robbins, and J. C. Saunders. 1970. Lands of the Goroka–Mount Hagen Area, territory of Papua and New Guinea. *Land Research Series* 27:1-172.
- Haber, W. 2000a. Appendix 1 - Vascular plants of Monteverde. Pages 457-518 in N. M. Nadkarni and N. Whellwright, editors. *Monteverde: Ecology and conservation of a tropical cloud forest*. Oxford University Press, New York.
- Haber, W. 2000b. Appendix 2 - Number of species with different plant growth form. Pages 519-522 in N. M. Nadkarni and N. Whellwright, editors. *Monteverde: Ecology and conservation of a tropical cloud forest*. Oxford University Press, New York.
- Haber, W. A. 2003. *Monteverde Natural History*.
<https://www.cs.umb.edu/~whaber/Monte/Plant/>.
- Hahn, W.J. 1996. Notes on Costa Rican Aquifoliaceae. *Novon* 6:181-184.
- Hallé, N. 1970. Famille de Rubiacees (2e partie). Muséum National d'Histoire Naturelle, Paris.
- Hallier, H. 1906. Neue indonesische Dikotyledonen. *Repertorium novarum specierum regni vegetabilis* 2:59-64.
- Hammel, B. E. 1999. Two new species of *Clusiella* (Clusiaceae) with a synopsis of the genus. *Novon* 9:349-359.
- Hammel, B. E., M. H. Grayum, C. Herrera, N. Zamora, and S. Troyo. 2003. *Gymnospermas y Monocotiledoneas (Agavaceae-Musaceae)*. Missouri Botanical Garden, St. Louis, Miss.
- Hardy, C. R. 2001. Systematics of *Cochliostema*, *Geogenanthus*, and an undescribed genus in the spiderwort family, *Commelinaceae*. Cornell University, Ithaca, NY, USA.
- Harris, R., G. Cassis, T. Auld, and I. Hutton. 2005. Floristics and structure of the mossy cloud forest of Mt Gower summit, Lord Howe Island. *Pacific Conservation Biology* 11:246-256.
- Harrison, R. D. pers. comm.
- Hawaii Tropical Botanical Garden. without year. <http://www.htbg.com/>.
- He, S.-Y., Li, P.-T., Lin, J.-Y., Yang, X.-H., 2009a. *Hoya persiciniconaria* (Apocynaceae, Asclepiadoideae), a new species from Hainan, China. *Novon* 19:475-478.
- He, S.-Y., Li, P.-T., Lin, J.-Y., Lin, G., Zeng, H.-L., 2011. *Hoya jianfenglingensis* (Apocynaceae), a new species from Hainan, China. *Novon* 21:343-346.
- He, S.-Y., X.-Y. Zhuang, P.-T. Li, J.-Y. Lin, and M. Li. 2009b. *Hoya baishaensis* (Apocynaceae), a new species from Hainan, China. *Annales Botanici Fennici* 46:155-158.
- Heads, M. 2003. Ericaceae in Malesia: vicariance biogeography, terrane tectonics and ecology. *Telopea* 10:311-449.
- Hecklau, E.F., Mori, S.A., Brown, J.L. 2005. Specific epithets of the flowering plants of central French Guiana. *Brittonia* 57:68-87.
- Helme, N. A., and P. J. Rakotomalaza. 1999. An overview of the botanical communities of the Reserve Naturelle Intégrale d'Andohahela, Madagascar. *Fieldiana Zoology* 30:11-23.
- Henderson, M. R. 1927. A list of plants from Cameron's Highlands, Pahang. *Journal of the Malayan Branch of the Royal Asiatic Society* 5:237-277.
- Hennequin, S., A. Ebihara, M. Ito, K. Iwatsuki, and J.-Y. Dubuisson. 2006. New insights into the phylogeny of the genus *Hymenophyllum* s.l. (Hymenophyllaceae): revealing the polyphyly of *Mecodium*. *Systematic Botany* 31:271-284.

- Hennequin, S., E. Schuettpelz, K. M. Pryer, A. Ebihara, and J. Y. Dubuisson. 2008. Divergence times and the evolution of epiphytism in filmy ferns (Hymenophyllaceae) revisited. *International Journal of Plant Sciences* 169:1278-1287.
- Herbarium_Berolinense_Virtual_Herbarium. <http://ww2.bgbm.org/herbarium/default.cfm>.
- Herbarium_Universität_Ulm. http://www.biologie.uni-ulm.de/cgi-bin/query_all/details.pl?id=47508&stufe=A&typ=PFL&sid=T&lang=d&pr=nix.
- Hermans, J., and P. Cribb. 2005. 519. *Paralophia epiphytica*. *Curtis's Botanical Magazine* 22:47-52.
- Hernandez, J. A., and M. A. Diaz. 2000. A new species of *Tetramicra* (Orchidaceae) from eastern Cuba. *Harvard Papers in Botany* 5:189-192.
- Herrero, A., C. Aedo, M. Velayos, and R. L. L. Viane. 2001. A new species of *Asplenium* (Aspleniaceae, Pteridophyta) from Equatorial Guinea. *Annales Botanici Fennici* 38:175-180.
- Hietz, P., and U. Hietz-Seifert. 1995. Composition and ecology of vascular epiphyte communities along an altitudinal gradient in central Veracruz, Mexico. *Journal of Vegetation Science* 6:487-498.
- Hietz-Seifert, U., P. Hietz, and S. Guevara. 1995. Epiphyte vegetation and diversity on remnant trees after forest clearance in Southern Veracruz, Mexico. *Biological Conservation* 75:103-111.
- Hilliard, O. M., and B. L. Burt. 2002. The genus *Agalmyla* (Gesneriaceae-Cyrtandroideae). *Edinburgh Journal of Botany* 59:1-210.
- Hirai, R. Y., and J. Prado. 2012. Monograph of *Moranopteris* (Polypodiaceae). *Taxon* 4:1-113.
- Hirai, R. Y., G. Rouhan, P. H. Labiak, T. A. Ranker, and J. Prado. 2011. *Moranopteris*: A new Neotropical genus of grammitid ferns (Polypodiaceae) segregated from Asian *Micropolypodium*. *Taxon* 60:1123-1137.
- Hoang, V. S. 2009. Uses and conservation of plant diversity in Ben En National Park Vietnam. PhD. Universiteit Leiden, Leiden.
- Hodge, W. H. 1941. Notes on Dominica Ferns (Conclusion). *American Fern Journal* 31:121-127.
- Hofstede, R. G. M., K. J. M. Dickinson, and A. F. Mark. 2001. Distribution, abundance and biomass of epiphyte-lianoid communities in a New Zealand lowland Nothofagus-podocarp temperate rain forest: tropical comparisons. *Journal of Biogeography* 28:1033-1049.
- Hokche, O., P. E. Berry, and O. Huber, editors. 2008. Nuevo catálogo de la flora vascular de Venezuela. Fundación Instituto Botánico de Venezuela, Caracas, Venezuela.
- Holbrook, N. M., and F. E. Putz. 1996a. From epiphyte to tree: differences in leaf structure and leaf water relations associated with the transition in growth form in eight species of hemiepiphytes. *Plant, Cell and Environment* 19:631-642.
- Holbrook, N. M., and F. E. Putz. 1996b. Water relations of epiphytic and terrestrially-rooted strangler figs in Venezuelan palm savanna. *Oecologia* 106:424-431.
- Holst, B. K. pers. comm.
- Holtermann, C. 1907. Der Einfluß des Klimas auf den Bau der Pflanzengewebe. Wilhelm Engelmann, Leipzig.
- Holtum, R. E. 1937. Further notes on *Stenochlaena*, *Lomariopsis* and *Teratophyllum*. *The Gardens' Bulletin, Straits Settlements* 9:139-144.
- Holtum, J. A. M., J. E. Aranda, A. Virgo, H. H. Gehrig, and K. Winter. 2004. $\delta^{13}\text{C}$ values and crassulacean acid metabolism in *Clusia* species from Panama. *Trees* 18:658-668.
- Homeier, J., F. A. Werner, S. R. Gradstein, S. W. Breckle, and M. Richter. 2008. Potential vegetation and floristic composition of Andean forests in South Ecuador, with a focus on the RBSF. *Ecological Studies* 198:87-100.

- Hopkins, M., R. Maulder, and B. Salomon. 1990. A real nice trip to Southeast Asia. CPNewsletter 19:19-28.
- Householder, E., J. Janovec, A. B. Mozambique, J. H. Maceda, J. Wells, R. Valega, H. Maruenda, and E. Christenson. 2010. Diversity, natural history, and conservation of Vanilla (Orchidaceae) in Amazonian wetlands of Madre de Dios, Peru. Journal of the Botanical Research Institute of Texas 4:227-243.
- Hovenkamp, P. 1986. A monograph of the fern genus Pyrrosia (Polypodiaceae). Leiden University Press, Leiden.
- Hovenkamp, P. 1997. Paraselliguea, a new genus of Malesian Polypodiaceae. Blumea 42:485-487.
- Hovenkamp, P. 1998. An account of the Malay-Pacific species of Selligiea. Blumea 43:1-108.
- Hovenkamp, P. 2011. A new species of Selligiea (Polypodiaceae) from Thailand. Edinburgh Journal of Botany 68:1-3.
- Hovenkamp, P., and B. C. Ho. 2012. A revision of the fern genus Oleandra (Oleandraceae) in Asia. PhytoKeys 11:1-37.
- Hovenkamp, P., and F. Miyamoto. 2005. A conspectus of the native and naturalized species of Nephrolepis (Nephrolepidaceae) in the world. Blumea 50:279-322.
- Hsieh, T.-Y., K. A. Hatch, and Y.-M. Chang. 2012. Phlegmariurus changii (Huperziaceae), a new hanging firmoss from Taiwan. American Fern Journal 102:283-288.
- Hsu, R. C. C., W. L. M. Tamis, N. Raes, G. R. de Snoo, J. H. D. Wolf, J. G. B. Oostermeijer, and S.-H. Lin. 2012. Simulating climate change impacts on forests and associated vascular epiphytes in a subtropical island of East Asia. Diversity and Distributions 18:334-347.
- Hsu, R., and J. H. D. Wolf. 2009. Diversity and phytogeography of vascular epiphytes in a tropical-subtropical transition island, Taiwan. Flora 204:612-627.
- Huang, Y.-M., V. B. Amoroso, F. P. Coritico, C.-W. Ko, T.-T. Kao, J. R. Callado, and W.-L. Chiou. 2015. Reproductive biology of Aglaomorpha cornucopia (Copel.) M.C. Roos, a rare and endemic fern from the Philippines. American Fern Journal 105:31-44.
- Huber, O., J. A. Steyermark, G. T. Prance, and C. Ales. 1984. The vegetation of the Sierra Parima, Venezuela-Brazil - some results of recent exploration. Brittonia 36:104-139.
- Hunt, D., N. Taylor, and G. Charles, editors. 2006. The new cactus lexicon. Descriptions and illustrations of the cactus family. dh Books, Milborne Port.
- Huxley, C. R., and M. H. P. Jebb. 1991a. The tuberous epiphytes of the Rubiaceae 2: The new genus Anthorrhiza. Blumea 36:21-41.
- Huxley, C. R., and M. H. P. Jebb. 1991b. The tuberous epiphytes of the Rubiaceae 3: A revision of Myrmephytum to include Myrmedoma. Blumea 36:43-52.
- Hyde, M. A., B. T. Wursten, P. Ballings, and M. Coates Palgrave. 2019. Flora of Zimbabwe: Home page. <https://www.zimbabweflora.co.zw/index.php>.
- Hyndman, D. C., and J. I. Menzies. 1990. Rain forests of the Ok Tedi headwaters, New Guinea: An ecological analysis. Journal of Biogeography 17:241-273.
- Ibisch, P. L. 1996. Neotropische Epiphytendiversität - das Beispiel Bolivien. Martina Galunder-Verlag, Wiehl, Germany.
- Ibrahim, H. 1999. Zingiberaceous species of Tawau Hill Parks, Sabah ASEAN Review of Biodiversity and Environmental Conservation (ARBEC) 6:1-10.
- India_Biodiversity_Portal. Huperzia hilliana <https://indiabiodiversity.org/species/show/267182>.
- Indonesian_Biodiversity_Information_System_(IBIS)_online. in construction. <http://ibis.biologi.lipi.go.id/>.
- Internet_Orchid_Species_Photo_Encyclopedia. <http://www.orchidspecies.com/>.

- Irume, M. V., M. de Lourdes da Costa Soares Morais, C. E. Zartman, and I. L. do Amaral. 2013. Floristic composition and community structure of epiphytic angiosperms in a terra firme forest in central Amazonia. *Acta Botanica Brasílica* 27:378-393.
- Ishii, H. R., T. Minamino, W. Azuma, K. Hotta, and A. Nakanishi. 2018. Large, retained trees of *Cryptomeria japonica* functioned as refugia for canopy woody plants after logging 350 years ago in Yakushima, Japan. *Forest Ecology and Management* 409:457-467.
- Isnard, S., J. Prosperi, S. Wanke, S. T. Wagner, M.-S. Samain, S. Trueba, L. Frenzke, C. Neinhuis, and N. P. Rowe. 2012. Growth form evolution in Piperales and its relevance for understanding Angiosperm diversification: an integrative approach combining plant architecture, anatomy, and biomechanics. *International Journal of Plant Sciences* 173:610-639.
- Iwatsuki, K. 2016. Studies on Asian ferns. *Journal of Japanese Botany*, 91:330-334.
- Iwatsuki, K., A. Ebihara, and M. Kato. 2019. Taxonomic studies of pteridophytes of Ambon and Seram (Moluccas) collected on Indonesian-Japanese botanical expeditions 1983-1986. XIII. Hymenophyllaceae. *PhytoKeys* 119:107-115.
- Iwatsuki, K., and M. G. Price. 1977. The pteridophytes of Mt Burnay and vicinity, Northern Luzon. *The southeast asian studies* 14:540-572.
- Jacques-Félix, H. 1977. Le genre *Preussiella* Gilg (Melastomataceae). *Adansonia sér.* 2 16:405-414.
- Jaman, R., and A. Latiff. 1998. On some pteridophytes of Sayap-Kinabalu Park, Sabah. *ASEAN Review of Biodiversity and Environmental Conservation (ARBEC)* IV:1-35.
- Jaman, R., and A. Latiff. 1999. The pteridophyte flora of Pulau Tioman, Peninsular Malaysia. *Raffles Bulletin of Zoology* 6:77-100.
- Jamir, S. A., and H. N. Pandey. 2003. Vascular plant diversity in the sacred groves of Jaintia Hills in northeast India. *Biodiversity and Conservation* 12:1497-1510.
- Janssen, T., and H. Schneider. 2005. Exploring the evolution of humus collecting leaves in drynarioid ferns (Polypodiaceae, Polypodiidae) based on phylogenetic evidence. *Plant Systematics and Evolution* 252:175-197.
- Janssens, S. B., E. Fischer, and T. Stévant. 2010. New insights into the origin of two new epiphytic *Impatiens* species (Balsaminaceae) from West Central Africa based on molecular phylogenetic analyses. *Taxon* 59:1508-1518.
- Jenny, R. 2007. *Cuitlauzina pendula* - History and culture of an orchid described in 1825. *Orchids* 76:296-301.
- Jermey, A. C., and K. P. Kavanagh. 1982. *Notulae et novitates Muluenses*. *Botanical Journal of the Linnean Society* 85:1-73.
- Jimeno Sevilla, H. D. 2008. El género *Echeveria* (Crassulaceae) en Veracruz, México. Universidad Veracruzana, Xalapa, Mexico.
- Jing, H. J., Q. Fan, W. Guo, and W. B. Liao. 2016. *Sorbus prunifolia* (Rosaceae), a new species from Yunnan Province, China. *Phytotaxa* 258:171-178.
- Jobson, R. W., J. Playford, K. M. Cameron, and V. A. Albert. 2003. Molecular phylogenetics of Lentibulariaceae inferred from plastid rps16 intron and trnL-F DNA sequences: Implications for character evolution and biogeography. *Systematic Botany* 28:157-171.
- Johansson, D. 1974. Ecology of vascular epiphytes in West African rain forest. *Acta Phytogeographica Suecica* 59:1-136.
- Johns, R. J., G. A. Shea, and P. Puradyatmika. 2012. Subalpine and alpine vegetation of Papua. Pages 1025-1053 in A. J. Marshall and B. M. Beehler, editors. *Ecology of Indonesian Papua Part II*.
- Johnson, P. N., and D. J. Campbell. 1975. Vascular plants of the Auckland Islands. *New Zealand Journal of Botany* 13:665-720.
- Jongkind, C. C. H. 2005. *Novitates Gabonenses* 54. A new species of *Gardenia* (Gardenieae, Rubiaceae) from Gabon. *Systematics and Geography of Plants* 75:259-262.

- Jørgensen, P. M., and S. León-Yáñez. 1999. Catalogue of the vascular plants of Ecuador. Missouri Botanical Garden.
- Jørgensen, P. M., M. Nee, and S. Beck, editors. 2014. Catálogo de las plantas vasculares de Bolivia. Missouri Botanical Press, St. Louis.
- Judd, W. S. 1986. Botany of the National Parks of Haiti. Department of Botany, University of Florida, Gainesville.
- Julius, A., A. Takano, M. Suleiman, and W. F. Tukin. 2010. Zingiberaceae of Maliau Basin, Sabah, Malaysia. *Journal of Tropical Biology and Conservation* 6:1-29.
- Kadereit, G. 2005. Revision of *Plethiandra* Hook. f.: A polystaminate, East Asian genus of Melastomataceae. *Edinburgh Journal of Botany* 62:127-144.
- Kadereit, G. pers. comm.
- Kang, Y. J. 2009. Species richness of epiphyte on forest vegetation in Jeju. PhD Dissertation. Jeju National University, Jeju, Korea, Jeju.
- Kanzaki, M., and K. Sri-ngernyuang. 2012. Diversity and dynamics of epiphyte, hemiepiphyte, and parasite in tropical forests of Doi Inthanon National Park (2008-2012). National Research Council of Thailand.
- Kartonegoro, A. 2010. A revision of *Dissochaeta* (Melastomataceae) in Java, Indonesia. *Reinwardtia* 13:125-145.
- Kato, M., and M. G. Price. 1990. Taxonomic studies of pteridophytes of Ambon and Seram (Moluccas) collected by Indonesian-Japanese Botanical Expeditions V.: Dipteridaceae and Polypodiaceae. *Acta Phytotaxonomica et Geobotanica* 41:61-76.
- Keller, H. W., P. G. Davison, C. H. Haufler, and D. B. Lesmeister. 2003. *Polypodium appalachianum*: An unusual tree canopy epiphyte in the Great Smoky Mountains National Park. *American Fern Journal* 93:36-41.
- Kelly, D. L. 1985. Epiphytes and climbers of a Jamaican rain forest: vertical distribution, life forms and life histories. *Journal of Biogeography* 12:223-241.
- Keppel, G., and N. T. Thomas. 2009. Composition and structure of the cloud forest on Mt Delaco, Gau, Fiji. *The South Pacific Journal of Natural Science* 27:28-34.
- Kersten, R. A., and S. M. Silva. 2006. The floristic compositions of vascular epiphytes of a seasonally inundated forest on the coastal plain of Ilha do Mel Island, Brazil. *Revista de Biología Tropical* 54:935-942.
- Kersten, R. d. A., and Y. S. Kuniyoshi. 2009. Conservação das florestas na bacia do alto iguaçu, Paraná – avaliação da comunidade de epífitas vasculares em diferentes estágios serais. *Floresta* 39:51-56.
- Kersten, R. D., Y. S. Kuniyoshi, and C. V. Roderjan. 2009. Vascular epiphytes of two nearby riverside forests, Iguacu River Basin, Parana. *Iheringia Serie Botanica* 64:33-43.
- Kessler, M. pers. comm.
- Kessler, M., A. L. M. Velazquez, M. Sundue, and P. H. Labiak. 2011. *Alansmia*, a new genus of grammitid ferns (Polypodiaceae) segregated from *Terpsichore*. *Brittonia* 63:233-244.
- Kessler, M., and A. R. Smith. 2017. Prodrum of a fern flora for Bolivia. X. *Hymenophyllaceae*. *Phytotaxa* 328:201-226.
- Kessler, M., and A. R. Smith. 2018. Prodrum of a fern flora for Bolivia. XXIX. *Aspleniaceae*. *Phytotaxa* 344:259-280.
- Kessler, M., R. Moran, J. T. Mickel, F. B. Matos, and A. R. Smith. 2018. Prodrum of a fern flora for Bolivia. XXXV. *Dryopteridaceae*. *Phytotaxa* 353:1-114.
- Khine, P. K., C. Fraser-Jenkins, S. Lindsay, D. Middleton, G. Miehe, P. Thomas, and J. Kluge. 2017. A contribution toward the knowledge of ferns and lycophytes from Northern and Northwestern Myanmar. *American Fern Journal* 107:219-256.
- Khine, P. K., S. Lindsay, C. Fraser-Jenkins, J. Kluge, M. Kyaw, and P. Hovenkamp. 2016. *Selliguea kachinensis* (Polypodiaceae), a new fern species of uncertain affinity from Northern Myanmar. *PhytoKeys*:73-81.

- Khwaiphan, W., and T. Boonkerd. 2008. The pteridophyte flora of Khao Khiao, Khao Yai National Park, Thailand. *The Natural History Journal of Chulalongkorn University* 8:69-82.
- Kidyoo, M. 2012. *Hoya mirabilis* Kidyoo, a New Species of *Hoya* (Asclepiadaceae) from Western Thailand. *Tropical Natural History* 12:21-28.
- Kidyoo, M. 2015. *Hoya rostellata* (Apocynaceae: Asclepiadoideae), a New Species from Thailand. *Taiwania* 60:3-42.
- Kidyoo, M., and J. Wai, Sae. 2009. A new record of *Hoya* (Asclepiadaceae) from southern Thailand. *The Natural History Journal of Chulalongkorn University* 9:235-238.
- Kiew, R. 1989. *Hoya endauensis* (Asclepiadaceae) and *Licuala dranifieldii* (Palmae), two new species from Ulu Endau, Peninsular Malaysia *Malayan Nature Journal* 42:261-265.
- Kiew, R., and S. Anthonysamy. 1987. A comparative study of vascular epiphytes in three epiphyte-rich habitats at Ulu Endau, Johore, Malaysia. *Malayan Nature Journal* 41:303-315.
- Kimmach, M. 1990. A new epiphytic cactus: *Trichocereus arboricola*. *Cactus and Succulent Journal* 62:3-5.
- King, R. M., and H. Robinson. 1969. A new genus, *Neomirandea*. *Phytologia* 19:305-310.
- Kirby, C. 2014. Field guide to New Zealand's epiphytes, vines & mistletoes Environmental Research Institute, University of Waikato, Hamilton, NZ.
- Kleijn, D., and R. Van Donkelaar. 2001. Notes on the taxonomy and ecology of the genus *Hoya* (Asclepiadaceae) in central Sulawesi. *Blumea* 46:457-483.
- Kloppenburger, R. D. 2004. *Hoya* species;
<https://hoyardk.wixsite.com/kloppenburger/publications>.
- Kloppenburger, R. D., S. V. Siar, J. M. Carandang, and A. S. Hadsall. 2011. *Hoya ilagii*, another new species (Apocynaceae, Asclepiadoideae) from Luzon Island, Philippines. *Asia Life Sciences* 20:263-269.
- Kluge, J. pers. comm.
- Knapp, R. 2011. Ferns and fern allies of Taiwan. KBCC Press, Goashu, Taiwan.
- Knapp, R., and T.-C. Hsu. 2017. Fern and fern allies of Taiwan, 2nd supplement. Dr. Cecilia Koo Botanic Conservation Center (KBCC Press), Taipei.
- Knapp, S. 1995. New taxa and combinations in the tribe Juanulloeae (Solanaceae). *Novon* 5:281-283.
- Knapp, S. 1998. New species and notes on the natural history of Markea (Solanaceae) from Colombia and Ecuador. *Novon* 8:152-161.
- Knapp, S., V. Persson, and S. Blackmore. 1997. A phylogenetic conspectus of the tribe Juanulloeae (Solanaceae). *Annals of the Missouri Botanical Garden* 84:67-89.
- Kolbek, J. 1995. Notes on epiphytic communities in forests of North Korea. *Preslia, Praha* 67:41-45.
- Kornas, J. 1976. Notes on African Hymenophyllaceae 2. *Trichomanes lenormandii* v. d. Bosch New to Continental Africa. *Bulletin du Jardin botanique national de Belgique / Bulletin van de National Plantentuin van Belgique* 46:393-397.
- Kornás, J. 1977. Life-forms and seasonal patterns in the pteridophytes in Zambia. *Acta Societatis Botanicorum Poloniae* 46:669-690.
- Kornás, J. 1984. Bulletin of pteridology — 4. *Taxon* 33:752-755.
- Kornas, J. 1993. Filmy ferns (Hymenophyllaceae) of Central Africa (Zaire, Rwanda, Burundi). 1. *Hymenophyllum*. *Fragmenta Floristica et Geobotanica Polonica* 38:3-19.
- Korotkova, N. pers. comm.
- Korotkova, N., L. Zabel, D. Quandt, and W. Barthlott. 2010. A phylogenetic analysis of Pfeiffera and the reinstatement of *Lymanbensonia* as an independently evolved lineage of epiphytic Cactaceae within a new tribe *Lymanbensoniaceae*. *Willdenowia* 40:151-S154.

- Köster, N. pers. comm.
- Köster, N., H. Kreft, J. Nieder, and W. Barthlott. 2013. Range size and climatic niche correlate with the vulnerability of epiphytes to human land use in the tropics. *Journal of Biogeography* 40:963–976.
- Kramer, K. U. 1972. The Lindsaeoid ferns of the Old World: IX Africa and its islands. *Bulletin du Jardin botanique national de Belgique / Bulletin van de National Plantentuin van België* 42:305-345.
- Krause, K. 1932. Araceae novae austro-americanae. *Notizblatt des Königlich botanischen Gartens und Museums zu Berlin* 11:606-622.
- Kreft, H., N. Köster, W. Küper, J. Nieder, and W. Barthlott. 2004. Diversity and biogeography of vascular epiphytes in Western Amazonia, Yasuní, Ecuador. *Journal of Biogeography* 31:1463-1476.
- Kreier, H. P., X. C. Zhang, H. Muth, and H. Schneider. 2008. The microsoroid ferns: Inferring the relationships of a highly diverse lineage of Paleotropical epiphytic ferns (Polypodiaceae, Polypodiopsida). *Molecular Phylogenetics and Evolution* 48:1155-1167.
- Krömer, T. 2003. Diversität und Ökologie der vaskulären Epiphyten in primären und sekundären Bergwäldern Boliviens. PhD. Universität Göttingen, Göttingen.
- Kumar, M. S. M. 1998. Studies on the fern flora of Kerala with special reference to Sylvan Valley, Munnar. Report 46, Kerala Forest Research Institute
- Kumar, M., and S. Sequiera. 2012. Observations on the epiphytic flora of Silent Valley
- Kunkel, G. 1962. Aufzeichnungen über die Vegetation des Mt. Wutivi-Gebietes (Nordwest-Liberia), unter besonderer Berücksichtigung der Pteridophyten. *Willdenowia* 3:151-168.
- Kunkel-Westphal, I., and P. Kunkel. 1979. Litter fall in a Guatemalan primary forest, with details of leaf-shedding by some common tree species. *Journal of Ecology* 67:665-686.
- Kvist, L. P. 1990. Revision of *Heppiella* (Gesneriaceae). *Systematic Botany* 15:720-735.
- Kvist, L. P., and L. E. Skog. 1992. Novae Gesneriaceae Neotropicarum IV: *Alloplectus purpureus* and *Columnea nematoloba* - New Gesneriaceae from Northwestern South-America. *Brittonia* 44:475-480.
- Labiak, P. H. 2013. Grammitid ferns (Polypodiaceae): I. *Lellingeria*. *Flora Neotropica*:1-130.
- Labiak, P. H., and J. P. S. Condack. 2008. *Ceradenia Itatiaiensis*, a new species of Polypodiaceae from Southeastern Brazil. *Novon* 18:495-497.
- Labiak, P. H., and J. Prado. 1998. Pteridófitas epífitas da Reserva Volta Velha, Itapoá - Santa Catarina, Brasil. *Boletim do Instituto de Botânica* 11:1-79.
- Labiak, P. H., and J. Prado. 2003. Grammitidaceae (Pteridophyta) no Brasil com ênfase nos gêneros *Ceradenia*, *Cochlidium* e *Grammitis*. *Hoehnea* 30:243-283.
- Labiak, P. H., and J. Prado. 2005a. As espécies de *Terpsichore* A.R. Sm. e *Zygophlebia* L.E. Bishop (Grammitidaceae) do Brasil. *Acta Botanica Brasílica* 19:867-887.
- Labiak, P. H., and J. Prado. 2005b. The species of the genus *Lellingeria* A.R. Sm. & R.C. Moran (Grammitidaceae - Pteridophyta) from Brazil. *Brazilian Journal of Botany* 28:1-22.
- Labiak, P. H., and J. Prado. 2008. New combinations in *Serpocaulon* and a provisional key for the atlantic rain forest species. *American Fern Journal* 98:139-159.
- Labiak, P. H., F. B. Matos, G. Rouhan, J. G. Hanks, and R. C. Moran. 2017. Notes on the taxonomy and growth habits of three species of *Campyloneurum* (Polypodiaceae) from Southeastern Brazil. *American Fern Journal* 107:1-20.
- Labiak, P. H., G. Rouhan, and M. Sundue. 2010a. Phylogeny and taxonomy of *Leucotrichum* (Polypodiaceae): A new genus of grammitid ferns from the Neotropics. *Taxon* 59:911-921.

- Labiak, P. H., M. Sundue, and G. Rouhan. 2010b. Molecular phylogeny, character evolution, and biogeography of the grammitid fern genus *Lellingeria* (Polypodiaceae). *American Journal of Botany* 97:1354-1364.
- LaFrankie, J. V. 1986. A new species of *Maianthemum* (Liliaceae) from Costa Rica with an upright and aerial rhizome. *American Journal of Botany* 73:1258-1260.
- Lamb, A., Gobilik, J., Ardiyani, M., Poulsen, A.D., 2013. A guide to ginger of Borneo. Natural History Publications (Borneo), Kota Kinabalu.
- Lambdon, P., and A. Darlow. 2012. Flowering plants & ferns of St Helena. Pisces Publications for St Helena Nature Conservation Group.
- Lammers, T. G. 1991. Systematics of *Clermontia* (Campanulaceae-Lobelioideae). *Systematic Botany Monographs* 32:1-97.
- Lammers, T. G. 2009. Revision of the endemic Hawaiian genus *Trematolobelia* (Campanulaceae: Lobelioideae). *Brittonia* 61:126-143.
- Lampard, S. 2016. *Pinguicula casabitoana* J. Jiménez Alm. Pages 402-681 in A. S. Robinson, editor. *Pinguicula of Latin America*. Redfern Natural History Productions, Poole, Dorset, England, pp. 402-681.
- Larsen, C., M. M. Ponce, and M. A. Scataglini. 2013. Revisión de las especies de *Hymenophyllum* (Hymenophyllaceae) del sur de Argentina y Chile. *Gayana. Botánica* 70:275-330.
- Lauterbach, C. 1919. Beiträge zur Flora von Papuasien. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie*:19-144.
- Lawton, R. O. 1986. The evolution of strangling by *Ficus crassiuscula*. *Brenesia* 25-26:273-278.
- Lazare, J.-J., J. Vivant, and C. Sastre. 1991. Les ptéridophytes de Guadeloupe: biodiversité, écologie, protection. *Bulletin de la Société Botanique de France. Actualités Botaniques* 138:197-214.
- Leal, E. d. S., and R. C. Forzza. 2012. Cyclanthaceae in Para State, Brazil. *Acta Botanica Brasílica* 26:822-835.
- Leao Barros, I. C., L. C. Cantarelli, R. d. P. Farias, A. F. de Novaes Pereira, and I. A. Araujo da Silva. 2014. Vertical distribution of epiphytic ferns in a fragment of the Atlantic Forest in the Brazilian Northeast. *Iheringia Serie Botanica* 69:143-153.
- Leena, K. R., and P. V. Madhusoodanan. 1998. Pteridophyte flora of Wynad District, Kerala: A preliminary survey on ecology and distribution. *Journal of Economic and Taxonomic Botany* 22:149-155.
- Leenhouts, P. W. 1962. Loganiaceae. *Flora Malesiana* 6:293-387.
- Leeuwenberg, A. J. M. 1958. The Gesneriaceae of Guiana - a critical revision with notes on species from adjacent regions. *Acta Botanica Neerlandica* 7:291-435.
- Lehnert, M. 2013. Grammitid ferns (Polypodiaceae): *Melpomene*. *Flora Neotropica Monograph* 112:1-121.
- Lehtonen, S., H. Tuomisto, G. Rouhan, and M. J. M. Christenhusz. 2010. Phylogenetics and classification of the pantropical fern family Lindsaeaceae. *Botanical Journal of the Linnean Society* 163:305-359.
- Leitman, P., A. Amorim, L. M. Neto, and R. C. Forzza. 2014. Epiphytic angiosperms in a mountain forest in southern Bahia, Brazil. *Biota Neotropica* 14:1-12.
- Lellinger, D. B. 1972. A revision of the fern genus *Niphidium*. *American Fern Journal* 62:101-121.
- Lellinger, D. B. 1984. New combinations and some new names in ferns. *American Fern Journal* 74:56-60.
- Lellinger, D. B. 1985. A field manual to the ferns and fern-allies of the United States and Canada. Smithsonian Institution Press, Washington, D.C.

- Lellinger, D. B. 1989. The ferns and fern-allies of Costa Rica, Panama, and the Chocó (Part I: Psilotaceae through Dicksoniaceae). American Fern Society, Washington, D.C.
- Lellinger, D. B. 1991. Notes on neotropical Hymenophyllaceae. American Fern Journal 81:24-36.
- Leme, E. M. C., and L. J. C. Kollmann. 2013. Miscellaneous new species of Brazilian Bromeliaceae. Phytotaxa 108:1-40.
- Leme, E. M. C., W. Till, L. J. C. Kollmann, R. L. De Moura, and O. B. C. Ribeiro. 2014. Miscellaneous new species of Brazilian Bromeliaceae - III. Phytotaxa 177:61-100.
- León, B. 1992. A taxonomic revision of the fern genus *Campyloneurum* (Polypodiaceae). Aarhus Universitet.
- León, B. 2004. Una especie nueva de *Campyloneurum* (Polypodiaceae) del norte del Perú. Revista Peruana de Biología 11:135-137.
- León, B. 2006a. Grammitidaceae endémicas del Perú. Revista Peruana de Biología 13:901-902.
- León, B. 2006b. Lycopodiaceae endémicas del Perú. Revista Peruana de Biología 13:911-912.
- León, B., and H. Beltrán. 2002. A New Microgramma Subgenus *Solanopteris* (Polypodiaceae) from Peru and a new combination in the subgenus. Novon 12:481-485.
- León-Parra, A. 2012. *Ceradenia* (Polypodiaceae). Flora de Colombia 29:1-83.
- Leroux, O., A. Bagniewska-Zadworna, S. K. Rambe, J. P. Knox, S. E. Marcus, E. Bellefroid, D. Stubbe, B. Chabbert, A. Habrant, M. Claeys, and R. L. L. Viane. 2011. Non-lignified helical cell wall thickenings in root cortical cells of *Aspleniaceae* (Polypodiales): histology and taxonomical significance. Annals of Botany 107:195-207.
- Lesica, P., and R. K. Antibus. 1990. The occurrence of mycorrhizae in vascular epiphytes of two Costa Rican rain forests. Biotropica 22:250-258.
- Li, F.-W., B. C. Tan, V. Buchbender, R. C. Moran, G. Rouhan, C.-N. Wang, and D. Quandt. 2009. Identifying a mysterious aquatic fern gametophyte. Plant Systematics and Evolution 281:77-86.
- Li, Y.-h., L.-n. Liu, and Q.-S. Ye. 2005. Effects of elevated CO₂ on the growth and photosynthesis of *Anthurium andraeanum*. Journal of Tropical and Subtropical Botany 13:343-346.
- Liede-Schumann, S., and U. Meve. 2006. The Genera of Asclepiadoideae, Secamonoideae and Periplocoideae (Apocynaceae) Descriptions, Illustrations, Identification, and Information Retrieval. Version: 21st September 2000. http://tftp://www.uni-bayreuth.de/departments/planta2/research/databases/delta_ascleps/index.html.
- Lijun, C., and L. Zhongjian. 2010. *Chroniochilus sinicus*, a new species of Orchidaceae from Yunnan, China. Novon 20:252-255.
- Lindau, G. 1921. Eine epiphytische Acanthacee, *Dischistocalyx epiphytica* Lindau nov. spec. . Notizblatt des Botanischen Gartens und Museums zu Berlin-Dahlem 70:494.
- Lindner, A., K. Stein, and M. Freiberg. 2010. Abundance and vigor of three selected understory species along environmental gradients in south-eastern Brazil. Ecotropica 16:101-112.
- Lindsay, S. 2003. *Antrophyum brassii* (Vittariaceae), a new fern species from Papua New Guinea. Harvard Papers in Botany 8:5-8.
- Lindsay, S., and D. J. Middleton. 2003. (1598) Proposal to conserve the name *Davallia repens* (L.f.) Kuhn (Davalliaceae) against *D. repens* (Bory) Desv. (Lindsaeaceae). Taxon 52:630-631.
- Lindsay, S., and D. J. Middleton. 2012 onwards. Ferns of Thailand, Laos and Cambodia. <http://rbg-web2.rbge.org.uk/thaiferns/>.
- Liogier, A. H., H. A. Liogier, and L. F. Martorell. 2000. Flora of Puerto Rico and adjacent islands: a systematic synopsis. La Editorial, UPR.
- Lipps, S. 2005. Madeira - Was hier alles wächst. Oliver Breda, Duisburg.

- Little, E. L. 1970. New tree species from Esmeralda, Ecuador. *Phytologia* 19:251-269.
- Liu, H.-M., R.-H. Jiang, J. Guo, P. Hovenkamp, L. R. Perrie, L. Shepherd, S. Hennequin, and H. Schneider. 2013. Towards a phylogenetic classification of the climbing fern genus *Arthropteris*. *Taxon* 62:688-700.
- Liu, W. Y., J. E. D. Fox, and Z. F. Xu. 2001. Community characteristics, species diversity and management of middle-mountain moist evergreen broad-leaved forest in the Ailao Mountains, Southwestern China. *Pacific Conservation Biology* 7:34-44.
- Löfstrand, S. D., A. Krüger, S. G. Razafimandimbison, and B. Bremer. 2014. Phylogeny and generic delimitations in the sister tribes Hymenodictyeae and Naucleaeae (Rubiaceae). *Systematic Botany Monographs* 39:304-315.
- Looser, G. 1948. The ferns of southern Chile. *American Fern Journal* 38:71-87.
- Lorence, D. H., and G. Rouhan. 2004. A revision of the Mascarene species of *Elaphoglossum* (Elaphoglossaceae). *Annals of the Missouri Botanical Garden* 91:536-565.
- Loriga, J., A. Vasco, L. Regalado, J. Heinrichs, and R. C. Moran. 2014. Phylogeny and classification of the Cuban species of *Elaphoglossum* (Dryopteridaceae), with description of *Elaphoglossum* sect. *Wrightiana* sect. nov. *Plant Systematics and Evolution* 300:937-951.
- Lovett, J. C., and D. Thomas. 1986. The ecology of pteridophytes in the Mwanihana Forest Reserve, Tanzania. *Fern Gazette* 13:103-107.
- Luis, D. G. P. 1976. Variation in Costa Rican *Ophioglossum palmatum* and nomenclature of the species. *American Fern Journal* 66:89-92.
- Luna-Vega, I., J. Daniel Tejero-Diez, R. Contreras-Medina, M. Heads, and G. Rivas. 2012. Biogeographical analysis of two *Polypodium* species complexes (Polypodiaceae) in Mexico and Central America. *Biological Journal of the Linnean Society* 106:940-955.
- Luo, W., X.-O. Song, H. Xu, H. Huang, C.-W. Yu, H. Hu, and H.-O. Chen. 2010. Floristic analysis of pteridophytes in Jianfengling Nature Reserve, Hainan Island. *Journal of Wuhan Botanical Research* 28:294-302.
- Lutelyn, J. L. 2002. Key to the species of Ericaceae of Bolivia, including two new species. *SIDA Contributions to Botany* 20:1-20.
- Lutelyn, J. L., and P. Pedraza-Peñalosa. 2012 onward. Blueberry relatives of the New World tropics (Ericaceae) (<http://sweetgum.nybg.org/ericaceae/index.php>). The New York Botanical Garden, Bronx, New York.
- Lüttge, U., E. Ball, M. Kluge, and B. L. Ong. 1986. Photosynthetic light requirements of various tropical vascular epiphytes. *Physiologie Végétale* 24:315-331.
- Mabberley, D. J. 1987. *The Plant-Book. A portable dictionary of the higher plants*. Cambridge University Press, Cambridge.
- Macbride, J. F. 1962. Solanaceae. *Flora of Peru*, Field Museum of Natural History, Botanical Series, Volume XIII.
- Maciel, J. R., R. Louzada, and M. Alves. 2014. *Aechmea nigribacteata* (Bromeliaceae), a new species from southern Bahia, Brazil. *Phytotaxa* 188:233-237.
- Madhusoodanan, P. V., and C. A. Hameed. 1998. *Crepidomanes agasthianum*, a new filmy fern species (Hymenophyllaceae — Pteridophyta) from India. *Nordic Journal of Botany* 18:169-170.
- Maesen, L. J. G. v. d., X. M. v. d. Burgt, and J. M. v. M. d. Rooy, editors. 1996. *The Biodiversity of African Plants*. Kluwer Academic Publishers.
- Maguire, B., and J. J. Wurdack. 1958. The botany of the Guyana Highland - Part III. *Memoirs of the New York Botanical Garden* 10:1-156.
- Mai, P., A. Rossado, J. M. Bonifacino, and J. L. Waechter. 2019. Catalogue of the vascular epiphytic flora of Uruguay. *Acta Botanica Brasílica*.

- Maia, V. C., and M. G. Santos. 2015. Record of insects in two fern species of the genus *Microgramma* (Polypodiaceae) in the Atlantic Rain Forest, Rio de Janeiro state, Brazil. *Brazilian Journal of Biology* 75:S253-S254.
- Maideen, H., Z. C. Desa, A. Damanhuri, A. Latiff, and G. Rusea. 2011. Diversity and habitat of pteridophytes in Angsi Forest Reserve, Negeri Sembilan. *Sains Malaysiana* 40:1341-1344.
- Makgomol, K. 2009. Ferns in the waterfall areas in Nam Nao National Park, Thailand. *KKU Science Journal* 37:56-61.
- Malizia, A. 2003. Host tree preference of vascular epiphytes and climbers in a subtropical montane cloud forest of Northwest Argentina. *Selbyana* 24:196-205.
- Maloney, B. K. 1983. The plant ecology of forest remnants on the Toba plateau of North Sumatra. *Indonesia Circle. School of Oriental & African Studies. Newsletter* 11:23-37.
- Mangambu, M. J. D. D. 2013. Taxonomie, biogeographie et ecologie des Pteridophytes de l'écosystème forestier des montagnes du Parc National de Kahuzi-Biega a l'Est de la R.D. Congo. Antwerpen.
- Mansfeld, R. 1936. Gesneriaceae austroamericanae novae vel rariores. III. Repertorium novarum specierum regni vegetabilis 41:145-150.
- Mantovani, A. 2000. Leaf orientation of epiphytic aroids: effect on water and temperature balances of the leaves. *Leandra* 15:91-103.
- Martínez, O. G., F. C. Assis, E. I. M. Torres, D. A. Cacharani, and D. G. Jaimez. 2016. El género *Pecluma* (Polypodiaceae) en Argentina. *Darwiniana* 4:234-251.
- Martínez-Meléndez, N., Martínez Camilo, R., Pérez Farrera, M.Á., Martínez Meléndez, J., 2011. Las epífitas de la Reserva El Triunfo, Chiapas. Colección Jaguar Unicach, Tuxtla Gutiérrez, Chiapas, Mexico.
- Massa, G. W. 1996. Factors affecting the distribution of a neotropical hemiepiphyte. MSc. San Jose State University, San Jose.
- Mathieu, G. 2008. Three new creeping *Peperomia* species (Piperaceae) from Ecuador. *Novon* 18:80-85.
- Mathieu, G., D. Vergara-Rodriguez, T. Krömer, and D. N. Karger. 2015. *Peperomia* (Piperaceae) novelties from Veracruz State, Mexico. *Phytotaxa* 205:268-276.
- Matos, F. B., A. M. Amorim, and P. H. Labiak. 2010. The ferns and lycophytes of a montane tropical forest in southern Bahia, Brazil. *Botanical Research Institute of Texas* 4:333-346.
- Matos, F. B., and A. Vasco. 2015. *Elaphoglossum fendleri* (Dryopteridaceae), a new species of *Elaphoglossum* sect. *Lepidoglossa* from Venezuela, and the identity of *Elaphoglossum ornatum*. *Brittonia* 67:144-149.
- Matos, F. B., and J. T. Mickel. 2014. The Brazilian species of *Elaphoglossum* section *Polytrichia* (Dryopteridaceae). *Brittonia* 66:371-395.
- Matos, F. B., and J. T. Mickel. 2018. The Brazilian species of *Elaphoglossum* section *Setosa* (Dryopteridaceae). *Brittonia* 70:173-205.
- Matos, F. B., P. H. Labiak, and L. S. Sylvestre. 2009. A new Brazilian species of the genus *Asplenium* L. (Aspleniaceae). *American Fern Journal* 99:101-105.
- Matos, F. B., P. H. Labiak, and R. C. Moran. 2019. The *Decursivum* group of *Elaphoglossum* sect. *Polytrichia* (Dryopteridaceae): Nonsubulate species nested in an otherwise subulatescaled clade1. *Annals of the Missouri Botanical Garden* 104:400-428.
- Maxon, W. R. 1914. Studies of tropical American ferns—no. 5. Contributions from the United States National Herbarium 17:391-425.
- Maxwell, J. F. 2009. Vegetation and vascular flora of the Mekong River, Kratie and Steung Treng Provinces, Cambodia. *Maejo International Journal of Science and Technology* 3:143-211.
- McMillan, P. D., G. Wyatt, and R. Morris. 2006. The *Begonia* of Veracruz: Additions and revisions. *Acta Botánica Mexicana* 75:77-99.

- McPherson, S. 2009. Pitcher Plants of the Old World. Redfern Natural History Productions, Poole, Dorset, England.
- Meave, J. A., M. A. Romero-Romero, A. Valle-Doménech, A. Rincón-Gutiérrez, E. Martínez, and C. H. Ramos. 2008. Plant diversity assessment in the Yaxchilán Natural Monument, Chiapas, Mexico. *Boletín de la Sociedad Botánica de México* 83:53-76.
- Medeiros, A. C., L. L. Loope, and S. J. Anderson. 1993. Differential colonization of epiphytes on native (*Cibotium* spp.) and alien (*Cyathea cooperi*) tree ferns in a Hawaiian rain forest. *Selbyana* 14:71-74.
- Meekiong, K., I. Ipor, C. S. Tawan, H. Ibrahim, M. R. Norhati, C. K. Lim, and A. Ampeng. 2011. Five new ginger species (Zingiberaceae) from the eastern part of Lanjak Entimau Wildlife Sanctuary, Sarawak, Borneo. *Folia Malaysiana* 12:9-26.
- Mellado, F., and B. León. 2007. New records and observations of some species of *Elaphoglossum* (Elaphoglossaceae) of Peru. *Revista Peruana de Biología* 14:21-23.
- Mellado-Mansilla, D., I. Díaz, and R. Moreno. 2015. Ampliación del rango de distribución de *Hymenophyllum ferrugineum colla* (Hymenophyllaceae) para Chile continental. *Chlorischile* 18:1.
- Mendoza-Ruiz, A., J. Ceja-Romero, and B. Pérez-García. 2016. Helechos y licofitos epífitos de Veracruz, México: riqueza y distribución. *Acta Botánica Mexicana* 114:87-136.
- Mendum, M. 2003. The Gesneriaceae of Sulawesi III: Three new species of *Aeschynanthus*. *Edinburgh Journal of Botany* 60:323-330.
- Menini Neto, L., R. C. Forzza, and D. Zappi. 2009. Angiosperm epiphytes as conservation indicators in forest fragments: A case study from southeastern Minas Gerais, Brazil. *Biodiversity and Conservation* 18:3785-3807.
- Menini Neto, L., S. G. Furtado, D. C. Zappi, A. T. de Oliveira Filho, and R. C. Forzza. 2016. Biogeography of epiphytic Angiosperms in the Brazilian Atlantic forest, a world biodiversity hotspot. *Brazilian Journal of Botany* 39:261-273.
- Menon, S., M. L. Khan, A. Paul, and A. T. Peterson. 2012. *Rhododendron* species in the Indian Eastern Himalayas: New approaches to understanding rare plant species distribution. *Journal of the American Rhododendron Society*:78-84.
- Menzel, F., and N. Blüthgen. 2010. Parabiotic associations between tropical ants: equal partnership or parasitic exploitation? *Journal of Animal Ecology* 79:71-81.
- Meyer, C. F. J., and G. Zotz. 2004. Does growth and survival of aerial roots limit the vertical distribution of hemiepiphytic aroids? *Biotropica* 36:483-491.
- Meza Torres, E. I., E. R. De la Sota, and M. S. Ferrucci. 2013. Synopsis of ferns and lycophytes of the Mburucuya National Park (Corrientes, Argentina). Keys to species. *Boletín de la Sociedad Argentina de Botánica* 48:121-136.
- Mickel, J. T. 1980. Nine new *Elaphoglossums* (Elaphoglossaceae) from Mexico. *Brittonia* 32:334-339.
- Mickel, J. T. 1985. The proliferous species of *Elaphoglossum* (Elaphoglossaceae) and their relatives. *Brittonia* 37:261-278.
- Mickel, J. T. 1992. New species of the fern genus *Elaphoglossum* from Mesoamerica. *Novon* 2:368-382.
- Mickel, J. T. 1993. Six new species of *Elaphoglossum* (Elaphoglossaceae). *Brittonia* 45:213-218.
- Mickel, J. T. 2008. Three new species of *Elaphoglossum* (Dryopteridaceae) from the Guianas. *Brittonia* 60:213-216.
- Mickel, J. T., and A. R. Smith. 2004. The pteridophytes of Mexico (Memoirs of The New York Botanical Garden 88). The New York Botanical Garden, Bronx, New York, USA.
- Middleton, B. A. 2007. A new species and a new combination in *Aeschynanthus* (Gesneriaceae) from Laos. *Edinburgh Journal of Botany* 64:45-50.

- Milbraed, J., editor. 1914. Wissenschaftliche Ergebnisse der Deutschen Zentral-Africa-Expedition, 1907-1908 unter Führung Adolf Friedrichs, Herzogs zu Mecklenburg. http://www.archive.org/stream/wissenschaftlich02adol/wissenschaftlich02adol_djvu.txt.
- Mildbraed, J. 1922. Wissenschaftliche Ergebnisse der Zweiten Deutschen Zentral-Afrika-Expedition 1910-1911. Klinkhardt & Biermann, Leipzig.
- Möller, M., and Q. C. B. Cronk. 2001. Evolution of morphological novelty: A phylogenetic analysis of growth patterns in *Streptocarpus* (Gesneriaceae). *Evolution* 55:918-929.
- Monro, A. K., D. Santamaria-Aguilar, F. Gonzalez, O. Chacon, D. Solano, A. Rodriguez, N. Zamora, E. Fedele, and M. Correa. 2017. A first checklist to the vascular plants of La Amistad International Park (PILA), Costa Rica-Panama. *Phytotaxa* 322:1-283.
- Moore, S.-J., C.-M. Chen, and J.-C. Wang. 2010. *Hymenophyllum pilosissimum* C. Chr. (Hymenophyllaceae), a new recorded fern from Taiwan. *American Fern Journal* 100:180-183.
- Moorea_Digital_Flora_Project. <http://ucjeps.berkeley.edu/moorea/pteridophytes.html>.
- Mora, M., R. Bernal, T. Croat, and J. Jacome. 2006. A phytogeographic analysis of Araceae of Cabo Corrientes (Choco, Colombia) and comparable lowland tropical American floras. *Annals of the Missouri Botanical Garden* 93:359-366.
- Moran, R. C. 2000. Monograph of the neotropical species of *Lomariopsis* (Lomariopsidaceae). *Brittonia* 52:55-111.
- Moran, R. C. 2010. *Pleopeltis coenosora* (Polypodiaceae), a new species from Peru. *Novon* 20:311-313.
- Moran, R. C. pers. comm.
- Moran, R. C., and B. Øllgaard. 1998. New species of ferns (Polypodiopsida) from Ecuador. *Nordic Journal of Botany* 18:431-439.
- Moran, R. C., and P. H. Labiak. 2017. The 1-pinnate species of *Campyloneurum* (Polypodiaceae). *Brittonia* 69:186-196.
- Moran, R. C., and R. Riba. 1995. "Flora Mesoamericana." Volume 1 Psilotaceae a Salviniaceae. Universidad Nacional Autónoma de México, Mexico.
- Moran, R., and M. Kimnach. 2004. *Echeveria nebularum* (Crassulaceae), a new cloud-forest species from Mexico. *Haseltonia* 10:67-70.
- Morat, P., J. M. Veillon, and H. S. MacKee. 1986. Floristic relationships of New Caledonian rainforest phanerogams. *Telopea* 2:631-679.
- Mori, S. A. 2002. *Dicotyledons*. New York Botanical Garden, Bronx.
- Mori, S. A., and J. A. Kallunki. 1977. Revision of the genus *Phyllonoma* (Grossulariaceae). *Brittonia* 29:69-84.
- Mori, S. A., E. F. Hecklau, and T. Kirchgessner. 2002. Life form, habitat, and nutritional mode of the flowering plants of central French Guiana. *Journal of the Torrey Botanical Society* 129:331-345.
- Morton, C. V. 1947. The American species of *Hymenophyllum*, section *Sphaerocionium*. *Systematic Plant Studies*:139-188.
- Morton, C. V. 1948. Notes on *Elaphoglossum*: II. The species of the French West Indies. *American Fern Journal* 38:202-214.
- Muasya, A., Y. Harvey, M. Cheek, K. Tah, and D. Simpson. 2010. *Coleochloa domensis* (Cyperaceae), a new epiphytic species from Cameroon. *Kew Bulletin* 65:323-325.
- Mueller-Dombois, D., and F. R. Fosberg. 1998. *Vegetation of the tropical Pacific islands*. Springer, New York.
- Murakami, N. 1995. Systematics and evolutionary biology of the fern genus *Hymenasplenium* (Aspleniaceae). *Journal of Plant Research* 108:257-268.
- Murakami, N., and R. C. Moran. 1993. Monograph of the neotropical species of *Asplenium* sect. *Hymenasplenium* (Aspleniaceae). *Annals of the Missouri Botanical Garden* 80:1-38.

- Murdock, A. G., and A. R. Smith. 2003. Pteridophytes of Moorea, French Polynesia, with a new species, *Tmesipteris gracilis* (Psilotaceae). *Pacific Science* 57:253-265.
- Murillo, M. T., and A. R. Smith. 2003. *Luisma*, a new genus of Grammitidaceae (Pteridophyta) from Colombia. *Novon* 13:313-317.
- Nadkarni, N. M. 1986. An ecological overview and checklist of vascular epiphytes in the Monteverde Cloud Forest Reserve, Costa Rica. *Brenesia* 24:55-62.
- Nagamasu, H., and S. Sakai. 1996. *Amomum roseisquamosum* (Zingiberaceae), a new epiphytic ginger from Borneo. *Edinburgh Journal of Botany* 53:39-42.
- Narayanan, M. K. R., J. P. Joseph, N. A. Kumar, M. Sivadasan, and A. H. Alfarhan. 2013. *Impatiens theuerkaufiana* (Balsaminaceae), a new scapigerous species from the Western Ghats, India. *Phytotaxa* 83:54-60.
- Narayanan, M. K. R., K. A. Sujana, B. V., R. Raj, Meera , and N. A. Kumar. 2012. *Impatiens mohana* (Balsaminaceae), a new scapigerous balsam from Wayanad, Western Ghats, India. *Edinburgh Journal of Botany* 69:281-285.
- Naturalis_Biodiversity_Center_(NL:Botany). Neotropical Flora
<https://serv.biokic.asu.edu/neotrop/plantae/index.php>.
- Nayar, M. P. 1966. Contributions to the knowledge of Indo-Malaysian and other Asiatic Melastomataceae VII. A synopsis of the genus *Hederella* Stapf. *Kew Bulletin* 20:235-244.
- Negrelle, R. R. B. 2002. The Atlantic forest in the Volta Velha Reserve: a tropical rain forest site outside the tropics. *Biodiversity and Conservation* 11:887-919.
- New_Zealand_Native_Orchids_Group. no year. New Zealand's native orchids;
<http://www.nativeorchids.co.nz/>.
- New_Zealand_Plant_Conservation_Network. 2019. <http://www.nzpcn.org.nz/default.aspx>.
- Nitta, J. H., and M. J. Epps. 2009. Hemi-epiphytism in *Vandenboschia collariata* (Hymenophyllaceae). *Brittonia* 61:392-397.
- Nooteboom, H. P. 1996. *Microsorium aurantiacum*, a new species of microsoroid ferns. *Blumea* 41:17 - 18.
- Nugent, G., K. W. Fraser, and P. J. Sweetapple. 1997. Comparison of red deer and possum diets and impacts in podocarp-hardwood forest, Waihaha Catchment, Pureora Conservation Park. Department of Conservation, Wellington, New Zealand.
- Nuraliev, M. S., G. V. Degtjareva, D. D. Sokoloff, A. A. Oskolski, T. H. Samigullin, and C. M. Valiejo-Roman. 2014. Flower morphology and relationships of *Schefflera subintegra* (Araliaceae, Apiales): an evolutionary step towards extreme floral polymery. *Botanical Journal of the Linnean Society* 175:553-597.
- O'Byrne, P. 2015. Notes on Borneo orchids 2. *Malesian Orchid Journal* 16:11-48.
- Ohwi, J. 1965. *Flora of Japan*. Smithsonian Institution Press, Washington, D.C.
- Oliveira, U. R., F. D. S. Do Espirito Santo, and I. A. Alvarez. 2015. Epiphytic community of *Syagrus coronata* (Mart.) Becc. (Arecaceae) in areas of pastures in Caatinga, Bahia. *Revista Caatinga* 28:84-91.
- Oliver, W. R. B. 1930. New Zealand epiphytes. *Journal of Ecology* 18:1-50.
- Øllgaard, B. 1992. Neotropical Lycopodiaceae-An Overview. *Annals of the Missouri Botanical Garden* 79:687-717.
- Øllgaard, B. 1994. Family 26. Lycopodiaceae. *Fieldiana Botany*:16-88.
- Omlor, R. 1996. Notes on Marsdenieae (Asclepiadaceae): A new, unusual species of *Hoya* from Northern Borneo. *Novon* 6:288-294.
- Omlor, R. 1998. *Generische Revision der Marsdenieae (Asclepiadaceae)*. Shaker Verlag.
- Orihuela, R. L. L., M. Molz, and J. L. Waechter. 2014. Links between environmental factors and hemiepiphytes along a slope of subtropical Atlantic forest. *Nordic Journal of Botany* 32:358-368.

- Ortiz, O. O., and T. B. Croat. 2017. Five new species of *Anthurium* (Araceae) from the Caribbean Forests of Panama. *Webbia* 72:71-81.
- Ortiz, O. O., R. M. Baldini, G. Berguido, and T. B. Croat. 2016. New species of *Anthurium* (Araceae) from Chucanti Nature Reserve, eastern Panama. *Phytotaxa* 255:47-56.
- OTS_Plant_Database. <https://archive.tropicalstudies.org/plantdatabase/>.
- Oyuela Torres, G. O., and T. B. Croat. 2015. Tres nuevas especies de *Anthurium*, (Araceae), para Colombia, Tolima, Ibagué, Cañon del Combeima. *Rodriguésia* 66:769-777.
- Pacific_Bulb_Society. 2017. <https://www.pacificbulbsociety.org/>.
- Palmer, D. D. 2002. Hawaii's ferns and fern allies. University of Hawaii Press.
- Panda, S. 2012. Checklist of Ericaceae in Tuensang District of Nagaland, India with special reference to Mt. Saramati. *Journal of Threatened Taxa* 4:3454-3461.
- Pande, P. C., and H. C. Pande. 1993. Systematics and distribution of epiphytic Pteridophytic flora of Kumaun Himalaya. *Indian Fern Journal* 10:17-29.
- Parmentier, I. 2005. Ecology and distribution of Melastomataceae in African rain forest inselbergs. *Biotropica* 37:364-372.
- Parra, M. J., R. Rodriguez, L. Cavieres, L. Munoz-Tapia, and C. Atala. 2015. Latitudinal patterns in pteridophyte distribution of Continental Chile. *Gayana Botanica* 72:58-69.
- Parris, B. S. 1971. Fourth re-examination of the permanent vegetation quadrats on Little Barrier Island. *Tane* 17:159-168.
- Parris, B. S. 1983. A taxonomic revision of the genus *Grammitis* Swartz (Grammitidaceae, Filicales) in New Guinea. *Blumea* 29:13-222.
- Parris, B. S. 1986. Grammitidaceae of peninsular Malaysia and Singapore. *Kew Bulletin* 41:491-517.
- Parris, B. S. 1997. Two new species of *Grammitis* from the Marquesas Islands. *Allertonia* 7:296-303.
- Parris, B. S. 1998. *Chrysogrammitis*, a new genus of Grammitidaceae (Filicales). *Kew Bulletin* 53:909-918.
- Parris, B. S. 2002. New species and new combinations in African Grammitidaceae (Filicales). *Kew Bulletin* 57:423-434.
- Parris, B. S. 2003. The distribution of Grammitidaceae (Filicales) inside and outside Malesia. *Telopea* 10:451-466.
- Parris, B. S. 2005. Grammitidaceae (Pteridophyta) of Mount Jaya, New Guinea, and other montane Malesian localities. *Fern Gazette* 17:183-203.
- Parris, B. S. pers. comm.
- Parris, B. S., and D. R. Given. 1976. A taxonomic revision of the genus *Grammitis* Sw. (Grammitidaceae: Filicales) in New Zealand. *New Zealand Journal of Botany* 14:85-111.
- Parris, B. S., R. S. Beaman, and J. H. Beaman. 1992. The plants of Mount Kinabalu: Ferns and fern allies. Royal Botanic Gardens, Kew.
- Partomihardjo, T., E. Suzuki, and J. Yukawa. 2004. Development and distribution of vascular epiphytes communities on the Krakatau islands, Indonesia. *South Pacific Studies* 25:7-26.
- Paul, A., M. L. Khan, A. Arunachalam, and K. Arunachalam. 2005. Biodiversity and conservation of Rhododendrons in Arunachal Pradesh in the Indo-Burma biodiversity hotspot. *Current Science (Bangalore)* 89:623-634.
- Paul_Shirley_Succulents. 2016. <https://paulshirleysucculents.wordpress.com/>.
- Pedraza-Penalosa, P. 2010. *Disterigma: Ericaceae: Vaccinieae* (Flora Neotropica Monograph, vol 108, New York).
- Pedraza-Penalosa, P. 2015. New blueberry and mortino relatives (Ericaceae) from northwestern Colombia. *PhytoKeys*:33-58.

- Pedraza-Penalosa, P., and J. L. Luteyn. 2011. Andean *Vaccinium* (Ericaceae: Vaccinieae): Seven new species from South America. *Brittonia* 63:257-275.
- Peña-Chocarro, M., C. Espada-Mateos, M. Vera, G. Céspedes, and S. Knapp. 2010. Updated checklist of vascular plants of the Mbaracayú Forest Nature Reserve (Reserva Natural del Bosque Mbaracayú), Paraguay. *Phytotaxa* 12:1-224.
- Pérez-Calix, E. 2008. Familia Crassulaceae. Instituto de Ecología A.C., Centro Regional del Bajío, Pátzcuaro, Michoacán, Mexico.
- Pérez-Farrera, M. A., R. Martínez-Camilo, N. Martínez-Melendez, O. Farrera-Sarmiento, and S. Maza-Villalobos. 2012. Check list of flora of the Mt. Quetzal (Polygon III) of the Triunfo Biosphere Reserve, Chiapas, Mexico. *Botanical Sciences* 90:113-142.
- Perret, M., A. Chautems, R. Spichiger, G. Kite, and V. Savolainen. 2003. Systematics and evolution of tribe Sinningieae (Gesneriaceae): Evidence from phylogenetic analyses of six plastid DNA regions and nuclear ncpGS. *American Journal of Botany* 90:445-460.
- Perrie, L., Brownsey, P.J., Lovis, J.D. 2010. *Tmesipteris horomaka*, a new octoploid species from Banks Peninsula. *New Zealand J. Bot.* 48: 15-29.
- Perrie, L. R., L. D. Shepherd, P. J. Brownsey, J. Larraín, B. Shaw, L. Thouvenot, and M. von Konrat. 2016. Rediscovery and reinstatement of the New Caledonian endemic filmy fern *Hymenophyllum pumilio* Rosenst. *New Zealand Journal of Botany* 54:1-10.
- Perrier de la Bathie, H. 1951. Mélastomatacées. Pages 1-326 in H. Humbert, editor. *Flore de Madagascar et des Comores*. Firmin-Didot, Paris.
- Peru_Checklist. 2019. <https://tropicos.org/projectwebportal.aspx?pagename=Introduction&projectid=5>.
- Pham, V. T., and L. V. Averyanov. 2011. *Hiepia corymbosa* – new genus of Apocynaceae (Asclepiadoideae) from Vietnam. *Turczaninowia* 14:5-10.
- Pham, V. T., and L. V. Averyanov. 2012. New species from Vietnam – *Hoya lockii* (Apocynaceae, Asclepiadoideae). *Taiwania* 57:49-54.
- Philippine_Medicinal_Plants. 2016. <http://www.stuartxchange.org/Abkel.html>.
- Phillipson, P. B., P. P. Lowry II, L. Andriamahefarivo, P. Antilahimena, and C. Birkinshaw. 2010. Floristic inventory of the Ambatovy-Analamay mine site and comparison to other sites in Madagascar. *Malagasy Nature* 3:44-76.
- Phutthai, T., M. Sands, and K. Sridith. 2009. Field surveys of natural populations of *Begonia* L. in Thailand. *Thai Forest Bulletin (Botany)*:186-196.
- Picheansoonthon, C., and P. Wongsuwan. 2011. A new species and a new record of *Hedychium* J. König (Zingiberaceae) from Thailand. *Taiwania* 56:231-239.
- Pichi Sermolli, R. E. G. 1977. *Fragmenta Pteridologiae*—VI. *Webbia* 31:237-259.
- Pichi Sermolli, R. E. G. 1982. *Microgonium erosum* and related species (Hymenophyllaceae) in continental Africa. *Webbia* 35:241-260.
- Piesschaert, F., S. Jansen, S. Huysmans, E. Smets, and E. Robbrecht. 1999. *Chassalia petitiiana* (Rubiaceae-Psychotrieae), an overlooked epiphytic species hidden in the African canopy. *Systematic Botany* 24:315-322.
- Piggott, A. G. 1980. The fern flora of oilpalm plantations in West Malaysia. *British Fern Gazette* 12:93-102.
- Pincheira-Ulbrich, J., J. R. Rau, and C. Smith-Ramirez. 2012. Vascular epiphytes and climbing plants diversity in an agroforestral landscape in southern Chile: a comparison among native forest fragments. *Boletín de la Sociedad Argentina de Botánica* 47:411-426.
- Pipoly, J. J., III, D. M. Kearns, and P. E. Berry. without year. Key to the Species and Sections of *Clusia* in the Venezuelan Guayana <http://www.mobot.org/mobot/research/ven-guayana/clusiaceae/clusia.html>.
- PlantNET. (The NSW Plant Information Network System). Royal Botanic Gardens and Domain Trust, Sydney. <http://plantnet.rbgsyd.nsw.gov.au>.

- Plunkett, G. M., P. P. Lowry, P. Fiaschi, D. G. Frodin, and A. N. Nicolas. 2020. Phylogeny, biogeography, and morphological evolution among and within the Neotropical and Asian clades of *Schefflera* (Araliaceae). *Taxon*.
- Ponce, M. M., C. Del Rio, A. Ebihara, and J. Y. Dubuisson. 2017. Discussion on taxonomy of the fern genera *Crepidomanes* and *Polyphlebium* (Hymenophyllaceae) in Argentina and south-eastern South America, and description of a new local variety for *Crepidomanes pyxidiferum*. *Botany Letters* 164:5-18.
- Porembski, S. 2011. Evolution, diversity, and habitats of poikilohydrous vascular plants. in U. Lüttge, E. Beck, and D. Bartels, editors. *Plant desiccation tolerance*. Springer, Berlin.
- Pos, E. T., and A. D. M. Slegers. 2010. Vertical distribution and ecology of vascular epiphytes in a lowland tropical rain forest of Brazil. *Boletim do Museu Paraense Emilio Goeldi Ciências Naturais*, Belém 5:335-344.
- POWO. 2019. "Plants of the World Online. Facilitated by the Royal Botanic Gardens, Kew. Published on the Internet; <http://www.plantsoftheworldonline.org/>
- Prasanthkumar, M. G., J. Skornickova, M. Sabu, and S. Jayasree. 2005. Conservation priority and phytogeographical significance of *Rhynchanthus longiflorus* Hook. f. (*Zingiberaceae*): A rare, endangered species from Mizo Hills, NE India. *Current Science* 88:977-980.
- Price, M. G. 1983. *Pecluma*, a new tropical American fern genus. *American Fern Journal* 73:109-116.
- Price, M. G. 1990. Four new Asian *Loxogramme*. *American Fern Journal* 80:4-8.
- Pridgeon, A. M., P. J. Cribb, J. M. Chase, and F. N. Rasmussen. 1999-2014. *Genera Orchidacearum*. Volume 1-6. Reprinted edition. Oxford University Press, Oxford.
- Pro. Summer's Web Garden. 2009. <http://fern.la.coocan.jp/>.
- Proctor, G. R. 1985. *Ferns of Jamaica*. British Museum (Natural History), London.
- Proctor, G. R. 1989. *Ferns of Puerto Rico and the Virgin islands*. New York Botanical Garden, New York.
- Profice, S. R., and P. Leitman. 2013. *Clistax bahiensis* (Acanthaceae), a new epiphytic species from Bahia, Brazil. *Phytotaxa* 132:47-52.
- Prósperi, J., C. Edelin, and G. Michaloud. 1995. Modalités d'occupation de l'espace aérien par les héli-épiphytes: Le rôle de la réitération caulinaire. *Revue d'écologie: la terre et la vie* 50:167-187.
- Puff, C. 2007. *Flora of Thailand: Rubiaceae*. https://homepage.univie.ac.at/christian.puff/_FTH-RUB/FTH-RUB_HOME.htm.
- Puff, C., and R. Buchner. 1998. *Lecananthus* and *Leucocodon*, two genera to be added to the tribe Schradereae (Rubiaceae). *Blumea* 43:265-286.
- Quan-Ru, L., M. Guan-Hua, G. Yuan, and X.-C. Zhang. 2008. A taxonomic revision of *Lepisorus* sect. *Hymenophyton* (Polypodiaceae) from China. *Journal of Systematics and Evolution* 46:906-915.
- Rafidah, A. R. 2012. *Rhododendron chamahensis* (Ericaceae), a new species from Peninsular Malaysia. *Blumea* 57:114-115.
- Rahayu, S. 2010. Diversity and conservation strategy of the epiphytic shrub *Hoya multiflora* Blume (Asclepiadaceae) at Gunung Gede Pangrango National Park, Indonesia based on AFLP analysis. http://atbc2010.org/doc/posters/p-15-8_rahayu.pdf
- Rajapaksha, R., C. Fraser-Jenkins, D. Pushpakumara, D. Wijesundara, and B. Parris. 2017. The endemic pteridophyte flora of Sri Lanka: Taxonomy, geographical distribution and conservation status. *Indian Fern Journal* 33:1-36.
- Rakotondrainibe, F. 1999. Pteridophytes of the eastern slope of the Reserve Naturelle Intégrée d'Andohahela, Madagascar: Distribution and floristic analysis - chapter 3. *Fieldiana Zoology*:25-49.

- Rakotondrainibe, F. 2010. Le genre *Rumohra* Raddi (Pteridophyta, Dryopteridaceae) à Madagascar: une espèce et deux combinaisons nouvelles. *Adansonia* 32:217-228, 212.
- Rakotondrainibe, F., A. Jouy, G. Rouhan, L. Bauret, and B. S. Parris. 2018. Taxonomical and nomenclatural novelties in the grammitid ferns (Pteridophyta, Polypodiaceae, Grammitidoideae) from Madagascar. *Adansonia* 40:141-162.
- Rakotondrainibe, F., A. Rasolohery, F. Raharimalala, and D. Florens. 2003. Les Ptéridophytes des forêts denses humides au nord et à l'ouest de la cuvette d'Andapa (Nord-Est de Madagascar): composition floristique et densité des peuplements, gradients de distribution des taxons. Nouveaux résultats d'inventaires biologiques faisant référence à l'altitude dans la région des massifs montagneux de Marojejy et d'Anjanaharibe-Sud. *Recherches pour le Développement, Sciences biologiques* 19:27-68.
- Rakotondrainibe, F., and A. Jouy. 2017. Nouveautés dans le genre *Lomariopsis* Fée (Pteridophyta, Lomariopsidaceae) à Madagascar : trois espèces nouvelles et une espèce africaine nouvellement signalée. *Adansonia* 39:101-110, 110.
- Rakotondrainibe, F., and F. Raharimalala. 1998. The pteridophytes of the Réserve Spéciale d'Anjanaharibe-Sud, Madagascar: Floristic analysis and altitudinal distribution. *Fieldiana Zoology*:17-38.
- Rakotondrainibe, F., and P. H. Hovenkamp. 2012. A new species of *Pyrrosia* (Polypodiaceae) from Madagascar. *Novon* 22:73, 75-77.
- Rakotondrainibe, F., and T. Deroin. 2006. Comparative morphology and rhizome anatomy of two new species of *Zygophlebia* (Grammitidaceae) from Madagascar and notes on the generic circumscription of *Zygophlebia* and *Ceradenia*. *Taxon* 55:145-152.
- Ramírez-Aguirre, E., S. Martén-Rodríguez, and J. F. Ornelas. 2016. Floral variation, nectar production, and reproductive success of two *Drymonia* (Gesneriaceae) species with mixed pollination syndromes. *International Journal of Plant Sciences* 177:469-480.
- Ranker, T. A., and C. Haufler, H. 2008. Biology and evolution of ferns and lycophytes. Cambridge University Press, New York.
- Ranker, T. A., J. M. O. Geiger, S. C. Kennedy, A. R. Smith, C. H. Haufler, and B. S. Parris. 2003. Molecular phylogenetics and evolution of the endemic Hawaiian genus *Adenophorus* (Grammitidaceae). *Molecular Phylogenetics and Evolution* 26:337-347.
- Rao, A. N. 1998. *India arunachalensis* Nageswara Rao (Orchidaceae): A new genus and species from Arunachal Pradesh (India). *Journal of Economic and Taxonomic Botany* 22:701-703.
- Rao, A. R., and P. Khare. 1965. Studies on Indian Hymenophyllaceae VII: Contributions to our knowledge of *Pleuromanes kurzii* (Bedd) Copeland Proceedings of the Indian National Science Academy B 31:165-174.
- Rao, A. R., and U. Sharma. 1966. Studies on Indian Hymenophyllaceae-Part IV: Contributions to our knowledge of *Mecodium badium* Copel Proceedings of the Indian National Science Academy B 32:208-217.
- Raskoti, B. B., and R. Ale. 2011. A new species of *Sunipia* (Orchidaceae) from Nepal. *Phytotaxa* 31:55-58.
- Rauh, W. 1973. *Solanopteris bismarckii* Rau. *Tropische und subtropische Pflanzenwelt* 5:1-38.
- Ravensberg, W. J., and E. Hennipman. 1986. The *Pyrrosia* species formerly referred to *Drymoglossum* and *Saxiglossum* (Filicales, Polypodiaceae). *Leiden Botanical Series* 9:281-310.
- Real_Jardin_Botanica_de_Madrid. without year. Flora de Guinea Ecuatorial: Base de datos de Flora de Guinea. <http://www.floradeguinea.com/index>
- Real_Jardin_Botanico. without year. Biblioteca digital. <https://bibdigital.rjb.csic.es/>

- Reeb, C., H. L. Ranarijaona, and J.-Y. Dubuisson. 2012. Ecological survey of the lycophytes and ferns of the Vohimana Reserve, Madagascar. *Plant Ecology and Evolution* 145:410-418.
- Regalado, J. C. 1990. Revision of *Medinilla* (Melastomataceae) of Borneo. *Blumea* 35:5-70.
- Regalado, J. C. 1995. Revision of Philippine *Medinilla* (Melastomataceae). *Blumea* 40:113-193.
- Reid, J. L., J. M. Chaves-Fallas, K. D. Holl, and R. A. Zahawi. 2016. Tropical forest restoration enriches vascular epiphyte recovery. *Applied Vegetation Science* 19:508-517.
- Renner, S. S. 1986. The neotropical epiphytic Melastomataceae: phytogeographic patterns, fruit types, and floral biology. *Selbyana* 9:104-111.
- Renner, S. S., G. Clausing, N. Cellin, and K. Meyer. 2001. Melastomataceae. *Flora of Thailand* 7:1-95.
- Rhind, P. M. 2010. Plant formations in the St Helena and Ascension BioProvince.
- Rhind, P. M. without year. Plant Formations in the Malayan BioProvince.
<http://www.terrestrial-biozones.net/Paleotropic%20Vegetation/Malay%20Vegetation.pdf>
- Rhind, P. M. without year. Plant formations in the Papuan BioProvince.
<http://www.terrestrial-biozones.net/Paleotropic%20Vegetation/Papuan%20Vegetation.pdf>
- Rhind, P. M. without year. Plant formations in the Philippinean BioProvince.
<http://www.terrestrial-biozones.net/Paleotropic%20Vegetation/Philippinean%20Vegetation.pdf>
- Rhind, P. M. without year. Plant formations in the Papuan BioProvince.
<http://www.terrestrial-biozones.net/Paleotropic%20Vegetation/Papuan%20Vegetation.pdf>
- Ribeiro, J. E. L. d. S. 1999. Flora da Reserva Ducke: guia de identificação das plantas vasculares de uma floresta de terra-firme na Amazônia Central. INPA-DFID, Manaus, Brazil.
- Richardson, A. 1971. Revision of *Louteridium* (Acanthaceae). *Toulane Studies in Zoology and Botany* 17:63.
- Richardson, K. A., R. S. Currah, and S. Hambleton. 1993. Basidiomycetous endophytes from the roots of neotropical epiphytic Orchidaceae. *Lindleyana* 8:127-137.
- Ridley, H. N. 1912. An expedition to Mount Menuang Casing, Selangor. *Journal of the Federated Malay States Museums*.
- Ridley, H. N. 1916. I. Report on the botany of the Wollaston Expedition to Dutch New Guinea, 1912-13. *Transactions of the Linnean Society - Second Series - Botany* IX:1-276.
- Ridley, H. N. 1920. New and rare Malayan Plants. *Journal of the Federated Malay States Museums* 10:128-156.
- Ridley, H. N. 1930. The dispersal of plants throughout the world. L. Reeve & Co, Ashford, Kent.
- Rintz, R. E. 1979. Three new species of Asclepiadaceae from Peninsular Malaya R.E. *Blumea* 25:225-231.
- Rintz, R. E. 1980. The Peninsular Malayan species of *Dischidia* (Asclepiadaceae). *Blumea* 26:81-126.
- Rio, C. D., S. Hennequin, G. Rouhan, A. Ebihara, P. P. Lowry II, J.-Y. Dubuisson, and M. Gaudeul. 2017. Origins of the fern genus *Hymenophyllum* (Hymenophyllaceae) in New Caledonia: Multiple independent colonizations from surrounding territories and limited in situ diversification. *Taxon* 66:1041-1064.

- Roalson, E. H., and W. R. Roberts. 2016. Distinct processes drive diversification in different clades of Gesneriaceae. *Systematic Biology* 65:662-684.
- Robert, Y., J.-M. Tamon, and H. Thomas. 2014. Confirmation of the presence of the fern species: *Lepisorus schraderi* (Mett.) Ching in Reunion island (Polypodiales: Polypodiaceae). *Cahiers scientifiques de l'océan Indien occidental* 5:21-26.
- Rocha, L. C. F., E. d. C. Smidt, M. A. N. Coelho, and L. G. Temponi. 2014. The genus *Anthurium* (Araceae) in Paraná State - Brazil. *Rodriguésia* 65:917-937.
- Rodda, M. 2012. Taxonomy of *Hoya lyi*, *Hoya yuennanensis* and *Hoya mekongensis* (Apocynaceae – Asclepiadoideae). *Edinburgh Journal of Botany* 69:83–93.
- Rodda, M. 2015. Two new species of *Hoya* R.Br. (Apocynaceae, Asclepiadoideae) from Borneo. *PhytoKeys* 53:83-93.
- Rodda, M., and N. S. Juhonewe. 2012. *Hoya vangviengiensis* (Apocynaceae, Asclepiadoideae), a new species from limestone formations of Vang Vieng, Lao PDR. *Webbia* 67:23-27.
- Rodda, M., and T. Nyhuus. 2009. *Hoya danumensis*, a new species of *Hoya* (Apocynaceae, Asclepiadoideae) from Borneo. *Webbia* 64:163-167.
- Rodda, M., N. Simonsson Juhonewe, and A. Lamb. 2015. *Hoya fauziana* (Apocynaceae: Asclepiadoideae), a new species from Sabah, Malaysian Borneo. *Webbia* 70:207-210.
- Rodda, M., N. Simonsson, and L. Wanntorp. 2011. *Hoya wongii* (Apocynaceae, Asclepiadoideae): a new campanulate flowered species from Brunei (Borneo). *Blumea - Journal of Plant Taxonomy and Plant Geography* 56:205-208.
- Rödl-Linder, G. 1990. A monograph of the fern genus *Goniophlebium*. *Blumea* 34:277-423.
- Rödl-Linder, G. 1994. Revision of the fern genus *Polypodiopteris*. *Blumea* 39:365-371.
- Rodríguez-González, A. 2005. Sinopsis de *Neomirandea* Asteraceae: Eupatorieae en Costa Rica. *Lankesteriana* 5:201-210.
- Rogalski, J. M., A. Reis, M. S. dos Reis, and M. H. de Queiroz. 2016. Spatial distribution of Araceae on phorophytes in different areas of submontane rainforest, Santa Catarina Island (SC), Southern Brazil. *Brazilian Journal of Botany*:1-10.
- Rojas-Alvarado, A. 2008a. Three new species in *Asplenium* (Aspleniaceae) from the Neotropics *Métodos en Ecología y Sistemática* 3:29-35.
- Rojas-Alvarado, A. F. 1996a. Aportes a la flora Pteridophyta costarricense. II. Taxones nuevos. *Brenesia* 45:33-50.
- Rojas-Alvarado, A. F. 1996b. Aportes a la flora pteridophyta neotrópica. 1. Notas sobre el género *Niphidium* (Polypodiaceae). *Brenesia* 45-46:27-32.
- Rojas-Alvarado, A. F. 2002. New species, new combinations and new distributions in neotropical species of *Elaphoglossum* (Lomariopsidaceae). *Revista de Biología Tropical* 50:969-1006.
- Rojas-Alvarado, A. F. 2003a. New taxa, new records and redefined concepts in the *Elaphoglossum* sect. *Elaphoglossum* subsec. *Pachyglossa* (Lomariopsidaceae) from Mexico and Central America. *Revista de Biología Tropical* 51:1-32.
- Rojas-Alvarado, A. F. 2003b. Notes on *Elaphoglossum* (Lomariopsidaceae) section *Polytrichia* subsection *Hybrida* in Mexico and Central America. *Revista de Biología Tropical* 51:33-48.
- Rojas-Alvarado, A. F. 2004. Una especie nueva de *Hymenophyllum* y una variedad nueva de *Trichomanes collariatum* Bosch (Filicales: Hymenophyllaceae) en Costa Rica. *Lankesteriana* 4:143-148.
- Rojas-Alvarado, A. F. 2006. Two new species of *Cochlidium* (Polypodiaceae) from Venezuela. *Métodos en Ecología y Sistemática* 2:13-17.
- Rojas-Alvarado, A. F. 2008b. Lista de los helechos (Pteridophyta) y lycófitos (Lycopodiophyta) de la Reserva Biológica Alberto Manuel Brenes, Costa Rica. *Métodos en Ecología y Sistemática* 3:1-12.

- Rojas-Alvarado, A., Tejedor, A., 2016. Tres especies nuevas de *Cyathea* (Cyatheales: Cyatheaceae) para Colombia. *Revista Biodiversidad Neotropical* 6:121-125.
- Rosenstock, E. 1910. Filices costaricenses. *Repertorium novarum specierum regni vegetabilis*, Beihefte 9:67-70.
- Rosenstock, E. 1912. Neue Farne der Insel Lombok. *Mededeelingen van's Rijks-Herbarium*, Leiden 14:31–33.
- Rosenstock, E. 1924. Neue Arten und Abarten brasilianischer Pteridophyten. *Repertorium novarum specierum regni vegetabilis* 20:89-95.
- Rouhan, G., D. H. Lorence, T. J. Motley, J. G. Hanks, and R. C. Moran. 2008. Systematic revision of *Elaphoglossum* (Dryopteridaceae) in French Polynesia, with the description of three new species. *Botanical Journal of the Linnean Society* 158:309-331.
- Rouhan, G., F. Rakotondrainibe, and R. C. Moran. 2007b. *Elaphoglossum nidusoides* (Dryopteridaceae), a new species of fern from Madagascar with an unusual phylogenetic position in the Squamipedia group. *Systematic Botany* 32:227-235.
- Rouhan, G., J. G. Hanks, D. McClelland, and R. C. Moran. 2007a. Preliminary phylogenetic analysis of the fern genus *Lomariopsis* (Lomariopsidaceae). *Brittonia* 59:115-128.
- Rouhan, G., P. H. Labiak, E. Randrianjohany, and F. Rakotondrainibe. 2012. Not so neotropical after all: the grammitid fern genus *Leucotrichum* (Polypodiaceae) is also paleotropical, as revealed by a new species from Madagascar. *Systematic Botany* 37:331-338.
- Roux, J. P. 2001. Conspectus of southern African Pteridophyta: an enumeration of the Pteridophyta of Angola, Botswana, Lesotho, Malawi, Mozambique, Namibia, South Africa (including the Marion Island group), Swaziland, Zambia and Zimbabwe. Project Coordinator, Southern African Botanical Diversity Network, c/o National Botanical Institute, Pretoria.
- Roux, J. P. 2011. The fern genus *Elaphoglossum* section *Lepidoglossa* (Dryopteridaceae) in Africa, Macaronesia, the mid-Atlantic and southern Indian Ocean Islands. *Botanical Journal of the Linnean Society* 165:20-63.
- Roux, J. P., and S. A. B. D. Network. 2003. Swaziland ferns and fern allies. National Botanical Institute, SABONET, Pretoria.
- Rusea, G., K. Claysius, S. Runi, U. Joanes, K. M. H. Maideen, and A. Latiff. 2009. Ecology and distribution of Lycopodiaceae Mirbel in Malaysia. *Blumea* 54:269-271.
- Rutishauser, R. 2016. Evolution of unusual morphologies in Lentibulariaceae (bladderworts and allies) and Podostemaceae (river-weeds): a pictorial report at the interface of developmental biology and morphological diversification. *Annals of Botany* 117:811-832.
- Säid, A. H., S. Hennequin, G. Rouhan, and J. Y. Dubuisson. 2017. Disentangling the diversity and taxonomy of Hymenophyllaceae (Hymenophyllales, Polypodiidae) in the Comoros. *European Journal of Taxonomy* 313.
- Sakai, S., and H. Nagamasu. 1998. Systematic studies of Bornean Zingiberaceae I. *Amomum* in Lambir Hills, Sarawak. *Edinburgh Journal of Botany* 55:45-64.
- Salino, A. 1998. New combinations in *Pecluma* (Polypodiaceae). *Novon* 8:296-297.
- Salino, A., T. E. Almeida, A. R. Smith, A. N. Gómez, H.-P. Kreier, and H. Schneider. 2008. A new species of *Microgramma* (Polypodiaceae) from Brazil and recircumscription of the genus based on phylogenetic evidence. *Systematic Botany* 33:630-635, 636.
- Samain, M.-S. pers. comm.
- Sampaio, A. F., M. G. Coelho de Souza, and M. R. Pietrobon. 2015. Ferns and lycophytes of the oil basin of Urucu, Coari, Amazon, Brazil. *Acta Biológica Catarinense* 2:33-48.
- Sánchez, C. 2001. A new filmy fern species and new unispecific section of *Trichomanes* (Hymenophyllaceae) ("filmy ferns") from Cuba. *Willdenowia* 31:125-127.
- Sandwith. 1931. *Bulletin of Miscellaneous Information* (Royal Gardens, Kew) 11:177.

- Sanín, D. 2018. *Serpocaulon* (Polypodiaceae). Universidad Nacional de Colombia, Bogotá.
- Sanín, D., J. C. Ospina, I. O. Moura, and A. Salino. 2019. A morphometric analysis of *Serpocaulon gilliesii* (Polypodiaceae) reveals a new species for Yungas Montane Forest, S. australe. *Systematic Botany* 44:90-100.
- Sanín, D., L. M. Á. Mejía, J. C. Mancera Santa, N. Castaño Rubiano, and G. G. Ocampo. 2008. Monilofitos y licofitos de la cuenca del río Chinchina (Caldas, Colombia). Clave para géneros y catálogo de las especies. *Revista de la Academia Colombiana de Ciencias* 32:331-352.
- Sastre, C., and J. J. Granville. 1974. Observations phytogéographiques sur les inselbergs du bassin supérieur du Maroni. *Comptes Rendus de la Société de Biogéographie* 444:7-15.
- Sathapattayanon, A., and T. Boonkerd. 2006. Pteridophyte diversity along a gradient of disturbance within mines in Thong Pha Phum District, Kanchanaburi Province. *BRT Research Report* 2549:1-11.
- Sazima, M., S. Buzato, and I. Sazima. 2003. *Dyssochroa viridiflorum* (Solanaceae): a reproductively bat-dependent epiphyte from the Atlantic rainforest in Brazil. *Annals of Botany* 92:725-730.
- Schäfer, H. 2005. Endemic vascular plants of the Azores: an updated list. *Hoppea* 66:275-283.
- Scheidel, A. 2005. Bestandsaufnahme von Epiphyten, Hemiepiphyten, Kletterer und Lianen in und um den Piedras Blancas National Park, Costa Rica. BSc-Arbeit. Universität Graz.
- Schelppe, E. A. C. L. E. 1966. The pteridophyta of Gorongosa Mountain, Southern Mozambique. *Boletim da Sociedade Broteriana* 40:149.
- Schimper, A. F. W. 1888. Die epiphytische Vegetation Amerikas. Gustav Fischer, Jena.
- Schlechter, R. 1911. Die Orchideen von Neu-Deutsch-Guinea - 24. Macodes. *Repertorium Specierum Novarum Regni Vegetabilis Beihefte* 1:70-74.
- Schlechter, R. 1915. Asclepidiaceae Philippinenses II. *Repertorium novarum specierum regni vegetabilis* 13:555-566.
- Schlechter, R. 1916. Neue Asclepiadaceen von Sumatra und Célebes. *Beihefte Botanisches Centralblatt* XXXIV. 2:1-18.
- Schlechter, R. 1919. Die Ericaceen von Neu-Deutsch-Guinea. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie*:137-194.
- Schlechter, R. 1923. *Sobralia pfavii* Schltr., nov. spec. *Repertorium novarum specierum regni vegetabilis*, Beihefte 19:272.
- Schmitt, J. L., and M. N. B. Goetz. 2010. Species richness of fern and lycophyte in an urban park in the Rio dos Sinos basin, Southern Brazil. *Brazilian Journal of Biology* 70:1161-1167.
- Schneider, H., H.-P. Kreier, P. Hovenkamp, and T. Janssen. 2008. Phylogenetic relationships of the fern genus *Christiopteris* shed new light onto the classification and biogeography of drynarioid ferns. *Botanical Journal of the Linnean Society* 157:645-656.
- Schneider, J. V. 2001. Diversity, structure, and biogeography of a successional and mature upper montane rain forest of the Venezuelan Andes. Der Andere Verlag, Osnabrück, Germany.
- Schneider, P. H., and J. L. Schmitt. 2011. Composition, community structure and vertical distribution of epiphytic ferns on *Alsophila setosa* Kaulf., in a semideciduous seasonal forest, Morro Reuter, RS, Brazil. *Acta Botanica Brasílica* 25:557-565.
- Schröter, H. 1938. Monographie der Gattung *Procris*. II. *Repertorium novarum specierum regni vegetabilis* 45:257-300.
- Schuettpelz, E., and D. W. Trapnell. 2006. Exceptional epiphyte diversity on a single tree in Costa Rica. *Selbyana* 27:65-71.
- Schuettpelz, E., C. W. Chen, M. Kessler, J. B. Pinson, G. Johnson, A. Davila, A. T. Cochran, L. Huiet, and K. M. Pryer. 2016. A revised generic classification of vittarioid ferns

- (Pteridaceae) based on molecular, micromorphological, and geographic data. *Taxon* 65:708-722.
- Schultz, G. P. 2005. Vascular flora of the El Eden Ecological Reserve, Quintana Roo, Mexico. *Journal of the Torrey Botanical Society* 132:311-322.
- Schwartsburd, P. B. 2014. The Identity of *Polypodium gyroflexum* (?*Pleopeltis gyroflexa*, Comb. Nov. – Polypodiaceae). *American Fern Journal* 104:16-21, 16.
- Schwartsburd, P. B., and A. R. Smith. 2013. Novelties in *Serpocaulon* (Polypodiaceae). *Journal of the Botanical Research Institute of Texas* 7:85-93.
- Schwartsburd, P. B., C. V. Miranda, and J. Prado. 2016. *Oleandra* (Oleandraceae) in the Brazilian Atlantic Forest. *American Fern Journal* 106:191-205.
- Scoupe, M. 2011. Composition floristique et diversité de la végétation de la zone Est du Parc National de Taï (Côte d'Ivoire). MSc. Université de Genève, Switzerland.
- Seidenfaden, G., and J. J. Wood. 1992. The orchids of peninsular Malaysia and Singapore. Olsen & Olsen, Fredensborg.
- Sen, G. C. 1963. The epiphytic flowering plants of Darjeeling Hills, other than orchids. *Nelumbo* 5:111-115.
- Senterre, B., Chew, M.Y., Chung, R.C., 2015. Flora and vegetation of Pulau Babi Tengah, Johor, Peninsular Malaysia. *Check List* 11:1714.
- Senterre, B., G. Rouhan, C. Morel, and J. Y. Dubuisson. 2017. New species and records in the fern genus *Didymoglossum* for the flora of Seychelles, with notes on the Southeast Asian *D. motleyi* complex (Hymenophyllaceae). *Phytotaxa* 292:201-217.
- Senterre, B., J. Gerlach, J. Mougil, and D. Matatiken. 2009. Old growth mature forest types and their floristic composition along the altitudinal gradient on Silhouette Island (Seychelles) - the telescoping effect on a continental mid-oceanic island. *Phytocoenologia* 39:157-174.
- Shaikh, S. D., and M. Dongare. 2010. Effects of mining on the diversity of the pteridophytes from the Western Ghats of Maharashtra (India). *World Applied Sciences Journal* 11:1547-1551.
- Shalimov, A. P., and A. I. Shmakov. 2014. To the taxonomy of the fern genus *Polypodiodes Ching* (Polypodiaceae). *Turczaninowia* 17:12-19.
- Shalisko, V., M. A. Sundue, A. R. Villalobos-Arambula, M. A. Muniz-Castro, and J. A. Vazquez-Garcia. 2019. Taxonomic novelties in grammitid ferns (Polypodiaceae) from the Neotropics and Madagascar supported by molecular data. *Phytotaxa* 394:176.
- Shao, W., and S.-G. Lu. 2011. *Himalayopteris*, a new fern genus from India and the adjacent Himalayas (Polypodiaceae, Polypodioideae). *Novon* 21:90-93.
- Shaw, J. 2008. Three new Crûg Farm introductions. *The Plantsman* 7:39-42.
- Sherff, E. E. 1950. Botanical Leaflets, a Series of Studies in the Systematic Botany of Miscellaneous Dicotyledonous Plants 2:5.
- Shi, J. P., and H. Zhu. 2009. Tree species composition and diversity of tropical mountain cloud forest in the Yunnan, southwestern China. *Ecological Research* 24:83-92.
- Shu-Gang, L. U. 1999. A new species of the genus *Polypodiodes Ching* (Polypodiaceae) from China. *Journal of Systematics and Evolution* 37:294-296.
- Sigel, E. M., M. D. Windham, C. H. Haufler, and K. M. Pryer. 2014. Phylogeny, divergence time estimates, and phylogeography of the diploid species of the *Polypodium vulgare* complex (Polypodiaceae). *Systematic Botany* 39:1042-1055.
- Silva, A. G. D., and P. B. Schwartsburd. 2017. Ferns of Viçosa, Minas Gerais State, Brazil: Polypodiaceae (Polypodiales, Filicopsida, Tracheophyta). *Hoehnea* 44:251-268.
- Silva, A., Junior, and J. F. S. Rorig. 2001. Estudo florístico-ecológico das pteridófitas da localidade de Picada Verao, Sapiranga - RS. *Pesquisas Botanica* 51:137-145.

- Silva, I. A., A. W. C. Ferreira, M. I. S. Lima, and J. J. Soares. 2010. Networks of epiphytic orchids and host trees in Brazilian gallery forests. *Journal of Tropical Ecology* 26:127-137.
- Simmonds, N. W. 1950. Notes on the biology of the Araceae of Trinidad. *Journal of Ecology* 38:277-291.
- Simmons, J. B. The Malesian Rhododendrons at Kew
<http://scholar.lib.vt.edu/ejournals/JARS/v23n4/v23n4-simmons.htm>.
- Singh, B., V. N. Singh, S. J. Phukan, B. K. Sinha, and S. K. Borthakur. 2012. Contribution to the pteridophytic flora of India: Nokrek Biosphere Reserve, Meghalaya. *Journal of Threatened Taxa* 4:2277-2294.
- Sledge, W. A. 1956. CXI - The Ceylon species of *Leptochilus*. *Journal of Natural History* 9:865-877.
- Sledge, W. A. 1968. The Hymenophyllaceae of Ceylon. *Journal of the Linnean Society of London, Botany* 60:289-308.
- Sleumer, H. 1934. Ericaceae americanae novae vel minus cognitae. *Notizblatt des Königl. botanischen Gartens und Museums zu Berlin* 12:119-140.
- Sleumer, H. 1957. Florae Malesianae praecursores XIV - A revision of the genus *Diplycosia* (Ericaceae). *Reinwardtia* 4:119-161.
- Sleumer, H. 1966. Ericaceae. *Flora Malesiana* 6:468-668.
- Sleumer, H. Ericaceae in *Flora Malesiana*,
http://www.archive.org/stream/floramalesiana64stee/floramalesiana64stee_djvu.txt.
- Smith, A. C. 1946. Studies of South American plants, XI. Noteworthy species of Hippocrateaceae and Vacciniaceae. *Journal of the Arnold Arboretum* 27:86-120.
- Smith, A. C. 1953. Studies of South American Plants. XIV. *American Journal of Botany* 40:469-475.
- Smith, A. R. 1990. Pteridophytes of the Venezuelan Guayana: New Species. *Annals of the Missouri Botanical Garden* 77:249-273.
- Smith, A. R. 1992. A review of the fern genus *Micropolypodium* (Grammitidaceae). *Novon* 2:419-425.
- Smith, A. R. 1993. New species and combinations in *Ceradenia* (Grammitidaceae). *Novon* 3:182-185.
- Smith, A. R., and J. D. Tejero-Díez. 2014. *Pleopeltis* (Polypodiaceae), a redefinition of the genus and nomenclatural novelties. *Botanical Sciences* 92:43-58.
- Smith, A. R., and M. Kessler. 2018a. Prodrum of a fern flora for Bolivia. III. Selaginellaceae. *Phytotaxa* 334:248-258.
- Smith, A. R., and M. Kessler. 2018b. Prodrum of a fern flora for Bolivia. XXXIII. Blechnaceae. *Phytotaxa* 334:99-117.
- Smith, A. R., B. León, H. Tuomisto, H. van der Werff, R. C. Moran, M. Lehnert, and M. Kessler. 2005. New records of pteridophytes for the flora of Peru. *SIDA, Contributions to Botany* 21:2321-2342.
- Smith, A. R., M. Kessler, B. Leon, T. E. Almeida, I. Jimenez-Perez, and M. Lehnert. 2018. Prodrum of a fern flora for Bolivia. XL. Polypodiaceae. *Phytotaxa* 354:1-67.
- Smith, J. F., and C. L. Carroll. 1997. A cladistic analysis of the tribe Episcieae (Gesneriaceae) based on *ndhF* sequences: Origin of morphological characters. *Systematic Botany* 22:713-724.
- Smith, R. M. 1987. A review of Bornean Zingiberaceae: III (Hedychieae). *Notes of the Royal Botanical Garden of Edinburgh* 44:202-232.
- Solano, G. R. 2011. Three new species of *Stelis* (Orchidaceae) from Mexico. *Brittonia* 63:325-333.

- Sonke, B., L. Zapfack, and D. C. Folefack. 2001. Distribution des epiphytes vasculaires sur les haies vives dans la region de Bafou (Cameroun). *Systematics and Geography of Plants* 71:209-222.
- Specht, C. D. 2006. Systematics and evolution of the tropical monocot family Costaceae (Zingiberales): A multiple dataset approach. *Systematic Botany* 31:89-106.
- Species_foundation_Rhododendrons. http://www.rhododendron.dk/rsf_ac.htm.
- Spooner, D. M., and K. J. Sytsma. 1992. Reexamination of series relationships of Mexican and Central American wild potatoes (*Solanum* sect. *Petota*): Evidence from chloroplast DNA restriction site variation. *Systematic Botany* 17:432-448.
- Spooner, D. M., V. Martínez, R. Hoekstra, and R. G. van den Berg. 1995. Report of expedition to collect wild species of potato in Guatemala USDA.
- Standley, P. C., L. Williams, and D. N. Gibson. 1974. Flora of Guatemala. *Fieldiana Botany* 24:http://www.archive.org/stream/floraofguatemala241034stan/floraofguatemala241034stan_djvu.txt.
- Stearn, W. T. 1954. *Lilium arboricola*, Kingdon Ward's epiphytic Burmese lily. *Garden Chronicle* 136:126-127.
- Steinhard, K. G. 1989. Kakteen und andere Sukkulente, Volumes 40-41.
- Stewart, T., J. P. Biseau, V. Cawoy, and V. Droisart. 2014. Taxonomy of Atlantic Central African orchids 3. A new species of *Bulbophyllum* Thouars (Orchidaceae) from the Monts de Cristal, Gabon. *Phytotaxa* 177:26-34.
- Stevens, P. F. 2004. New taxa in *Paphia* and *Dimorphanthera* (Ericaceae) in Papuaia and the problem of generic limits in *Vaccinieae*. *Edinburgh Journal of Botany* 60:267-298.
- Stevens, P. F. 1990. Family 233: Ericaceae.
- Steyermark, J. A. 1953-1957. Botanical exploration in Venezuela Iii, Iv *Fieldiana Botany* 28:http://www.archive.org/stream/contributionstof2822stey/contributionstof2822stey_djvu.txt.
- Steyermark, J. A. 1974. The summit vegetation of Cerro Autana. *Biotropica* 6:7-13.
- Steyermark, J. A., P. E. Berry, B. K. Holst, and K. Yatskievych, editors. 1995. Flora of the Venezuelan Guayana: Pteridophytes, Spermatophytes Acanthaceae-Araceae. Timber Press.
- Sugden, A. M., and R. J. Robins. 1979. Aspects of the ecology of vascular epiphytes in two Colombian cloud forests, I. The distribution of the vascular flora. *Biotropica* 11:173-188.
- Sugumaran, M. 2010. A systematic study of *Fagraea* sensu lato and a revision of the peninsular Malaysian taxa. Ph.D. thesis. University of Malaya.
- Sulasmi, E. S., R. S. Wicaksono, Sulisetijono, A. N. Rochmah, H. Baidlowi, K. Murdianto, and K. Mawaddah. 2020. Morphological and anatomical characteristic of epiphyte ferns in Ronggo Soerjo Botanical Forest. Page 040030 in AIP Conference Proceedings. AIP Publishing LLC.
- Sundue, M. A. 2007. Two new species of *Pleopeltis* (Polypodiaceae) from Andean South America. *American Fern Journal* 97:107-112, 106.
- Sundue, M. A. 2010. A monograph of *Ascogrammitis*, a new genus of grammitid ferns (Polypodiaceae). *Brittonia* 62:357-399.
- Sundue, M. A. 2014. *Mycopteris*, a new Neotropical genus of grammitid ferns (Polypodiaceae). *Brittonia* 66:174-185.
- Sundue, M. A. 2017a. *Ceradenia spectabilis* (Polypodiaceae), a new species from Cerro del Torra, Colombia. *American Fern Journal* 107:193-199.
- Sundue, M. A. 2017b. Three new species of grammitid ferns (Polypodiaceae) from the fern hunter's paradise: Sierra Juárez, Oaxaca, Mexico. *Systematic Botany* 42:160-168, 169.
- Sundue, M. A., B. S. Parris, T. A. Ranker, A. R. Smith, E. L. Fujimoto, D. Zamora-Crosby, C. W. Morden, W.-L. Chiou, C.-W. Chen, G. Rouhan, R. Y. Hirai, and J. Prado. 2014.

- Global phylogeny and biogeography of grammitid ferns (Polypodiaceae). *Molecular Phylogenetics and Evolution* 81:195-206.
- Sundue, M. A., P. H. Labiak, J. Mostacero, and A. R. Smith. 2012. *Galactodenia*, a new genus of grammitid ferns segregated from *Terpsichore* (Polypodiaceae). *Systematic Botany* 37:339-346.
- Sundue, M. A., W. L. Testo, and T. A. Ranker. 2015. Morphological innovation, ecological opportunity, and the radiation of a major vascular epiphyte lineage. *Evolution* 69:2482-2495.
- Sundue, M., I. Olivares, and M. Kessler. 2018. *Ascogrammitis lehnertii* (Polypodiaceae): A new and dominant understory-species from a diverse community of Grammitid ferns in the Andes of Ecuador. *Systematic Botany* 43:709-716.
- Sutherland, C. H. N. 2001. Plantas descritas originalmente de Honduras y sus nomenclaturas equivalentes actuales. *Ceiba* 42:1-71.
- Sykes, W. R., and J. C. Game. 1996. *Phymatosorus* (Polypodiaceae) in the Cook Islands. *New Zealand Journal of Botany* 34:143-146.
- Synnott, T. J. 1985. A checklist of the flora of Budongo Forest Reserve, Uganda, with notes on ecology and phenology. Dept. of Forestry, Commonwealth Forestry Institute, University of Oxford, Oxford, UK.
- Sytsma, K. J. 1987. The shrubby Gentian genus *Macrocarpaea* in Panama. *Annals of the Missouri Botanical Garden* 74:310-313.
- Szlachetko, D. L., and M. Kolanowska. 2014. Notes on the taxonomy of the genus *Cyrtochilum* (Orchidaceae, Oncidieae) and description of two new species of the *Cyrtochilum divaricatum*-alliance from Colombia. *Wulfenia* 21:55-62.
- Tagawa, M. 1943. *Loxogramme* of Japan. *Acta Phytotaxonomica et Geobotanica* 13:118-129.
- Tagawa, M., and K. Iwatsuki. 1967. Enumeration of Thai pteridophytes collected during 1965-66. *South East Asian Studies*: 23-120.
- Takeuchi, W. 2007. Some notes on Ericaceae from recent expeditions to New Guinea summit environments. *Harvard Papers in Botany* 12:163-171.
- Takeuchi, W. 2008. *Vaccinium obatapaquini* (Ericaceae), a new species from limestone environments in the Southern Fold Belt, Papua New Guinea. *Harvard Papers in Botany* 13:273-275.
- Tamura, M. N., M. Ogisu, and J.-m. Xu. 1997. *Heteropolygonatum*, a New Genus of the Tribe Polygonateae (Convallariaceae) from West China. *Kew Bulletin* 52:949-956.
- Tan, J., Wang, Y., Wu, W., Li, Y.Q., Zhao, W.Y., Liao, W.B., Fan, Q., Liao, B.Y. 2020. The complete genome sequence of *Sorbus insignis* (Rosaceae: Amygdaloideae), an epiphytic shrub in this genus. *Mitochondrial DNA Part B-Resources* 5:3257-3258.
- Tanaka, N., T. Ohi-Toma, and J. Murata. 2010. A new species of *Argostemma* (Rubiaceae) from Mount Victoria, Myanmar. *Blumea* 55:65-67.
- Tavares, A. R., P. Giampaoli, J. Machado, S. Kanashiro, E. P. Chu, and P. B. Pita. 2015. Structural analysis of the foliar epidermis during acclimatization of *Aechmea blanchetiana* (Bromeliaceae) in vitro cultured. *Horticultura Brasileira* 33:45-50.
- Taylor, C. M. 1994a. Revision of *Hillia* (Rubiaceae). *Annals of the Missouri Botanical Garden* 81:571-609.
- Taylor, C. M. 2001. Overview of the neotropical genus *Notopleura* (Rubiaceae: Psychotrieae), with the description of some new species. *Annals of the Missouri Botanical Garden* 88:478-515.
- Taylor, C. M. without year. *Didymochlamys* Hook.f.
<https://www.mobot.org/MOBOT/Research/taylor/didymochlamys.shtml>.
- Taylor, C. M., and R. E. Gereau. 2011. *Rubiacearum Americanarum Magna Hama Pars XXVI*: New species of *Hoffmannia* (Hameliaceae) and more comments on the genus. *Novon* 21:94-117.

- Taylor, N., G. Olsthoom, D. Zappi, G. Khew, and D. Quandt. 2014. A remarkable new *Rhipsalis* (Cactaceae) from eastern Brazil. *Bradleya* 32:2-12.
- Taylor, P. 1994b. The genus *Utricularia* - A taxonomic monograph. Royal Botanic Gardens, Kew, London.
- Teixeira, M. D. R., P. Fiaschi, and A. M. Amorim. 2014. Flora of Bahia: Marcgraviaceae. *Sitientibus série Ciências Biológicas* 13:10.13102/scb13219.
- Tenorio, V., C. M. Sakuragui, and R. C. Vieira. 2012. Stem anatomy of *Philodendron* Schott (Araceae) and its contribution to the systematics of the genus. *Plant Systematics and Evolution* 298:1337-1347.
- Testo, W. L., A. R. Field, E. B. Sessa, and M. Sundue. 2019. Phylogenetic and morphological analyses support the resurrection of *Dendroconche* and the recognition of two new genera in Polypodiaceae subfamily Microsoroideae. *Systematic Botany* 44:737-752, 716.
- The_Gesneriad_Reference_Web. <https://gesneriads.info/>.
- Timberlake, J., F. Dowsett-Lemaire, J. Bayliss, T. Alves, S. Baena, C. Bento, K. Cook, J. Francisco, T. Harris, P. Smith, and C. de Sousa. 2009. MT Nambuli, Mozambique: Biodiversity and conservation. Royal Botanic Gardens, Kew, London.
- Tindale, M. D., and S. K. Roy. 2002. A cytotaxonomic survey of the Pteridophyta of Australia. *Australian Systematic Botany* 15:839-937.
- Tomlinson, P. B. 1986. The botany of mangroves. Cambridge University Press, Cambridge.
- Tomlinson, P. B., J. B. Fisher, F. Halle, and R. Villalobos. 2005. Development of woody branch attachments in *Schefflera* (Araliaceae or Apiaceae). *American Journal of Botany* 92:1765-1773.
- Triana-Moreno, L. A. 2011. Novedades en Pecluma (Polypodiaceae). *Brittonia* 63:62.
- Tropicos. Missouri Botanical Garden. 22 May 2019 <<http://www.tropicos.org>>.
- Trusty, J. L., H. C. Kesler, and G. H. Delgado. 2006. Vascular flora of Isla del Coco, Costa Rica. *Proceedings of the California Academy of Sciences* 57:247-355.
- Tryon, R. M. 1989a. Pteridophyta of Peru. *Fieldiana Botany* 22:1-124.
- Tryon, R. M. 1989b. Pteridophyta of Peru. Part I. 1. Ophioglossaceae-12. Cyatheaceae. *Fieldiana Bot NS* 20:111-138.
- Tryon, R. M. 1993. Pteridophyta of Peru: 18. Aspleniaceae-21. Polypodiaceae. Part V. Field Museum of Natural History.
- Tryon, R. M., and R. G. Stolze. 1991. Pteridophyta of Peru. Part IV. 17. Dryopteridaceae. *Fieldiana, Botany* 27:1-176.
- Tsutsumi, C., and M. Kato. 2006. Evolution of epiphytes in Davalliaceae and related ferns. *Botanical Journal of the Linnean Society* 151:495-510.
- Tsutsumi, C., C. W. Chen, A. Larsson, Y. Hirayama, and M. Kato. 2016. Phylogeny and classification of Davalliaceae on the basis of chloroplast and nuclear markers. *Taxon* 65:1236-1248.
- Turner, I. M. 1995. A catalogue of the vascular plants of Malaya. *Gardens' Bulletin, Singapore* 47:1-757.
- Turner, I. M., F. W. Xing, and R. T. Corlett. 2000. An annotated check-list of the vascular plants of the South China Sea and its shores. *Raffles Bulletin of Zoology* 8:23-116.
- Tutul, E., M. Z. Uddin, M. O. Rahman, and M. A. Hassan. 2010. Angiospermic flora of Runtia sal forest, Bangladesh. II. Magnoliopsida (dicots). *Bangladesh Journal of Plant Taxonomy* 17:33-53.
- Uhl, C. H., R. Moran, and M. Kimnach. 2005. Chromosomes and hybrids of *Echeveria* Dc. IX. series *Spicatae* (Baker) Berger. *Haseltonia*:138-149.
- Ule, E. 1904. Epiphyten des Amazonasgebietes. *Vegetationsbilder* 2.
- University_of_Connecticut. 2019. Biodiversity Education & Research Greenhouses. <http://titanarum.uconn.edu/198500305.html>.

- Valadares, R. T., and C. M. Sakuragui. 2014. A new species of *Anthurium* (Araceae) sect. *Urospadix* subsect. *Obscureviridia* from Espirito Santo, Eastern Brazil. *Systematic Botany* 39:31-35.
- Valdivia, P. E. 1977. Estudio botánico y ecológico de la región del Río Uxpanapa, Veracruz. No 4. Las epífitas. *Biotica* 2:55-81.
- Van Alderwerelt van Rosenburgh, C. R. W. K. 1916. The Aniboina Pteridophyta collected by C. B. Robinson. *Philippine Journal of Science* 11:101-124.
- van den Berghe, E. P., and I. G. d. van den Berghe. 2008. Orchids of Nicaragua.
- van den Brink, R. C. B. 1965. Flora of Java. P. Noordhoff, Groningen.
- van der Bosch, R. B. 1861. *Hymenophyllaceae Javanicae*. Van der Post, Amsterdam.
- Van Velzen, H. P., W. H. Wijtzes, and M. Kappelle. 1993. List of secondary vegetation species of the Pacific montane belt, Cordillera de Talamanca, Costa Rica. *Brenesia* 39-40:147-161.
- Vander Kloet, S. P. 1977. Contribution to the Papuan flora: Notes on the ecology and taxonomy of the *Vaccinioideae* of the Bulldog Road, Morose District Papua New Guinea. *Proceedings of the Nova Scotian Institute of Science* 28:105-117.
- Vander Kloet, S. P., T. A. Dickinson, and W. Strickland. 2003. From Nepal to Formosa, a much larger foot print for *Vaccinium* sect. *Aethopus*. *Acta Botanica Yunnanica* 25:1-24.
- Varadarajan, G. S., and E. Forero. 1987. Genus *Pitcairnia* (Bromeliaceae): New taxa from Colombia, nomenclatural changes, and species complexes. *Systematic Botany* 12:410-415.
- Vasco, A. 2011. Taxonomic revision of *Elaphoglossum* subsection *Muscosa* (Dryopteridaceae). *Blumea* 56:165-202.
- Vasco, A., J. Loriga, G. Rouhan, B. A. Ambrose, and R. C. Moran. 2015. Divided leaves in the genus *Elaphoglossum* (Dryopteridaceae): A phylogeny of *Elaphoglossum* section *Squamipedia*. *Systematic Botany* 40:46-55.
- Vasco, A., R. C. Moran, and G. Rouhan. 2009a. Circumscription and phylogeny of the *Elaphoglossum ciliatum* group (E. sect. *Lepidoglossa*, Dryopteridaceae) based on cpDNA sequences. *Taxon* 58:825-834.
- Vasco, A., R. C. Moran, and G. Rouhan. 2009b. Monograph of the *Elaphoglossum ciliatum* group (Dryopteridaceae). *Brittonia* 61:241-272.
- Vasques, D. T., A. Ebihara, and M. Ito. 2017. The felt fern genus *Pyrrosia* Mirbel (Polypodiaceae): a new subgeneric classification with a molecular phylogenetic analysis based on three plastid markers. *Acta Phytotaxonomica et Geobotanica* 68:65-82.
- Vasques, D. T., A. Ebihara, R. Y. Hirai, J. Prado, and I. Motomi. 2019. Phylogeny of *Hymenophyllum* subg. *Mecodium* (Hymenophyllaceae), with special focus on the diversity of the *Hymenophyllum polyanthos* species complex. *Plant Systematics and Evolution* 305:811-825.
- Vásquez, R. C. 2003. *Zygostates riefenstahliae* (Orchidaceae) – a new species from an isolated dry inter-Andean valley in Bolivia. *Revista de la Sociedad Boliviana de Botánica* 4:41-44.
- Vásquez, R., P. L. Ibsch, and B. Gerkmann. 2003. Diversity of Bolivian Orchidaceae - a challenge for taxonomic, floristic and conservation research. *Organisms Diversity & Evolution* 3:93-102.
- Vazquez-Garcia, J. A., G. Nieves-Hernandez, J. Padilla-Lepe, A. T. Nuno-Rubio, and M. Chazaro-Basanez. 2014. *Echeveria munizii* (Crassulaceae) a new species of epiphyte from tropical Volcan de Colima, Mexico. *Phytotaxa* 191:165-171.
- Vazquez-Perez, N., A. Mendoza-Ruiz, and B. Perez-Garcia. 2012. Morphogenesis of the sexual phase of seven epiphytic taxa of *Polypodium* (Polypodiaceae) from Mexico *Acta Botánica Mexicana* 98:5-21.

- Velayos, M., C. Aedo, F. Cabezas, and M. d. I. Estrella. 2008. Oleandraceae. Pages 1-9 in M. Velayos, editor. Flora de Guinea Ecuatorial, online.
- Veldkamp, J. F. 1978. The *Medinilla myrtilloides*-alliance (Melastomataceae). *Blumea* 24:447-454.
- Venter, F., and A. Arihafa. 2015. Vascular plants. Pages 14-43 in S. J. Richards and N. Whitmore, editors. A Rapid Biodiversity Assessment of Papua New Guinea's Hindenburg Wall Region. Wildlife Conservation Society, Goroka.
- Venter, H. J. T., and R. L. Verhoeven. 2001. Diversity and relationships within the Periplocoideae (Apocynaceae). *Annals of the Missouri Botanical Garden* 88:550-568.
- Verma, D., S. K. Singh, B. S. Kholia, B. K. Sinha, and S. Panday. 2014. Pteridophytes of Khawnglung Wildlife Sanctuary, Mizoram, India. *Indian Journal of Forestry* 37:219-234.
- Vibrans, A. C., A. Bonnet, E. Cagliioni, A. L. d. Gasper, and D. V. Lingner. 2013. Inventário florístico florestal de Santa Catarina: epífitos vasculares da Floresta Ombrófila Densa. Universidade Regional de Blumenau, Blumenau.
- Vicent, M., J. M. Gabriel y Galán, and E. B. Sessa. 2017. Phylogenetics and historical biogeography of *Lomaridium* (Blechnaceae: Polypodiopsida). *Taxon* 66:1304-1316.
- Villagrán, C., R. Rodríguez, and E. Barrera. 1986. Pteridophyta del Archipiélago de Chiloé: Localidades interesantes para Chile. *Gayana* 43:3-14.
- Villaverde, C. S. 1987. Tres nuevas especies de Hymenophyllaceae para Cuba. *Revista del Jardín Botánico Nacional* 8:9-16.
- von Mueller, B. F. 1886. New *Vacciniae* from New Guinea. *Journal of Botany, British and Foreign* 24:289-291.
- Voss, R., M. Turner, M. Fisher, and R. Cort. 1980. Floral biology of *Markea neurantha* Hemsley (Solanaceae), a bat-pollinated epiphyte. *American Midland Naturalist* 103:262-268.
- Vumba_Nature. <http://www.vumba-nature.com/>.
- Waechter, J. L. 1986. Epífitos vasculares da mata paludosa do Faxinal, Torres, RS, Brasil. *Iheringia, Serie Botanica* 34:39-50.
- Wagner, W. H., Jr. 1972. *Solanopteris brunei*, a little known fern epiphyte with dimorphic stems. *American Fern Journal* 62:33-43.
- Wagner, W. L. 1999. Manual of the flowering plants of Hawai'i. University of Hawai'i at Manoa.
- Wallace, B. J. 1981. The Australian vascular epiphytes: flora and ecology. Ph.D. University of New England.
- Walters, G., G. Dauby, T. Stevart, S. Dessein, R. Niangadouma, and O. Lachenaud. 2011. *Novitates Gabonenses* 80. Additions and corrections to the flora of Gabon. *Edinburgh Journal of Botany* 68:423-442.
- Wanek, W., W. Huber, S. K. Arndt, and M. Popp. 2002. Mode of photosynthesis during different life stages of hemiepiphytic *Clusia* species. *Functional Plant Biology* 29:725-732.
- Wanntorp, L., A. Kocyan, R. van Donkelaar, and S. S. Renner. 2006. Towards a monophyletic *Hoya* (Marsdenieae, Apocynaceae): Inferences from the chloroplast trnL region and the rbcL-atpB spacer. *Systematic Botany* 31:586-596.
- Wanntorp, L., and P. I. Forster. 2007. Phylogenetic relationships between *Hoya* and the monotypic genera *Madangia*, *Absolmsia*, and *Micholitzia* (Apocynaceae, Marsdenieae): Insights from flower morphology. *Annals of the Missouri Botanical Garden* 94:36-55.
- Wardle, P., and A. H. MacRae. 1966. Biological Flora of New Zealand - 1. *Weinmannia racemosa* Linn. f. (Cunoniaceae). *Kamahi. New Zealand Journal of Botany* 4:114-131.

- Wasserthal, L. T. 1997. The pollinators of the Malagasy star orchids *Angraecum sesquipedale*, *A. sororium* and *A. compactum* and the evolution of extremely long spurs by pollinator shift. *Botanica Acta* 110:343-359.
- Watkins Jr., J. E., and R. C. Moran. 2019. Gametophytes of the fern genera *Dracoglossum* and *Lomariopsis* (Lomariopsidaceae) and their phylogenetic significance. *International Journal of Plant Science* 180:1004-1015.
- Wattana, S. 2001. A new species of *Agapetes* (Ericaceae) from Thailand. *Edinburgh Journal of Botany* 58:423-427.
- Wattana, S. 2010. The genus *Diplycosia* Blume (Ericaceae) in Thailand. *Thai Journal of Botany* 2:127-132.
- Wattana, S., and C. Trisonthi. 1999. *Agapetes inopinata* Airy Shaw (Ericaceae), a new record for Thailand. *Thai Forest Bulletin (Botany)* 27:19-23.
- Watts, J. L., R. C. Moran, and J. E. Watkins, Jr. 2019. *Hymenasplenium volubile*: documentation of its gametophytes and the first record of a hemiepiphyte in the Aspleniaceae. *Annals of Botany* 124:829-835.
- WCSP. 2012. World Checklist of Selected Plant Families. Facilitated by the Royal Botanic Gardens, Kew. Published on the Internet; <http://apps.kew.org/wcsp/> Retrieved Feb 2, 2012.
- WCSP. 2014. World Checklist of Selected Plant Families. Facilitated by the Royal Botanic Gardens, Kew. Published on the Internet; <http://apps.kew.org/wcsp/> Retrieved Dec 2014.
- WCSP. 2019. World Checklist of Selected Plant Families. Facilitated by the Royal Botanic Gardens, Kew. Published on the Internet; <http://apps.kew.org/wcsp/> Retrieved September 2019.
- Weatherby, C. A. 1939. The group of *Polypodium polypodioides*. *Contributions from the Gray Herbarium of Harvard University* 124:22-35.
- Weber, A., and L. E. Skog. 2007 onw. The genera of Gesneriaceae. Basic information with illustration of selected species. Ed. 2. <http://www.genera-gesneriaceae.at>.
- Webster, G. L., and R. M. Rhode. 2001. Plant diversity of an Andean cloud forest - checklist of the vascular flora of Maquipucuna, Ecuador. University of California Press, Berkeley.
- Wee-Lek, C. 1963. A revision of the genus *Poikilospermum*. *Garden's Bulletin Singapore* 20:1-103.
- Wegner, C., M. Wunderlich, M. Kessler, and M. Schawe. 2003. Foliar C:N ratio of ferns along an Andean elevational gradient. *Biotropica* 35:486-490.
- Wei, X.-P., and X.-C. Zhang. 2013. Species delimitation in the fern genus *Lemmaphyllum* (Polypodiaceae) based on multivariate analysis of morphological variation. *Journal of Systematics and Evolution* 51:485-496.
- Weissflog, A., J. Moog, W. Federle, M. Werner, R. Hashim, and U. Maschwitz. 1999. *Hoya mitrata* Kerr (Asclepiadaceae): a new myrmecotrophic epiphyte from southeast Asia with unique multileaved domatium. *Ecotropica* 5:221-225.
- Went, F. W. 1895. Über Haft- und Nährwurzeln bei Kletterpflanzen und Epiphyten. *Annales du Jardin Botanique de Buitenzorg* 12:1-72.
- Went, F. W. 1940. Soziologie der Epiphyten eines tropischen Regenwaldes. *Annales du Jardin Botanique de Buitenzorg* 50:1-98.
- Werner, F., J. Homeier, and S. R. Gradstein. 2005. Diversity of vascular epiphytes on isolated remnant trees in the montane forest belt of southern Ecuador. *Ecotropica* 11:21-40.
- Wessels Boer, J. G. 1962. The new world species of *Trichomanes* sect. *Didymoglossum* and *Microgonium*. *Acta Botanica Neerlandica* 11:277-330.

- Wester, S., G. Mendieta Leiva, L. Nauheimer, W. Wanek, H. Kreft, and G. Zotz. 2011. Diversity and biogeography of vascular epiphytes at Río Changuinola, Panama Flora 206:66–79.
- Wiehler, H. 1975. *Rufodorsia*, a new Central-American genus in the Gesneriaceae. *Selbyana* 1:138-149.
- Wilbur, R. L. 1975. Synopsis of Costa Rican species of *Burmeistera* (Campanulaceae: Lobelioideae). *Bulletin of the Torrey Botanical Club* 102:225-231.
- Wilbur, R. L. 1976. Flora of Panama - IX. Family 183. - Campanulaceae. *Annals of the Missouri Botanical Garden* 63:593-655.
- Wilbur, R. L., and J. L. Luteyn. 2008. A synopsis of the Mexican and Central American species of *Vaccinium* (Ericaceae). *Journal of the Botanical Research Institute of Texas* 2:207-241.
- Wilder, G. J. 1986. Anatomy of first-order roots in the Cyclanthaceae (Monocotyledoneae). I. Epidermis, cortex, and pericycle. *Canadian Journal of Botany* 64:2622-2644.
- Williams, C. B., and S. C. Sillett. 2007. Epiphyte communities on redwood (*Sequoia sempervirens*) in northwestern California. *Bryologist* 110:420-452.
- Williams, K. J., W. J. Kress, and T. Htun. 2003. A striking new epiphytic *Hedychium* (Zingiberaceae) from Myanmar with a discussion on some anomalous related genera. *Edinburgh Journal of Botany* 60:43-48.
- Williams, L. O. 1975. Tropical American Plants, XVII, http://www.archive.org/stream/tropicalamerican3610will/tropicalamerican3610will_djvu.txt. *Fieldiana Botany* 36.
- Windisch, P. G. 1992. *Trichomanes crispum* L. (Pteridophyta, Hymenophyllaceae) and allied species. *Bradea* VI:78-117.
- Windisch, P. G. 2014. Hymenophyllaceae (Polypodiopsida) no estado do Rio Grande do Sul. *Pesquisas, Botânica* 65:15-48.
- Winter, K., B. J. Wallace, G. C. Stocker, and Z. Roksandic. 1983. Crassulacean acid metabolism in Australian vascular epiphytes and some related species. *Oecologia* 57:129-141.
- Wolf, J. H. D., and A. Flamenco-S. 2006. Vascular epiphytes and their potential as a conservation tool in Pine-oak forests of Chiapas, Mexico. Pages 375-391 in M. Kappelle, editor. *Ecology and Conservation of Neotropical montane oak forests*. Springer, Berlin.
- Wong, K. M., and J. B. Sugau. 1996. A revision of *Fagraea* (Loganiaceae) in Borneo, with notes on related Malesian species and 21 new species. *Sandakania* 8:1-93.
- Wright, A. E., and E. K. Cameron. 1985. Botanical features of northeastern Great Barrier island. *Journal of the Royal Society of New Zealand* 15:251-278.
- Wunderlin, R. P., B. F. Hansen, A. R. Franck, and F. B. Essig. 2019. Atlas of Florida Plants (<http://florida.plantatlas.usf.edu/>). [S. M. Landry and K. N. Campbell (application development), USF Water Institute.] Institute for Systematic Botany, University of South Florida, Tampa.
- Wursten, B., J. Timberlake, and I. Darbyshire. 2017. The Chimanimani Mountains: an updated checklist. *Kirkia* 19:70-100.
- Xu, C.-D., and S.-G. Lu. 2007. *Arthromeris cyrtomioides* S. G. Lu & C. D. Xu, a new species of Polypodiaceae from Yunnan, China. *Acta Phytotaxonomica Sinica* 45:82-85.
- Xu, H. Q., and W. Y. Liu. 2005. Species diversity and distribution of epiphytes in the montane moist evergreen broad-leaved forest in Ailao Mountain, Yunnan. *Biodiversity Science* 13:137-147.
- Young, R. P. 2008. A biodiversity assessment of the Centre Hills, Montserrat. Durrell Wildlife Conservation Trust, Jersey, Channel Islands.

- Yuyen, Y., and T. Boonkerd. 2002. Pteridophyte flora of Huai Yang waterfall National Park, Prachuap Khiri Khan province, Thailand. 2:39-49.
- Zapfack, L., and S. Engwald. 2008. Biodiversity and spatial distribution of vascular epiphytes in two biotopes of the Cameroonian semi-deciduous rain forest. *Plant Ecology* 195:117-130.
- Zhang, D. 1999. Systematics of *Burmannia* L. (Burmanniaceae) in the Old World. Hong Kong University, Hong Kong.
- Zhang, S.-B., Z.-J. Guan, M. Sun, J.-J. Zhang, K.-F. Cao, and H. Hu. 2012. Evolutionary association of stomatal traits with leaf vein density in *Paphiopedilum*, Orchidaceae. *PLoS ONE* 7:e40080
- Zhou, X. M., L. Zhang, C. W. Chen, C. X. Li, Y. M. Huang, D. K. Chen, N. T. Lu, D. Ciczuzza, R. Knapp, T. T. Luong, J. H. Nitta, X. F. Gao, and L. B. Zhang. 2017. A plastid phylogeny and character evolution of the Old World fern genus *Pyrrosia* (Polypodiaceae) with the description of a new genus: *Hovenkampia* (Polypodiaceae). *Molecular Phylogenetics and Evolution* 114:271-294.
- Zona, S., and M. J. M. Christenhusz. 2015. Litter-trapping plants: filter-feeders of the plant kingdom. *Botanical Journal of the Linnean Society* 179:554-586.
- Zou, L.-H., Z.-J. Liu, and J.-X. Huang. 2014. *Vanda malipoensis*, a new species of *Vanda* (Orchidaceae: Epidendroideae; Vandeeae) from China: evidence from morphological and molecular phylogenetic analyses. *Phytotaxa* 186:87-96.
- Zubek, S., K. Piatek, P. Naks, W. Heise, M. Wayda, and P. Mleczko. 2010. Fungal root endophyte colonization of fern and lycophyte species from the Celaque National Park in Honduras. *American Fern Journal* 100:126-136.
- Zuloaga, F. O., O. Morrone, and M. J. Belgrano. without year. Catalogo de las plantas vasculares de cono sur; <http://www.darwin.edu.ar/Proyectos/FloraArgentina/fa.htm>.

V.G. History of data set usage

V.G.1. Data request history: N/A

V.G.2. Data set update history: to be updated continuously

V.G.3. Review history: N/A

V.G.4. Questions and comments from secondary users:

Acknowledgements:

We thank the following colleagues for their advice on particular taxa: Frank Almeda, Robbin Moran, David Neill, Barbara Parris, Charlotte Taylor, Alen Eskov, Diego Giraldo-Cañas, Cheng-Wei Chen, Michael Sundue, and Weston Testo. See also additional acknowledgements in Zotz (2013).

Literature Cited:

- Allard, D. J., M. Petru, and R. R. Mill. 2005. An ecological study of *Pedicularis dendrothauma*, an arboreal hemiparasitic epiphyte from Nepal. *Folia Geobotanica* 40:135-149.

- Balslev, H., R. Valencia, G. Paz y Miño, H. Christensen, and I. Nielsen. 1998. Species count of vascular plants in one hectare of humid lowland forest in Amazonian Ecuador. Pages 585-594 in F. Dallmeier and J. A. Comiskey, editors. Forest biodiversity in North, Central and South America, and the Caribbean. Parthenon Publishing Group, Paris.
- Brazilian Flora 2020 in construction. continuously updated. - available in: <http://floradobrasil.jbrj.gov.br/>, Rio de Janeiro Botanical Garden.
- Burns, K. C. 2008. Meta-community structure of vascular epiphytes in a temperate rainforest. *Botany* **86**:1252-1259.
- Burns, K. C. 2010. How arboreal are epiphytes? A null model for Benzing's classifications. *New Zealand Journal of Botany* **48**:185 - 191.
- Cascante-Marín, A. and A. Estrada-Chavarría. 2012. Las plantas vasculares de El Rodeo, Costa Rica. *Brenesia* **77**:71-128.
- Cascante-Marín, A. and A. Nivia-Ruiz. 2013. Neotropical flowering epiphyte diversity: local composition and geographic affinities. *Biodiversity and Conservation* **22**:113-125.
- Coode, M. J. E. 2005. *Elaeocarpus* for Flora Malesiana: *E. crenulatus*, *E. myrtoides* and *E. amabilis* from New Guinea, reconsidered. *Kew Bulletin* **60**:305-311.
- Costa, L. M. S., M. Goetze, A. V. Rodrigues, G. Dubal, S. Seger, and F. Bered. 2020. Global rheophytes data set: angiosperms and gymnosperms. *Ecology*:e03056.
- Crayn, D. M., K. Winter, and J. A. C. Smith. 2004. Multiple origins of crassulacean acid metabolism and the epiphytic habit in the Neotropical family Bromeliaceae. *Proceedings of the National Academy of Sciences of the United States of America* **101**:3703-3708.
- da Fonseca-da-Silva, T. L., J. Lovo, D. C. Zappi, M. F. Moro, E. D. Leal, C. Maurity, and P. L. Viana. 2020. Plant species on Amazonian canga habitats of Serra Arqueada: the contribution of an isolated outcrop to the floristic knowledge of the Carajas region, Para, Brazil. *Brazilian Journal of Botany* **43**:315-330.
- Derroire, G., L. Schmitt, J. N. Riviere, J. M. Sarrailh, and J. Tassin. 2007. The essential role of tree-fern trunks in the regeneration of *Weinmannia tinctoria* in rain forest on Reunion, Mascarene Archipelago. *Journal of Tropical Ecology* **23**:487-492.
- Díaz, I. A., K. E. Sieving, M. E. Peña-Foxon, J. Larraín, and J. J. Armesto. 2010. Epiphyte diversity and biomass loads of canopy emergent trees in Chilean temperate rain forests: A neglected functional component. *Forest Ecology and Management* **259**:1490-1501.
- Engemann, K., B. Sandel, B. Boyle, B. J. Enquist, P. M. Jørgensen, J. Kattge, B. J. McGill, N. Morueta-Holme, R. K. Peet, N. J. Spencer, C. Violle, S. K. Wiser, and J.-C. Svenning. 2016. A plant growth form dataset for the New World. *Ecology* **97**:3243-3243.
- Fall, P. L. and T. D. Drezner. 2020. Vascular plant species of the Kingdom of Tonga by vegetation type, species origin, growth form, and dispersal mechanism. *Ecology* **101**. e02902
- Flora of China. 2008. Published on the Internet <http://www.efloras.org> [last accessed January 2020] Missouri Botanical Garden, St. Louis, MO & Harvard University Herbaria, Cambridge, MA.
- Gentry, A. H. and C. H. Dodson. 1987. Diversity and biogeography of Neotropical vascular epiphytes. *Annals of the Missouri Botanical Garden* **74**:205-233.
- Gentry, H. S. 2004. Agaves of continental North America. University of Arizona Press.
- Giraldo-Cañas, D. 2018. Circunscripción morfológica, diversidad, patrones de distribución y catálogo de la familia neotropical Marcgraviaceae (Ericales). *Biota Colombiana* **19**:49-69.
- Givnish, T. J., M. H. J. Barfuss, B. V. Ee, R. Riina, K. Schulte, R. Horres, P. A. Gonsiska, R. S. Jabaily, D. M. Crayn, J. A. C. Smith, K. Winter, G. K. Brown, T. M. Evans, B. K.

- Holst, H. Luther, W. Till, G. Zizka, P. E. Berry, and K. J. Sytsma. 2014. Adaptive radiation, correlated and contingent evolution, and net species diversification in Bromeliaceae. *Molecular Phylogenetics and Evolution* **71**:55–78.
- Givnish, T. J., D. Spalink, M. Ames, S. P. Lyon, S. J. Hunter, A. Zuluaga, W. J. D. Iles, M. A. Clements, M. T. K. Arroyo, J. Leebens-Mack, L. Endara, R. Kriebel, K. M. Neubig, W. M. Whitten, N. H. Williams, and K. M. Cameron. 2015. Orchid phylogenomics and multiple drivers of their extraordinary diversification. *Proceedings of the Royal Society of London B: Biological Sciences* **282**: 20151553.
- Grigore, M.-N., L. Ivanescu, and C. Toma. 2014. *Halophytes: An integrative anatomical study*. Springer International Publishing Switzerland.
- Hargis, H., S. G. Gotsch, P. Porada, G. W. Moore, B. Ferguson, and J. T. Van Stan. 2019. Arboreal epiphytes in the soil-atmosphere interface: How often are the biggest “buckets” in the canopy empty? *Geosciences* **9**:342.
- Hassler, M. and B. Schmitt. 2019. Checklist of ferns and lycophytes of the world, <https://worldplants.webarchiv.kit.edu/ferns/>, accessed June 23, 2020.
- Hoeber, V., M. Klinghardt, and G. Zotz. 2020. Drought resistance does not explain epiphytic abundance of accidental epiphytes. *Plant Ecology & Diversity* **13**:175-187.
- Hoeber, V., T. Weichgrebe, and G. Zotz. 2019. Accidental epiphytism in the Harz Mountains, Central Europe. *Journal of Vegetation Science* **30**:765-775.
- Hofstede, R. G. M., K. J. M. Dickinson, and A. F. Mark. 2001. Distribution, abundance and biomass of epiphyte-lianoid communities in a New Zealand lowland *Nothofagus*-podocarp temperate rain forest: tropical comparisons. *Journal of Biogeography* **28**:1033-1049.
- Ibisch, P. L. 1996. *Neotropische Epiphytendiversität - das Beispiel Bolivien*. Martina Galunder-Verlag, Wiehl, Germany.
- Irume, M. V., M. de Lourdes da Costa Soares Morais, C. E. Zartman, and I. L. do Amaral. 2013. Floristic composition and community structure of epiphytic angiosperms in a terra firme forest in central Amazonia. *Acta Botanica Brasílica* **27**:378-393.
- Kelly, D. L., E. V. J. Tanner, E. M. Nic Lughadha, and V. Kapos. 1994. Floristics and biogeography of a rain forest in the Venezuelan Andes. *Journal of Biogeography* **21**:421-440.
- Khine, P. K., C. Fraser-Jenkins, S. Lindsay, D. Middleton, G. Miehe, P. Thomas, and J. Kluge. 2017. A contribution toward the knowledge of ferns and lycophytes from Northern and Northwestern Myanmar. *American Fern Journal* **107**:219-256.
- Kress, W. J. 1986. The systematic distribution of vascular epiphytes: An update. *Selbyana* **9**:2-22.
- Kuijt, J. 1963. On the ecology and parasitism of the Costa Rican tree mistletoe, *Gaiadendron punctatum* (Ruiz & Pavon) G.Don. *Canadian Journal of Botany* **41**:927-938.
- Lens, F., S. Dressler, S. Vinckier, S. Janssens, S. Dessein, L. Van Evelghem, and E. Smets. 2005. Palynological variation in balsaminoid Ericales. I. Marcgraviaceae. *Annals of Botany* **96**:1047-1060.
- Linares-Palomino, R., V. Cardona, E. I. Hennig, I. Hensen, D. Hoffmann, J. Lendzion, D. Soto, S. K. Herzog, and M. Kessler. 2009. Non-woody life-form contribution to vascular plant species richness in a tropical American forest. *Plant Ecology* **201**:87-99.
- Madison, M. 1977. Vascular epiphytes: their systematic occurrence and salient features. *Selbyana* **2**:1-13.
- Martínez-Quezada, D. M., S. Arias, N. Korotkova, and T. Terrazas. 2020. The phylogenetic significance of the stem morpho-anatomy in the Hylocereeae (Cactoideae, Cactaceae). *Plant Systematics and Evolution* **306**:8.
- Massa, G. W. 1996. *Factors affecting the distribution of a neotropical hemiepiphyte*. MSc. San Jose State University, San Jose.

- McCracken, S. F. and M. R. J. Forstner. 2014. Herpetofaunal community of a high canopy tank bromeliad (*Aechmea zebrina*) in the Yasuní Biosphere Reserve of Amazonian Ecuador, with comments on the use of “arboreal” in the herpetological literature. *Amphibian & Reptile Conservation* **8**:65–75.
- Melnychuk, M. and D. S. Srivastava. 2002. Vertical distribution of a bromeliad-dwelling damselfly larva (*Mecistogaster modesta*) in a Costa Rican rainforest (Odonata: Pseudostigmatidae). *International Journal of Odonatology* **5**:81-97.
- Mota, N. F. d. O., M. T. C. Watanabe, D. C. Zappi, A. L. Hiura, J. Pallos, R. S. Viveros, A. M. Giuliatti, and P. L. Viana. 2018. Cangas da Amazônia: a vegetação única de Carajás evidenciada pela lista de fanerógamas. *Rodriguésia* **69**:1435-1488.
- Munn-Estrada, D. X. 2017. Contribution to the floristic knowledge of the Sierra Mazateca of Oaxaca, Mexico. *Lundellia* **20**:25-59.
- Nadkarni, N. M. and T. J. Matelson. 1991. Fine litter dynamics within the tree canopy of a tropical cloud forest. *Ecology* **72**:2071-2082.
- Nadkarni, N. M. and T. J. Matelson. 1992. Biomass and nutrient dynamics of epiphytic litterfall in a neotropical montane forest, Costa Rica. *Biotropica* **24**:24-30.
- Nguyen, D. V., T. B. Croat, H. T. Luu, C. Lee, J. Lee, and R. de Kok. 2013. Two new species of *Alocasia* (Araceae, Colocasieae) from Vietnam. *Willdenowia* **43**:293 – 299.
- Nieder, J., S. Engwald, M. Klawun, and W. Barthlott. 2000. Spatial distribution of vascular epiphytes (including hemiepiphytes) in a lowland Amazonian rain forest (Surumoni Crane Plot) of Southern Venezuela. *Biotropica* **32**:385-396.
- Nitta, J. H. and M. J. Epps. 2009. Hemi-epiphytism in *Vandenboschia collariata* (Hymenophyllaceae). *Brittonia* **61**:392-397.
- Ogburn, R. M. and E. J. Edwards. 2010. The ecological water-use strategies of succulent plants. Pages 179-225 in J. C. Kader and M. Delseny, editors. *Advances in Botanical Research*.
- Penneys, D. S. and W. S. Judd. 2011. Phylogenetics and morphology in the Blakeaceae (Melastomataceae). *International Journal of Plant Sciences* **172**:78-106.
- Plunkett, G. M., P. P. Lowry, P. Fiaschi, D. G. Frodin and A. N. Nicolas 2020. Phylogeny, biogeography, and morphological evolution among and within the Neotropical and Asian clades of *Schefflera* (Araliaceae). *Taxon* **68**: 1278-1313.
- Poltz, K. and G. Zotz. 2011. Vascular epiphytes on isolated pasture trees along a rainfall gradient in the lowlands of Panama. *Biotropica* **43**:165-172.
- Prescott, G. W., D. P. Edwards, and W. A. Foster. 2015. Retaining biodiversity in intensive farmland: epiphyte removal in oil palm plantations does not affect yield. *Ecology and Evolution* **5**:1944-1954.
- Pridgeon, A. M., P. J. Cribb, J. M. Chase, and F. N. Rasmussen. 1999-2014. *Genera Orchidacearum*. Volume 1-6. Reprinted edition. Oxford University Press, Oxford.
- Ramos, F. N. and S. R. Mortara and N. Monalisa-Francisco and J. P. C. Elias and L. M. Neto and L. Freitas and R. Kersten and A. M. Amorim and F. B. Matos and A. F. Nunes-Freitas and S. Alcantara and M. H. N. Alexandre and R. J. de Almeida-Scabbia and O. J. G. de Almeida and F. E. Alves and R. M. de Oliveira Alves and F. S. Alvim and A. C. S. de Andrade and S. de Andrade and L. Y. S. Aona and A. C. Araujo and K. C. T. de Araújo and V. Ariati and J. C. Assis and C. O. de Azevedo and B. F. Barbosa and D. E. F. Barbosa and F. d. R. Barbosa and F. de Barros and G. A. Basilio and F. A. Bataghin and F. Bered and J. S. Bianchi and C. T. Blum and C. R. Boelter and A. Bonnet and P. H. S. Brancalion and T. B. Breier and C. d. T. Brion and C. R. Buzatto and A. Cabral and T. J. Cadorin and E. Cagliani and L. Canêz and P. H. Cardoso and F. S. de Carvalho and R. G. Carvalho and E. L. M. Catharino and S. J. Ceballos and M. T. Cerezini and R. G. César and C. Cestari and C. J. N. Chaves and V. Citadini-Zanette and L. F. M. Coelho and J. V. Coffani-Nunes and R. Colares and G. D.

Colletta and N. d. M. Corrêa and A. F. da Costa and G. M. da Costa and L. M. S. Costa and N. G. S. Costa and D. R. Couto and C. Cristofolini and A. C. R. da Cruz and L. A. Del Neri and M. di Pasquo and A. dos Santos Dias and L. d. C. D. Dias and R. Dislich and M. C. Duarte and J. R. Fabricante and F. H. A. Farache and A. P. G. de Faria and C. Faxina and M. T. M. Ferreira and E. Fischer and C. R. Fonseca and T. Fontoura and T. M. Francisco and S. G. Furtado and M. Galetti and M. L. Garbin and A. L. de Gasper and M. Goetze and J. Gomes-da-Silva and M. F. A. Gonçalves and D. R. Gonzaga and A. C. G. e. Silva and A. d. C. Guaraldo and E. d. S. G. Guarino and A. V. Guislon and L. B. Hudson and J. G. Jardim and P. Jungbluth and S. d. S. Kaeser and I. M. Kessous and N. M. Koch and Y. S. Kuniyoshi and P. H. Labiak and M. E. Lapate and A. C. L. Santos and R. L. B. Leal and F. S. Leite and P. Leitman and A. P. Liboni and D. Liebsch and D. V. Lingner and J. A. Lombardi and E. Lucas and J. d. R. Luzzi and P. Mai and L. F. Mania and W. Mantovani and A. G. Maragni and M. C. M. Marques and G. Marquez and C. Martins and L. d. N. Martins and P. L. S. S. Martins and F. F. F. Mazziere and C. d. A. Melo and M. M. F. de Melo and A. F. Mendes and L. Mesacasa and L. P. C. Morellato and V. d. S. Moreno and A. Muller and M. M. d. S. Murakami and E. Cecconello and C. Nardy and M. H. Nervo and B. Neves and M. G. C. Nogueira and F. R. Nonato and A. T. de Oliveira-Filho and C. P. L. de Oliveira and G. E. Overbeck and G. M. Marcusso and M. L. B. Paciencia and P. Padilha and P. T. Padilha and A. C. A. Pereira and L. C. Pereira and R. A. S. Pereira and J. Pincheira-Ulbrich and J. S. R. Pires and M. A. Pizo and K. C. Pôrto and L. Rattis and J. R. d. M. Reis and S. G. d. Reis and T. C. da Rocha-Pessôa and C. F. D. Rocha and F. S. Rocha and A. R. P. Rodrigues and R. R. Rodrigues and J. M. Rogalski and R. L. Rosanelli and A. Rossado and D. R. Rossatto and D. C. Rother and C. R. Ruiz-Miranda and F. Z. Saiter and M. B. Sampaio and L. D. Santana and J. S. d. Santos and R. Sartorello and M. Sazima and J. L. Schmitt and G. Schneider and B. G. Schroeder and L. Sevegnani and V. O. S. Júnior and F. R. da Silva and M. J. da Silva and M. P. P. Silva and R. G. Silva and S. M. Silva and R. B. Singer and G. Siqueira and L. E. Soares and H. C. de Sousa and A. Spielmann and V. R. Tonetti and M. T. Z. Toniato and P. S. B. Ulguim and C. van den Berg and E. van den Berg and I. G. Varassin and I. B. V. da Silva and A. C. Vibrans and J. L. Waechter and E. W. Weissenberg and P. G. Windisch and M. Wolowski and A. Yañez and V. N. Yoshikawa and L. R. Zandoná and C. M. Zanella and E. M. Zanin and D. C. Zappi and V. B. Zipparro and J. P. F. Zorzanelli and M. C. Ribeiro. 2019. Atlantic epiphytes: A data set of vascular and non-vascular epiphyte plants and lichens from the Atlantic Forest. *Ecology* **100**:e02541.

Richardson, B. A., M. J. Richardson, F. N. Scatena, and W. H. McDowell. 2000. Effects of nutrient availability and other elevational changes on bromeliad populations and their invertebrate communities in a humid tropical forest in Puerto Rico. *Journal of Tropical Ecology* **16**:167-188.

Schimper, A. F. W. 1888. *Die epiphytische Vegetation Amerikas*. Gustav Fischer, Jena.

Schuettpelz, E., H. Schneider, A. R. Smith, P. Hovenkamp, J. Prado, G. Rouhan, A. Salino, M. Sundue, T. E. Almeida, B. Parris, E. B. Sessa, A. R. Field, A. L. de Gasper, C. J. Rothfels, M. D. Windham, M. Lehnert, B. Dauphin, A. Ebihara, S. Lehtonen, P. B. Schwartzburd, J. Metzgar, L.-B. Zhang, L.-Y. Kuo, P. J. Brownsey, M. Kato, M. D. Arana, F. C. Assis, M. S. Barker, D. S. Barrington, H.-M. Chang, Y.-H. Chang, Y.-S. Chao, C.-W. Chen, D.-K. Chen, W.-L. Chiou, V. A. d. O. Dittrich, Y.-F. Duan, J.-Y. Dubuisson, D. R. Farrar, S. Fawcett, J. M. Gabriel y Galan, L. A. d. A. Goes-Neto, J. R. Grant, A. L. Grusz, C. Haufler, W. Hauk, H. He, S. Hennequin, R. Y. Hirai, L. Huiet, M. Kessler, P. Korall, P. H. Labiak, A. Larsson, B. Leon, C.-X. Li, F.-W. Li, M. Link-Perez, H.-M. Liu, L. Ngan Thi, E. I. Meza-Torres, X.-Y. Miao, R. Moran, C.

- M. Mynssen, N. Nagalingum, B. Ollgaard, A. M. Paul, J. B. d. S. Pereira, L. R. Perrie, M. Ponce, T. A. Ranker, C. Schulz, W. Shinohara, A. Shmakov, E. M. Sigel, F. S. de Souza, L. d. S. Sylvestre, W. Testo, L. A. Triana-Moreno, C. Tsutsumi, H. Tuomisto, I. A. Valdespino, A. Vasco, R. S. Viveros, A. Weakley, R. Wei, S. Weststrand, P. G. Wolf, G. Yatskievych, X.-G. Xu, Y.-H. Yan, L. Zhang, X.-C. Zhang, X.-M. Zhou, and G. Pteridophyte Phylogeny. 2016. A community-derived classification for extant lycophytes and ferns. *Journal of Systematics and Evolution* **54**:563-603.
- Solano, R., H. Huerta-Espinoza, G. Cruz-García, and F. Ortiz-Riveros. 2019. A new natural hybrid in the genus *Laelia* (Orchidaceae) from Oaxaca, Mexico. *Phytotaxa* **402**:232-240.
- Sperotto, P., P. Acevedo-Rodríguez, T. N. C. Vasconcelos, and N. Roque. 2020. Towards a standardization of terminology of the climbing habit in plants. *The Botanical Review*. 1- 31, 10.1007/s12229-020-09218-y
- Stuntz, S., U. Simon, and G. Zotz. 2002. Rainforest airconditioning: the moderating influence of epiphytes on the microclimate in tropical tree crowns. *International Journal of Biometeorology* **46**:53-59.
- Sugden, A. M. and R. J. Robins. 1979. Aspects of the ecology of vascular epiphytes in two Colombian cloud forests, I. The distribution of the vascular flora. *Biotropica* **11**:173-188.
- Van Stan, J. T., II, A. Stubbins, T. Bittar, J. S. Reichard, K. A. Wright, and R. B. Jenkins. 2015. *Tillandsia usneoides* (L.) L. (Spanish moss) water storage and leachate characteristics from two maritime oak forest settings. *Ecohydrology* **8**:988-1004.
- Vidal-Russell, R. and D. L. Nickrent. 2008. The first mistletoes: Origins of aerial parasitism in Santalales. *Molecular Phylogenetics and Evolution* **47**:523-537.
- Watts, J. L., R. C. Moran, and J. E. Watkins, Jr. 2019. *Hymenasplenium volubile*: documentation of its gametophytes and the first record of a hemiepiphyte in the Aspleniaceae. *Annals of Botany* **124**:829-835.
- WCSP. 2019. World Checklist of Selected Plant Families. Facilitated by the Royal Botanic Gardens, Kew. Published on the Internet; <http://apps.kew.org/wcsp/> Retrieved September 2019.
- WCVF. 2020. World Checklist of Vascular Plants, version 2.0. Facilitated by the Royal Botanic Gardens, Kew. Published on the Internet; <http://wcvp.science.kew.org/> Retrieved 17 July 2020.
- Weigelt, P., C. König, and H. Kreft. 2020. GIFT – a global inventory of floras and traits for macroecology and biogeography. *Journal of Biogeography* **47**:16-43.
- Whitmore, T. C., R. Peralta, and K. Brown. 1985. Total species count in a Costa Rican tropical rain forest. *Journal of Tropical Ecology* **1**:375-378.
- Zotz, G. 2002. Categories and CAM - blurring divisions, increasing understanding? *New Phytologist* **156**:4-6.
- Zotz, G. 2013. The systematic distribution of vascular epiphytes – a critical update. *Botanical Journal of the Linnean Society* **171**:453-481.
- Zotz, G. 2016. *Plants on plants. The biology of vascular epiphytes*. Springer International Publishing Switzerland, Cham.
- Zotz, G., A. P. Bautista, J. Kohlstruck, and T. Weichgrebe. 2020. Life forms in aroids - natural variability vs. terminological confusion. *Aroideana* **43**:315-333..
- Zotz, G., P. Hietz, and G. Schmidt. 2001. Small plants, large plants - the importance of plant size for the physiological ecology of vascular epiphytes. *Journal of Experimental Botany* **52**:2051-2056.
- Zotz, G. and S. Schultz. 2008. The vascular epiphytes of a lowland forest in Panama - species composition and spatial structure. *Plant Ecology* **195**:131-141.

Tables:

Table 1. Systematic distribution of epiphytic seed plants (Magnoliopsida and Cycadopsida). For each family and genus, the numbers of epiphytic/hemiepiphytic species (“E/H”) are given as 1) all species in **Epilist 1.0**, as 2) all epiphytic species that are either “accepted” or “ambiguous” in World Flora Online (wfo), and 3) all epiphytic species that are “accepted” in wfo. For the latter two categories the total number of species (“Total”) are given and the percentage of epiphytic species (“%Total”). An “H” marks genera, which have many or only hemiepiphytes. The following example of Acanthaceae clarifies the use of the table. **Epilist 1.0** contains 8 species of this family. Six of them are accepted in wfo, which is equivalent to 0.2 % of the 3944 accepted species. One more epiphytic species is found in wfo as "ambiguous". These 7 accepted / ambiguous epiphytic species (all) represent < 0.1% of the 8920 accepted / ambiguous species listed in wfo. An 8th epiphytic species of **Epilist 1.0** is not found in wfo but accepted in plants of the world online (powo).

Class/family	Genus	E/H	wfo (all)			wfo (accepted)		
			E/H	Total	% Total	E/H	Total	% Total
Magnoliopsida		28227	27144	508447	5.3	25318	303775	8.3
Acanthaceae		8	7	8920	<0.1	6	3944	0.2
	<i>Clistax</i>	1	0	2	<0.1	0	2	<0.1
	<i>Dischistocalyx</i>	2	2	16	12.5	2	13	15.4
	<i>Justicia</i>	1	1	1393	<0.1	1	652	0.2
	<i>Louteridium</i>	1	1	11	9.1	1	9	11.1
	<i>Ruellia</i>	1	1	686	<0.1	1	269	0.4
	<i>Stenandrium</i>	1	1	78	1.3	1	49	2.0
	<i>Stenostephanus</i>	1	1	69	1.4		40	<0.1
Alzateaceae		1	1	1	100	1	1	100
	<i>Alzatea</i>	1	1	1	100	1	1	100
Amaryllidaceae		10	10	2603	0.4	10	2246	0.4
	<i>Clivia</i>	1	1	10	10.0	1	7	14.3
	<i>Cyrtanthus</i>	1	1	62	1.6	1	56	1.8
	<i>Hippeastrum</i>	4	4	106	3.8	4	94	4.3
	<i>Pamianthe</i>	3	3	3	100	3	3	100

	<i>Scadoxus</i>	1	1	13	7.7	1	9	11.1	
Apocynaceae		269	253	9347	2.7	91	5556	1.6	
	<i>Absolmsia</i>	1	1	2	50.0	1	1	100	
	<i>Anatropanthus</i>	1	1	1	100	1	1	100	
	<i>Centrostemma</i>	1	1	1	100		0		
	<i>Conchophyllum</i>	1	1	10	10.0		0		
	<i>Dischidia</i>	36	35	128	27.3	34	116	29.3	
	<i>Dischidiopsis</i>	8	8	9	88.9	8	8	100	
	<i>Epistemma</i>	3	3	4	75.0	2	3	66.7	
	<i>Heynella</i>	1	1	1	100		0		
	<i>Hiepia</i>	1	1	1	100	1	1	100	
	<i>Hoya</i>	208	193	466	41.4	37	51	72.5	
	<i>Mandevilla</i>	4	4	199	2.0	4	191	2.1	
	<i>Micholitzia</i>	1	1	1	100	1	1	100	
	<i>Oreosparte</i>	1	1	1	100		0		
	<i>Physostelma</i>	1	1	5	20.0	1	1	100	
	<i>Sarcorrhiza</i>	1	1	1	100	1	1	100	
Aquifoliaceae		3	3	882	0.3	3	480	0.6	H
	<i>Ilex</i>	3	3	823	0.4	3	475	0.6	H
Araceae		747	717	4670	15.4	685	3366	20.4	
	<i>Anthurium</i>	664	634	1616	39.2	603	926	65.1	
	<i>Arisaema</i>	3	3	190	1.6	3	180	1.7	
	<i>Philodendron</i>	56	56	661	8.5	55	486	11.3	
	<i>Remusatia</i>	4	4	4	100	4	4	100	
	<i>Scindapsus</i>	5	5	37	13.5	5	35	14.3	
	<i>Stenospermaton</i>	14	14	79	17.7	14	54	25.9	
	<i>Syngonium</i>	1	1	40	2.5	1	35	2.9	
Araliaceae		94	94	1981	4.7	94	1533	6.1	H
	<i>Aralia</i>	7	7	99	7.1	7	74	9.5	H
	<i>Dendropanax</i>	1	1	104	1.0	1	99	1.0	H

	<i>Oreopanax</i>	14	14	164	8.5	14	152	9.2	H
	<i>Pentapanax</i>	1	1	2	50.0	1	1	100	
	<i>Polyscias</i>	2	2	184	1.1	2	176	1.1	H
	<i>Pseudopanax</i>	1	1	7	14.3	1	7	14.3	H
	<i>Raukaua</i>	2	2	8	25.0	2	8	25.0	H
	<i>Schefflera</i>	66	66	598	11.0	66	591	11.2	H
Asparagaceae		25	24	3465	0.7	24	2916	0.9	
	<i>Cordyline</i>	1	1	36	2.8	1	26	3.8	
	<i>Drimia</i>	1	1	110	0.9	1	99	1.0	
	<i>Heteropolygonatum</i>	7	6	6	100	6	6	100	
	<i>Hosta</i>	1	1	43	2.3	1	23	4.3	
	<i>Maianthemum</i>	7	7	39	17.9	7	39	17.9	
	<i>Polygonatum</i>	7	7	81	8.6	7	71	9.9	
	<i>Yucca</i>	1	1	96	1.0	1	49	2.0	
Asphodelaceae		1	1	1510	<0.1	1	1232	0.1	
	<i>Dianella</i>	1	1	42	2.4	1	41	2.4	
Asteliaceae		10	9	37	24.3	9	37	24.3	
	<i>Astelia</i>	6	5	26	19.2	5	26	19.2	
	<i>Collospermum</i>	4	4	5	80.0	4	5	80.0	
Asteraceae		38	38	58232	<0.1	37	32704	0.1	
	<i>Aristeguietia</i>	1	1	19	5.3	1	19	5.3	
	<i>Brachyglottis</i>	1	1	43	2.3	1	38	2.6	
	<i>Chromolaena</i>	1	1	180	0.6	1	175	0.6	
	<i>Dahlia</i>	1	1	58	1.7	1	42	2.4	
	<i>Nelsonianthus</i>	1	1	1	100		0		
	<i>Neomirandea</i>	22	22	30	73.3	22	30	73.3	
	<i>Pachythamnus</i>	1	1	1	100	1	1	100	
	<i>Pentacalia</i>	3	3	263	1.1	3	223	1.3	
	<i>Senecio</i>	2	2	3035	<0.1	2	1583	0.1	
	<i>Sinclairia</i>	2	2	32	6.3	2	28	7.1	

	<i>Solanecio</i>	1	1	17	5.9	1	16	6.3
	<i>Tuberostylis</i>	2	2	2	100	2	2	100
Balsaminaceae		35	34	1237	2.7	10	488	2.0
	<i>Impatiens</i>	35	34	1222	2.8	10	487	2.1
Begoniaceae		58	56	1809	3.1	56	1601	3.5
	<i>Begonia</i>	58	56	1795	3.1	56	1600	3.5
Bromeliaceae		1985	1970	3613	54.5	1943	3319	58.5
	<i>Aechmea</i>	277	274	298	91.9	274	283	96.8
	<i>Androlepis</i>	2	2	2	100	2	2	100
	<i>Araeococcus</i>	9	9	9	100	9	9	100
	<i>Billbergia</i>	65	64	94	68.1	64	66	97.0
	<i>Brocchinia</i>	5	5	23	21.7	5	21	23.8
	<i>Bromelia</i>	1	1	68	1.5	1	61	1.6
	<i>Canistrum</i>	22	21	22	95.5	21	21	100
	<i>Catopsis</i>	21	21	24	87.5	21	21	100
	<i>Cipuroopsis</i>	2	2	2	100	2	2	100
	<i>Connellia</i>	1	1	6	16.7	1	6	16.7
	<i>Disteganthus</i>	5	3	3	100	3	3	100
	<i>Fascicularia</i>	1	1	1	100	1	1	100
	<i>Fernseea</i>	1	1	2	50.0	1	2	50.0
	<i>Glomeropitcairnia</i>	2	2	2	100	2	2	100
	<i>Greigia</i>	1	1	36	2.8	1	35	2.9
	<i>Guzmania</i>	203	202	224	90.2	202	218	92.7
	<i>Hohenbergia</i>	44	44	63	69.8	44	62	71.0
	<i>Hohenbergiopsis</i>	1	1	1	100	1	1	100
	<i>Lymania</i>	8	8	9	88.9	8	9	88.9
	<i>Mezobromelia</i>	8	8	9	88.9	8	8	100
	<i>Navia</i>	4	4	92	4.3	4	91	4.4
	<i>Neoregelia</i>	117	116	126	92.1	116	124	93.5
	<i>Nidularium</i>	42	42	60	70.0	42	55	76.4

	<i>Orthophytum</i>	3	3	68	4.4	3	68	4.4	
	<i>Pitcairnia</i>	40	40	421	9.5	40	402	10.0	
	<i>Portea</i>	2	2	8	25.0	2	8	25.0	
	<i>Quesnelia</i>	19	19	23	82.6	19	22	86.4	
	<i>Racinaea</i>	21	21	21	100	0	0		
	<i>Ronnbergia</i>	23	11	11	100	11	11	100	
	<i>Tillandsia</i>	665	659	751	87.7	659	692	95.2	
	<i>Vriesea</i>	316	315	396	79.5	315	361	87.3	
	<i>Wallisia</i>	2	2	NA		0	NA		
	<i>Werauhia</i>	10	10	11	90.9		0		
	<i>Wittmackia</i>	42	41	NA		0	NA		
Burmanniaceae		2	2	165	1.2	2	163	1.2	
	<i>Burmannia</i>	2	2	56	3.6	2	56	3.6	
Burseraceae		1	1	1039	<0.1	1	649	0.2	H
	<i>Bursera</i>	1	1	136	0.7	1	120	0.8	H
Cactaceae		130	129	7672	1.7	119	2230	5.4	
	<i>Disocactus</i>	11	11	25	44.0	11	15	73.3	
	<i>Epiphyllum</i>	12	12	33	36.4	12	15	80.0	
	<i>Hatiora</i>	6	6	12	50.0	6	8	75.0	
	<i>Hylocereus</i>	12	12	20	60.0	12	15	80.0	
	<i>Lepismium</i>	4	4	27	14.8	4	8	50.0	
	<i>Lymanbensonia</i>	3	3	3	100		0		
	<i>Pfeiffera</i>	4	4	8	50.0	3	3	100	
	<i>Pseudorhipsalis</i>	6	6	7	85.7	6	7	85.7	
	<i>Rhipsalidopsis</i>	1	1	2	50.0		0		
	<i>Rhipsalis</i>	39	38	120	31.7	37	52	71.2	
	<i>Schlumbergera</i>	5	5	17	29.4	2	7	28.6	
	<i>Selenicereus</i>	18	18	34	52.9	18	28	64.3	
	<i>Trichocereus</i>	1	1	51	2.0	1	9	11.1	
	<i>Weberocereus</i>	8	8	11	72.7	7	9	77.8	

Calophyllaceae	9	9	528	1.7	9	302	3.0	H
<i>Clusiella</i>	9	9	9	100	9	9	100	H
Campanulaceae	32	31	2703	1.1	31	2383	1.3	
<i>Burmeistera</i>	18	17	106	16.0	17	103	16.5	
<i>Canarina</i>	1	1	4	25.0	1	3	33.3	
<i>Clermontia</i>	12	12	25	48.0	12	24	50.0	
<i>Trematolobelia</i>	1	1	10	10.0	1	8	12.5	
Clusiaceae	70	69	1212	5.6	68	868	7.7	H
<i>Clusia</i>	69	68	347	19.5	67	307	21.9	H
<i>Quapoya</i>	1	1	12	8.3	1	5	20.0	H
Commelinaceae	3	3	810	0.4	3	727	0.4	
<i>Belosynapsis</i>	1	1	6	16.7	1	6	16.7	
<i>Cochliostema</i>	2	2	2	100	2	2	100	
Costaceae	6	6	145	4.1	6	139	4.3	
<i>Costus</i>	6	6	111	5.4	6	105	5.7	
Crassulaceae	30	29	3163	0.9	27	1482	1.8	
<i>Aichryson</i>	2	2	27	7.4	2	18	11.1	
<i>Bryophyllum</i>	1	1	47	2.1	1	46	2.2	
<i>Echeveria</i>	12	11	242	4.5	11	170	6.5	
<i>Kalanchoe</i>	6	6	211	2.8	5	133	3.8	
<i>Sedum</i>	9	9	807	1.1	8	392	2.0	
Cunoniaceae	2	2	593	0.3	2	245	0.8	H
<i>Leiospermum</i>	1	1	3	33.3	1	1	100	H
<i>Weinmannia</i>	1	1	219	0.5	1	117	0.9	H
Cyclanthaceae	60	60	261	23.0	60	231	26.0	
<i>Asplundia</i>	7	7	104	6.7	7	100	7.0	
<i>Chorigyne</i>	7	7	7	100	7	7	100	
<i>Ludovia</i>	2	2	3	66.7	2	3	66.7	
<i>Sphaeradenia</i>	40	40	54	74.1	40	53	75.5	
<i>Stelestylis</i>	3	3	4	75.0	3	4	75.0	

Cyperaceae	<i>Thoracocarpus</i>	1	1	1	100	1	1	100
		1	1	6433	<0.1	1	5783	<0.1
	<i>Coleochloa</i>	1	1	8	12.5	1	8	12.5
Ericaceae		741	729	7241	10.1	531	3553	14.9
	<i>Agapetes</i>	78	78	161	47.9	78	147	52.5
	<i>Anthopteropsis</i>	1	1	1	100	1	1	100
	<i>Anthopterus</i>	7	7	14	50.0	7	12	58.3
	<i>Calopteryx</i>	2	2	2	100	2	2	100
	<i>Cavendishia</i>	82	82	125	65.6	81	113	71.7
	<i>Ceratostema</i>	26	26	73	35.6	26	36	72.2
	<i>Costera</i>	8	8	10	80.0	8	10	80.0
	<i>Demosthenesia</i>	3	3	14	21.4	3	13	23.1
	<i>Didonica</i>	4	4	4	100	4	4	100
	<i>Dimorphanthera</i>	14	14	78	17.9	14	77	18.2
	<i>Diogenesia</i>	4	4	14	28.6	3	13	23.1
	<i>Diplycosia</i>	46	41	120	34.2	40	111	36.0
	<i>Disterigma</i>	30	30	47	63.8	25	38	65.8
	<i>Gaultheria</i>	4	4	165	2.4	4	141	2.8
	<i>Gonocalyx</i>	5	5	11	45.5	5	11	45.5
	<i>Lateropora</i>	2	2	3	66.7	2	3	66.7
	<i>Lebetanthus</i>	1	1	1	100	1	1	100
	<i>Macleania</i>	18	18	43	41.9	18	37	48.6
	<i>Mycerinus</i>	1	1	3	33.3	1	3	33.3
	<i>Oreanthes</i>	7	7	7	100	7	7	100
	<i>Orthaea</i>	15	15	35	42.9	15	34	44.1
	<i>Paphia</i>	2	2	21	9.5		0	
	<i>Plutarchia</i>	1	1	11	9.1	1	10	10.0
	<i>Prionotes</i>	1	1	2	50.0		0	
	<i>Psammisia</i>	35	33	69	47.8	31	45	68.9
	<i>Rhododendron</i>	169	166	1500	11.1	35	641	5.5

	<i>Rusbya</i>	1	1	1	100	1	1	100	
	<i>Satyria</i>	12	10	28	35.7	10	24	41.7	
	<i>Semiramisia</i>	1	1	5	20.0	1	3	33.3	
	<i>Siphonandra</i>	1	1	5	20.0	1	3	33.3	
	<i>Sphyrropermum</i>	19	19	24	79.2	19	24	79.2	
	<i>Themistoclesia</i>	16	16	38	42.1	14	27	51.9	
	<i>Thibaudia</i>	20	20	118	16.9	19	62	30.6	
	<i>Utleya</i>	1	1	1	100	1	1	100	
	<i>Vaccinium</i>	104	104	767	13.6	53	223	23.8	
Gentianaceae		28	24	3044	0.8	27	1688	1.6	H
	<i>Fagraea</i>	25	21	77	27.3	25	72	34.7	H
	<i>Macrocarpaea</i>	3	3	115	2.6	2	75	2.7	
Gesneriaceae		628	626	4912	12.7	616	3108	19.8	
	<i>Acanthonema</i>	1	1	2	50.0	1	2	50.0	
	<i>Aeschynanthus</i>	199	199	201	99.0	194	194	100	
	<i>Agalmyla</i>	91	91	96	94.8	91	96	94.8	
	<i>Alloplectus</i>	6	6	29	20.7	6	6	100	
	<i>Asteranthera</i>	1	1	1	100	1	1	100	
	<i>Codonanthe</i>	21	21	23	91.3	20	20	100	
	<i>Codonanthopsis</i>	4	4	4	100	4	4	100	
	<i>Columnnea</i>	191	191	254	75.2	191	195	97.9	
	<i>Crantzia</i>	1	1	11	9.1	1	2	50.0	
	<i>Cremosperma</i>	2	2	27	7.4	2	23	8.7	
	<i>Cyrtandra</i>	2	2	1043	0.2	2	732	0.3	
	<i>Drymonia</i>	14	14	82	17.1	14	75	18.7	
	<i>Episcia</i>	3	3	42	7.1	2	12	16.7	
	<i>Fieldia</i>	2	2	2	100	2	2	100	
	<i>Glossoloma</i>	5	5	27	18.5	5	27	18.5	
	<i>Heppiella</i>	1	1	4	25.0	1	4	25.0	
	<i>Kohleria</i>	2	2	50	4.0	2	40	5.0	

	<i>Loxostigma</i>	1		7	<0.1		7	<0.1	
	<i>Lysionotus</i>	13	13	35	37.1	13	23	56.5	
	<i>Nematanthus</i>	31	30	45	66.7	30	35	85.7	
	<i>Neomortonia</i>	3	3	3	100	3	3	100	
	<i>Oerstedina</i>	3	3	3	100	3	3	100	
	<i>Oreocharis</i>	1	1	67	1.5	1	55	1.8	
	<i>Paradrymonia</i>	10	10	46	21.7	10	31	32.3	
	<i>Primulina</i>	2	2	121	1.7	2	121	1.7	
	<i>Rufodorsia</i>	4	4	4	100	4	4	100	
	<i>Sarmienta</i>	1	1	2	50.0	1	1	100	
	<i>Sinningia</i>	5	5	87	5.7	4	70	5.7	
	<i>Streptocarpus</i>	5	5	202	2.5	4	134	3.0	
	<i>Trichantha</i>	2	2	5	40.0	2	3	66.7	
	<i>Trichosporum</i>	1	1	88	1.1		0		
Griselinaceae		3	3	7	42.9	3	7	42.9	H
	<i>Griselinia</i>	3	3	7	42.9	3	7	42.9	H
Grossulariaceae		2	2	513	0.4	2	195	1.0	
	<i>Ribes</i>	2	2	431	0.5	2	194	1.0	
Hydrangeaceae		2	2	419	0.5	2	236	0.8	
	<i>Hydrangea</i>	2	2	107	1.9	2	52	3.8	
Lamiaceae		1	1	9390	<0.1	1	7879	<0.1	
	<i>Callicarpa</i>	1	1	187	0.5	1	172	0.6	
Lentibulariaceae		28	28	499	5.6	27	312	8.7	
	<i>Pinguicula</i>	6	6	153	3.9	5	73	6.8	
	<i>Utricularia</i>	22	22	296	7.4	22	218	10.1	
Liliaceae		2	2	834	0.2	2	745	0.3	
	<i>Lilium</i>	2	2	129	1.6	2	111	1.8	
Malvaceae		2	2	9185	<0.1	2	4462	<0.1	H
	<i>Spirotheca</i>	2	2	5	40.0	2	5	40.0	H
Marantaceae		1	1	643	0.2	1	569	0.2	

	<i>Monotagma</i>	1	1	45	2.2	1	39	2.6	
Marcgraviaceae		38	13	184	7.1	35	136	25.7	H
	<i>Marcgravia</i>	9	7	86	8.1	8	61	13.1	H
	<i>Marcgraviastrum</i>	6	1	14	7.1	5	11	45.5	H
	<i>Norantea</i>	1	1	13	7.7	1	7	14.3	H
	<i>Pseudosarcopera</i>	1	1	1	100	1	1	100	H
	<i>Ruyschia</i>	3	1	13	7.7	3	10	30.0	H
	<i>Sarcopera</i>	3	1	7	14.3	3	7	42.9	H
	<i>Schwartzia</i>	10	2	19	10.5	9	18	50.0	H
	<i>Souroubea</i>	5	1	28	3.6	5	21	23.8	H
Melastomataceae		378	376	9704	3.9	214	4077	5.2	
	<i>Acinodendron</i>	1	1	529	0.2	1	491	0.2	
	<i>Bertolonia</i>	1	1	35	2.9	1	9	11.1	
	<i>Blakea</i>	90	90	127	70.9	68	81	84.0	
	<i>Calvoa</i>	7	7	25	28.0	5	9	55.6	
	<i>Clidemia</i>	9	9	341	2.6	8	173	4.6	
	<i>Dicellandra</i>	1	1	5	20.0	1	3	33.3	
	<i>Diplectria</i>	2	2	26	7.7	1	1	100	
	<i>Graffenrieda</i>	2	2	62	3.2	1	54	1.9	H
	<i>Gravesia</i>	2	2	117	1.7	2	109	1.8	
	<i>Guyonia</i>	1	1	2	50.0	1	1	100	
	<i>Kendrickia</i>	1	1	1	100		0		
	<i>Macrocentrum</i>	4	4	25	16.0	4	24	16.7	
	<i>Medinilla</i>	147	146	509	28.7	52	85	61.2	
	<i>Memecylon</i>	1	1	474	0.2		106	<0.1	
	<i>Miconia</i>	9	9	1216	0.7	9	881	1.0	
	<i>Monolena</i>	8	8	16	50.0	8	8	100	
	<i>Pachycentria</i>	8	8	28	28.6		0		
	<i>Phainantha</i>	4	4	5	80.0	4	5	80.0	
	<i>Pleiochiton</i>	12	11	11	100	5	5	100	

	<i>Plethiandra</i>	7	7	11	63.6		0		
	<i>Preussiella</i>	1	1	2	50.0	1	2	50.0	
	<i>Salpinga</i>	3	3	12	25.0	2	9	22.2	
	<i>Topobea</i>	55	55	85	64.7	38	49	77.6	H
	<i>Triolena</i>	2	2	20	10.0	2	20	10.0	
Moraceae		389	389	2002	19.4	383	1217	31.5	H
	<i>Dorstenia</i>	2	2	136	1.5	2	117	1.7	
	<i>Ficus</i>	384	384	983	39.1	380	841	45.2	H
	<i>Pourouma</i>	1	1	38	2.6	1	29	3.4	H
Myrtaceae		3	3	6532	<0.1	3	5970	0.1	H
	<i>Metrosideros</i>	2	2	65	3.1	2	55	3.6	H
	<i>Xanthomyrtus</i>	1	1	25	4.0	1	25	4.0	
Nepenthaceae		30	30	186	16.1	30	186	16.1	
	<i>Nepenthes</i>	30	30	186	16.1	30	186	16.1	
Onagraceae		8	8	1775	0.5	8	828	1.0	
	<i>Fuchsia</i>	8	8	133	6.0	8	113	7.1	
Orchidaceae		21169	21010	31300	67.1	20956	27793	75.4	
	<i>Abdominea</i>	1	1	1	100	1	1	100	
	<i>Acampe</i>	8	8	8	100	8	8	100	
	<i>Acianthera</i>	286	223	224	99.6	221	221	100	
	<i>Acineta</i>	17	17	19	89.5	17	17	100	
	<i>Acriopsis</i>	9	9	9	100	9	9	100	
	<i>Acrochaene</i>	1	1	1	100	1	1	100	
	<i>Acrorchis</i>	1	1	1	100	1	1	100	
	<i>Ada</i>	2	2	2	100	1	1	100	
	<i>Adamantina</i>	1	1	1	100	1	1	100	
	<i>Adenoncos</i>	17	17	17	100	17	17	100	
	<i>Adrorhizon</i>	1	1	1	100	1	1	100	
	<i>Aerangis</i>	58	51	56	91.1	51	53	96.2	
	<i>Aeranthes</i>	45	45	47	95.7	44	45	97.8	

<i>Aerides</i>	35	35	62	56.5	32	32	100
<i>Aeridostachya</i>	1	1	0		1	0	
<i>Aetheorrhyncha</i>	1	1	1	100	1	1	100
<i>Aganisia</i>	4	4	6	66.7	4	4	100
<i>Aglossorrhyncha</i>	13	13	13	100	13	13	100
<i>Agrostophyllum</i>	131	102	104	98.1	102	102	100
<i>Alamania</i>	1	1	1	100	1	1	100
<i>Alatiliparis</i>	5	5	5	100	5	5	100
<i>Ambrella</i>	1	1	1	100	1	1	100
<i>Amesiella</i>	3	3	3	100	3	3	100
<i>Amoana</i>	2		0			0	
<i>Anathallis</i>	165	153	153	100	153	153	100
<i>Ancistrochilus</i>	2	2	2	100	2	2	100
<i>Ancistrorhynchus</i>	18	18	19	94.7	18	18	100
<i>Andinia</i>	73	13	13	100	13	13	100
<i>Angraecopsis</i>	22	22	22	100	22	22	100
<i>Angraecum</i>	219	218	242	90.1	217	221	98.2
<i>Anguloa</i>	1	1	16	6.3	1	13	7.7
<i>Ansellia</i>	1	1	1	100	1	1	100
<i>Anthogonium</i>	1	1	1	100	1	1	100
<i>Appendicula</i>	140	134	142	94.4	134	136	98.5
<i>Arachnis</i>	13	11	12	91.7	11	12	91.7
<i>Archivea</i>	1	1	1	100	1	1	100
<i>Armodorum</i>	4	4	4	100	4	4	100
<i>Arnottia</i>	1	1	4	25.0	1	4	25.0
<i>Arpophyllum</i>	3	3	3	100	3	3	100
<i>Artorima</i>	1	1	1	100	1	1	100
<i>Ascidieria</i>	8	8	8	100	8	8	100
<i>Ascocentrum</i>	13	13	13	100	13	13	100
<i>Ascochilopsis</i>	2	2	2	100	2	2	100

<i>Ascochilus</i>	6	6	7	85.7	6	6	100
<i>Ascoglossum</i>	1	1	1	100	1	1	100
<i>Aspasia</i>	7	7	7	100	7	7	100
<i>Aspidogyne</i>	1	1	46	2.2	1	46	2.2
<i>Barbosella</i>	19	19	20	95.0	19	19	100
<i>Barkeria</i>	16	16	17	94.1	16	17	94.1
<i>Batemannia</i>	5	5	5	100	5	5	100
<i>Beclardia</i>	2	2	2	100	2	2	100
<i>Beloglottis</i>	2	2	7	28.6	2	7	28.6
<i>Benthamia</i>	2	2	38	5.3	2	33	6.1
<i>Benzingia</i>	9	9	9	100	9	9	100
<i>Biermannia</i>	11	11	11	100	11	11	100
<i>Bifrenaria</i>	21	21	28	75.0	21	22	95.5
<i>Bogoria</i>	4	4	4	100	4	4	100
<i>Bolusiella</i>	7	6	6	100	6	6	100
<i>Brachionidium</i>	81	75	78	96.2	75	75	100
<i>Brachypeza</i>	12	7	7	100	7	7	100
<i>Bracisepalum</i>	2	2	2	100	2	2	100
<i>Braemia</i>	1	1	1	100	1	1	100
<i>Brasiliorchis</i>	15	15	15	100	15	15	100
<i>Brassavola</i>	24	22	25	88.0	22	23	95.7
<i>Brassia</i>	68	63	65	96.9	63	63	100
<i>Bromheadia</i>	30	30	30	100	30	30	100
<i>Broughtonia</i>	6	6	9	66.7	6	7	85.7
<i>Bryobium</i>	8	8	8	100	8	8	100
<i>Bulbophyllum</i>	2043	1870	1921	97.3	1866	1884	99.0
<i>Bulleyia</i>	1	1	1	100	1	1	100
<i>Caladenia</i>	1	1	288	0.3	1	285	0.4
<i>Calanthe</i>	15	15	238	6.3	15	220	6.8
<i>Callostylis</i>	5	5	5	100	5	5	100

<i>Caluera</i>	3	3	3	100	3	3	100
<i>Calymmanthera</i>	5	5	5	100	5	5	100
<i>Calypetrochilum</i>	2	2	2	100	2	2	100
<i>Camaridium</i>	83	83	83	100	83	83	100
<i>Campanulorchis</i>	5	5	5	100	5	5	100
<i>Campylocentrum</i>	83	65	67	97.0	65	65	100
<i>Capanemia</i>	9	9	10	90.0	9	10	90.0
<i>Cardiochilos</i>	1	1	1	100	1	1	100
<i>Catasetum</i>	156	156	215	72.6	156	194	80.4
<i>Cattleya</i>	108	108	265	40.8	108	186	58.1
<i>Caucaea</i>	15	9	9	100	9	9	100
<i>Caularthron</i>	4	4	4	100	4	4	100
<i>Centroglossa</i>	5	5	6	83.3	5	6	83.3
<i>Ceratocentron</i>	1	1	1	100	1	1	100
<i>Ceratochilus</i>	1	1	1	100	1	1	100
<i>Ceratostylis</i>	148	147	147	100	147	147	100
<i>Chamaeangis</i>	11	11	11	100	11	11	100
<i>Chamaeanthus</i>	2	2	2	100	2	2	100
<i>Chamelophyton</i>	1	1	1	100	1	1	100
<i>Chaseella</i>	1	1	1	100	1	1	100
<i>Chaubardia</i>	3	3	3	100	3	3	100
<i>Chaubardiella</i>	8	8	8	100	8	8	100
<i>Chauliodon</i>	1	1	1	100	1	1	100
<i>Cheiradenia</i>	1	1	1	100	1	1	100
<i>Cheirostylis</i>	2	2	55	3.6	2	54	3.7
<i>Chelonistele</i>	15	13	13	100	13	13	100
<i>Chilopogon</i>	2	2	2	100	2	2	100
<i>Chiloschista</i>	20	20	21	95.2	20	20	100
<i>Chondrorhyncha</i>	7	7	8	87.5	7	7	100
<i>Chondroscaphe</i>	14	14	14	100	14	14	100

<i>Christensonella</i>	17	17	17	100	17	17	100
<i>Chroniochilus</i>	5	5	5	100	5	5	100
<i>Chrysoglossum</i>	3	3	4	75.0	3	4	75.0
<i>Chysis</i>	10	10	14	71.4	10	10	100
<i>Chytroglossa</i>	3	3	3	100	3	3	100
<i>Cirrhaea</i>	7	7	12	58.3	7	7	100
<i>Cischweinfia</i>	11	11	11	100	11	11	100
<i>Claderia</i>	1	1	3	33.3	1	2	50.0
<i>Cleisocentron</i>	6	5	5	100	5	5	100
<i>Cleisomeria</i>	2	2	2	100	2	2	100
<i>Cleisostoma</i>	96	88	96	91.7	88	88	100
<i>Cleisostomopsis</i>	2	2	2	100	2	2	100
<i>Clowesia</i>	7	7	7	100	7	7	100
<i>Cochleanthes</i>	4	4	4	100	4	4	100
<i>Coelia</i>	3	3	5	60.0	3	5	60.0
<i>Coeliopsis</i>	1	1	1	100	1	1	100
<i>Coelogyne</i>	200	200	216	92.6	200	200	100
<i>Comparettia</i>	81	78	79	98.7	78	79	98.7
<i>Conchidium</i>	11	11	10	100	10	10	100
<i>Constantia</i>	6	6	6	100	6	6	100
<i>Cordiglottis</i>	7	7	7	100	7	7	100
<i>Coryanthes</i>	60	60	67	89.6	59	59	100
<i>Corybas</i>	1	1	136	0.7	1	133	0.8
<i>Cottonia</i>	1	1	1	100	1	1	100
<i>Crepidium</i>	9	7	260	2.7	7	260	2.7
<i>Cribbia</i>	4	4	4	100	4	4	100
<i>Crossoglossa</i>	2	2	27	7.4	2	26	7.7
<i>Cryptarrhena</i>	3	3	3	100	3	3	100
<i>Cryptocentrum</i>	20	20	20	100	20	20	100
<i>Cryptochilus</i>	5	5	5	100	5	5	100

<i>Cryptopus</i>	4	4	4	100	4	4	100
<i>Cryptopylos</i>	1	1	1	100	1	1	100
<i>Cryptostylis</i>	1	1	23	4.3	1	23	4.3
<i>Cuitlauzina</i>	8	7	7	100	7	7	100
<i>Cyclopogon</i>	4	4	86	4.7	4	83	4.8
<i>Cycnoches</i>	34	34	36	94.4	34	34	100
<i>Cylindrolobus</i>	16	3	3	100	0	0	
<i>Cymbidiella</i>	2	2	3	66.7	2	3	66.7
<i>Cymbidium</i>	56	56	119	47.1	56	80	70.0
<i>Cynorkis</i>	2	2	168	1.2	2	158	1.3
<i>Cyphochilus</i>	1	1	8	12.5	1	8	12.5
<i>Cypholoron</i>	3	3	3	100	3	3	100
<i>Cypripedium</i>	1	1	240	0.4	1	59	1.7
<i>Cyrtidiorchis</i>	5	5	5	100	5	5	100
<i>Cyrtochiloides</i>	3	3	3	100	3	3	100
<i>Cyrtochilum</i>	179	135	141	95.7	135	138	97.8
<i>Cyrtopodium</i>	37	37	50	74.0	37	48	77.1
<i>Cyrtorchis</i>	19	18	19	94.7	18	18	100
<i>Daiotyla</i>	4	4	4	100	4	4	100
<i>Deceptor</i>	1	1	1	100	1	1	100
<i>Dendrobium</i>	1557	1485	1601	92.8	1484	1523	97.4
<i>Dendrochilum</i>	284	275	279	98.6	275	278	98.9
<i>Dendrolirium</i>	3	1	1	100	0	0	
<i>Dendrophylax</i>	15	14	14	100	14	14	100
<i>Devogelia</i>	1	1	1	100	1	1	100
<i>Diaphananthe</i>	34	23	23	100	22	22	100
<i>Dichaea</i>	122	119	123	96.7	119	120	99.2
<i>Dickasonia</i>	1	1	1	100	1	1	100
<i>Dienia</i>	1	1	6	16.7	1	6	16.7
<i>Dilochia</i>	5	5	8	62.5	5	8	62.5

<i>Dilochiopsis</i>	1	1	1	100	1	1	100
<i>Dilomilis</i>	5	5	5	100	5	5	100
<i>Dimerandra</i>	8	8	8	100	8	8	100
<i>Dimorphorchis</i>	8	5	5	100	5	5	100
<i>Dinema</i>	1	1	1	100	1	1	100
<i>Dinklageella</i>	3	3	4	75.0	3	4	75.0
<i>Diodonopsis</i>	6	5	5	100	5	5	100
<i>Diplocentrum</i>	2	2	2	100	2	2	100
<i>Diploprora</i>	2	2	2	100	2	2	100
<i>Dipodium</i>	22	22	28	78.6	22	24	91.7
<i>Disa</i>	1	1	192	0.5	1	185	0.5
<i>Disperis</i>	1	1	82	1.2	1	79	1.3
<i>Distylodon</i>	2	1	1	100	1	1	100
<i>Domingoa</i>	4	4	5	80.0	4	5	80.0
<i>Draconanthes</i>	2	2	2	100	2	2	100
<i>Dracula</i>	138	128	132	97.0	128	130	98.5
<i>Dresslerella</i>	13	13	16	81.3	13	16	81.3
<i>Dressleria</i>	13	12	12	100	11	11	100
<i>Dryadella</i>	56	54	54	100	54	54	100
<i>Dryadorchis</i>	5	5	5	100	5	5	100
<i>Drymoanthus</i>	4	4	4	100	4	4	100
<i>Drymoda</i>	3	3	3	100	3	3	100
<i>Dyakia</i>	1	1	1	100	1	1	100
<i>Earina</i>	5	5	8	62.5	5	7	71.4
<i>Echinorhyncha</i>	5	5	5	100	5	5	100
<i>Echinosepala</i>	14	11	11	100	11	11	100
<i>Eclecticus</i>	1	1	1	100	1	1	100
<i>Eggelingia</i>	3	3	3	100	3	3	100
<i>Elleanthus</i>	112	112	112	100	112	112	100
<i>Eloyella</i>	10	10	10	100	10	10	100

<i>Embreea</i>	2	2	2	100	2	2	100
<i>Encyclia</i>	177	160	188	85.1	160	172	93.0
<i>Entomophobia</i>	1	1	1	100	1	1	100
<i>Eparmatostigma</i>	1	1	1	100	1	1	100
<i>Epiblastus</i>	21	21	24	87.5	21	22	95.5
<i>Epidendrum</i>	1305	1303	1560	83.5	1303	1434	90.9
<i>Epilyna</i>	2	2	2	100	2	2	100
<i>Epistephium</i>	1	1	22	4.5	1	21	4.8
<i>Erasanthe</i>	1	1	1	100	1	1	100
<i>Eria</i>	241	234	258	90.7	234	238	98.3
<i>Eriodes</i>	1	1	1	100	1	1	100
<i>Eriopsis</i>	5	5	5	100	5	5	100
<i>Erycina</i>	7	7	8	87.5	7	7	100
<i>Esmeralda</i>	3	3	3	100	3	3	100
<i>Euanthe</i>	1	1	1	100	1	1	100
<i>Euchile</i>	3	3	3	100	3	3	100
<i>Eulophiella</i>	3	3	7	42.9	3	5	60.0
<i>Euryblema</i>	2	2	2	100	2	2	100
<i>Eurychone</i>	2	2	2	100	2	2	100
<i>Eurystyles</i>	19	19	21	90.5	19	20	95.0
<i>Fernandezia</i>	58	52	52	100	51	51	100
<i>Frondaria</i>	1	1	1	100	1	1	100
<i>Galeandra</i>	39	39	43	90.7	39	39	100
<i>Galeottia</i>	12	12	14	85.7	12	12	100
<i>Gastrochilus</i>	59	55	64	85.9	55	57	96.5
<i>Gastrorchis</i>	3	3	8	37.5	3	8	37.5
<i>Geesinkorchis</i>	4	4	4	100	4	4	100
<i>Genyorchis</i>	10	10	10	100	10	10	100
<i>Glomera</i>	167	130	133	97.7	130	131	99.2
<i>Gomesa</i>	125	117	134	87.3	117	124	94.4

<i>Gomphichis</i>	2	2	25	8.0	2	24	8.3
<i>Gongora</i>	76	74	77	96.1	74	74	100
<i>Goodyera</i>	4	4	113	3.5	4	99	4.0
<i>Grammangis</i>	2	2	2	100	2	2	100
<i>Grammatophyllum</i>	12	12	14	85.7	12	12	100
<i>Grandiphyllum</i>	8	7	7	100	7	7	100
<i>Graphorkis</i>	4	4	4	100	4	4	100
<i>Grastidium</i>	1	1	2	50.0	1	1	100
<i>Grobya</i>	5	5	5	100	5	5	100
<i>Grosourdya</i>	23	11	11	100	11	11	100
<i>Guanchezia</i>	1	1	1	100	1	1	100
<i>Guarianthe</i>	4	4	5	80.0	4	5	80.0
<i>Gunnarella</i>	9	9	9	100	9	9	100
<i>Gynoglottis</i>	1	1	1	100	1	1	100
<i>Habenaria</i>	1	1	902	<0.1	1	837	0.1
<i>Hagsatera</i>	2	2	2	100	2	2	100
<i>Haraella</i>	1	1	1	100	1	1	100
<i>Hederorkis</i>	2	2	2	100	2	2	100
<i>Helleriella</i>	2	2	2	100	2	2	100
<i>Herminium</i>	1	1	27	3.7	1	22	4.5
<i>Heterotaxis</i>	13	13	13	100	13	13	100
<i>Hintonella</i>	1	1	1	100	1	1	100
<i>Hippeophyllum</i>	10	10	10	100	10	10	100
<i>Hoehneella</i>	2	2	2	100	2	2	100
<i>Hofmeisterella</i>	3	2	2	100	2	2	100
<i>Holcoglossum</i>	18	14	14	100	14	14	100
<i>Homalopetalum</i>	9	8	9	88.9	8	8	100
<i>Horichia</i>	1	1	1	100	1	1	100
<i>Horvatia</i>	1	1	1	100	1	1	100
<i>Houlletia</i>	8	8	11	72.7	8	9	88.9

<i>Huntleya</i>	16	14	15	93.3	14	14	100
<i>Hygrochilus</i>	1	1	1	100	1	1	100
<i>Hylaeorchis</i>	1	1	1	100	1	1	100
<i>Hymenorchis</i>	11	11	12	91.7	11	12	91.7
<i>India</i>	1	1	1	100	1	1	100
<i>Inti</i>	2	2	2	100	2	2	100
<i>Ionopsis</i>	6	6	8	75.0	6	6	100
<i>Isabelia</i>	3	3	4	75.0	3	4	75.0
<i>Ischnogyne</i>	1	1	1	100	1	1	100
<i>Isochilus</i>	13	13	14	92.9	13	13	100
<i>Ixyophora</i>	6	5	5	100	5	5	100
<i>Jacquinia</i>	12	12	12	100	12	12	100
<i>Jejewoodia</i>	6	6	6	100	6	6	100
<i>Jejosephia</i>	1	1	1	100	1	1	100
<i>Jumellea</i>	60	60	67	89.6	59	64	92.2
<i>Kefersteinia</i>	71	70	72	97.2	70	70	100
<i>Kegeliella</i>	4	4	4	100	4	4	100
<i>Koellensteinia</i>	12	12	17	70.6	12	17	70.6
<i>Kraenzlinella</i>	10	10	10	100	9	9	100
<i>Lacaena</i>	2	2	2	100	2	2	100
<i>Laelia</i>	25	24	62	38.7	24	24	100
<i>Lankesterella</i>	11	11	11	100	11	11	100
<i>Lemurella</i>	4	4	4	100	4	4	100
<i>Lemurorchis</i>	1	1	1	100	1	1	100
<i>Leochilus</i>	12	12	14	85.7	12	12	100
<i>Lepanthes</i>	1165	1095	1135	96.5	1090	1092	99.8
<i>Lepanthopsis</i>	47	43	45	95.6	43	43	100
<i>Leptotes</i>	9	9	10	90.0	9	9	100
<i>Liparis</i>	367	367	452	81.4	367	428	86.0
<i>Listrostachys</i>	1	1	4	25.0	1	1	100

<i>Lockhartia</i>	33	28	30	93.3	28	28	100
<i>Loefgrenianthus</i>	1	1	1	100	1	1	100
<i>Lueckelia</i>	1	1	1	100	1	1	100
<i>Lueddemannia</i>	3	3	3	100	3	3	100
<i>Luisia</i>	43	40	42	95.2	40	40	100
<i>Lycaste</i>	35	34	56	60.7	34	45	75.6
<i>Lycomormium</i>	6	5	5	100	5	5	100
<i>Macradenia</i>	13	11	12	91.7	11	11	100
<i>Macroclinium</i>	45	41	42	97.6	41	42	97.6
<i>Macropodanthus</i>	10	8	8	100	8	8	100
<i>Malaxis</i>	6	6	200	3.0	6	183	3.3
<i>Malleola</i>	34	34	34	100	34	34	100
<i>Mapinguari</i>	4	4	5	80.0	4	5	80.0
<i>Margelliantha</i>	6	6	6	100	6	6	100
<i>Masdevallia</i>	638	597	625	95.5	596	598	99.7
<i>Maxillaria</i>	400	321	358	89.7	317	319	99.4
<i>Maxillariella</i>	45	45	46	97.8	45	46	97.8
<i>Mediocalcar</i>	17	17	18	94.4	17	17	100
<i>Megalotus</i>	1	1	1	100	1	1	100
<i>Meiracyllium</i>	2	2	2	100	2	2	100
<i>Mesospinidium</i>	1	1	2	50.0		0	
<i>Microchilus</i>	1	1	137	0.7	1	137	0.7
<i>Microcoelia</i>	33	30	30	100	30	30	100
<i>Microepidendrum</i>	1	1	1	100	1	1	100
<i>Micropera</i>	22	22	23	95.7	21	21	100
<i>Microsaccus</i>	12	12	12	100	12	12	100
<i>Microtatorchis</i>	51	51	51	100	51	51	100
<i>Microterangis</i>	7	7	7	100	7	7	100
<i>Miltonia</i>	12	12	21	57.1	12	18	66.7
<i>Miltoniopsis</i>	5	5	6	83.3	5	5	100

<i>Mobilabium</i>	1	1	1	100	1	1	100
<i>Monomeria</i>	3	3	3	100	3	3	100
<i>Mormodes</i>	83	83	93	89.2	83	83	100
<i>Mormolyca</i>	26	26	27	96.3	26	26	100
<i>Muscarella</i>	7	3	3	100	0	0	
<i>Mycaranthes</i>	35	35	36	97.2	35	36	97.2
<i>Myoxanthus</i>	50	48	49	98.0	48	48	100
<i>Myrmecophila</i>	10	10	13	76.9	10	11	90.9
<i>Mystacidium</i>	10	10	12	83.3	10	10	100
<i>Nabalua</i>	3	3	3	100	3	3	100
<i>Neobathiea</i>	6	6	6	100	5	5	100
<i>Neocogniauxia</i>	2	2	2	100	2	2	100
<i>Neofinetia</i>	3	3	3	100	3	3	100
<i>Neogardneria</i>	1	1	1	100	1	1	100
<i>Neogyna</i>	1	1	1	100	1	1	100
<i>Neomoorea</i>	1	1	1	100	1	1	100
<i>Nephrangis</i>	2	2	2	100	2	2	100
<i>Nidema</i>	2	2	2	100	2	2	100
<i>Nitidobulbon</i>	3	3	3	100	3	3	100
<i>Nohawilliamsia</i>	1	1	1	100	1	1	100
<i>Notheria</i>	1	1	1	100	1	1	100
<i>Nothodoritis</i>	1	1	1	100	1	1	100
<i>Notylia</i>	56	56	59	94.9	56	56	100
<i>Notyliopsis</i>	1	1	1	100	1	1	100
<i>Oberonia</i>	333	322	330	97.6	322	324	99.4
<i>Octarrhena</i>	54	52	52	100	52	52	100
<i>Octomeria</i>	166	160	174	92.0	160	162	98.8
<i>Oeceoclades</i>	1	1	41	2.4	1	38	2.6
<i>Oeonia</i>	5	5	5	100	5	5	100
<i>Oeoniella</i>	2	2	2	100	2	2	100

<i>Oestlundia</i>	4	4	4	100	4	4	100
<i>Oliveriana</i>	13	6	6	100	6	6	100
<i>Omoea</i>	2	2	2	100	2	2	100
<i>Oncidium</i>	342	335	393	85.2	334	341	97.9
<i>Ophioglossella</i>	1	1	1	100	1	1	100
<i>Oreorchis</i>	1	1	20	5.0	1	16	6.3
<i>Orleanesia</i>	9	9	9	100	9	9	100
<i>Ornithidium</i>	56	56	63	88.9	56	57	98.2
<i>Ornithocephalus</i>	56	56	58	96.6	56	56	100
<i>Ornithochilus</i>	3	3	5	60.0	3	3	100
<i>Ossiculum</i>	1	1	1	100	1	1	100
<i>Otochilus</i>	5	5	5	100	5	5	100
<i>Otoglossum</i>	21	13	15	86.7	13	15	86.7
<i>Otostylis</i>	1	1	4	25.0	1	4	25.0
<i>Oxystophyllum</i>	36	36	36	100	36	36	100
<i>Pabstia</i>	5	5	5	100	5	5	100
<i>Pabstiella</i>	73	28	35	80.0	28	29	96.6
<i>Pachyphyllum</i>	1	1	4	25.0		0	
<i>Panisea</i>	12	10	11	90.9	10	11	90.9
<i>Paphinia</i>	16	16	16	100	16	16	100
<i>Paphiopedilum</i>	7	7	136	5.1	7	118	5.9
<i>Papilionanthe</i>	9	9	11	81.8	9	11	81.8
<i>Papillilabium</i>	1	1	1	100	1	1	100
<i>Paralophia</i>	2	2	2	100	2	2	100
<i>Paraphalaenopsis</i>	4	4	5	80.0	4	5	80.0
<i>Parapteroceras</i>	8	8	8	100	8	8	100
<i>Pedilochilus</i>	38	38	38	100	36	36	100
<i>Pelatantheria</i>	8	8	8	100	8	8	100
<i>Pelexia</i>	1	1	79	1.3	1	78	1.3
<i>Penkimia</i>	1	1	1	100	1	1	100

<i>Pennilabium</i>	18	15	15	100	15	15	100
<i>Peristeranthus</i>	1	1	1	100	1	1	100
<i>Peristeria</i>	13	13	14	92.9	13	13	100
<i>Pescatoria</i>	25	25	30	83.3	25	26	96.2
<i>Phaius</i>	1	1	58	1.7	1	45	2.2
<i>Phalaenopsis</i>	74	61	77	79.2	61	70	87.1
<i>Phloeophila</i>	12	11	12	91.7	11	11	100
<i>Pholidota</i>	38	38	45	84.4	38	39	97.4
<i>Phragmipedium</i>	8	8	36	22.2	8	27	29.6
<i>Phragmorchis</i>	1	1	1	100	1	1	100
<i>Phreatia</i>	220	211	214	98.6	211	211	100
<i>Phyllorchis</i>	1	1	32	3.1	1	12	8.3
<i>Phymatidium</i>	10	10	11	90.9	10	10	100
<i>Physoceras</i>	1	1	12	8.3	1	12	8.3
<i>Pinalia</i>	116	101	106	95.3	101	106	95.3
<i>Pityphyllum</i>	5	5	5	100	5	5	100
<i>Platanthera</i>	3	3	172	1.7	3	152	2.0
<i>Platyrrhiza</i>	1	1	1	100	1	1	100
<i>Platystele</i>	112	101	102	99.0	101	101	100
<i>Plectorrhiza</i>	3	3	3	100	3	3	100
<i>Plectrelminthus</i>	1	1	1	100	1	1	100
<i>Plectrophora</i>	10	10	10	100	10	10	100
<i>Pleione</i>	12	12	28	42.9	12	26	46.2
<i>Pleurothallis</i>	618	559	615	90.9	556	557	99.8
<i>Pleurothallopsis</i>	18	18	18	100	18	18	100
<i>Plocoglottis</i>	1	1	42	2.4	1	41	2.4
<i>Poaephyllum</i>	7	6	6	100	6	6	100
<i>Podangis</i>	2	1	1	100	1	1	100
<i>Podochilus</i>	63	61	66	92.4	61	62	98.4
<i>Polycycnis</i>	17	17	17	100	17	17	100

<i>Polyotidium</i>	1	1	1	100	1	1	100
<i>Polystachya</i>	249	241	253	95.3	241	245	98.4
<i>Pomatocalpa</i>	25	25	25	100	25	25	100
<i>Ponera</i>	8	8	14	57.1	8	8	100
<i>Ponerorchis</i>	1	1	20	5.0	1	20	5.0
<i>Ponthieva</i>	2	2	68	2.9	2	65	3.1
<i>Porpax</i>	21	13	13	100	13	13	100
<i>Porphyrodesme</i>	3	3	3	100	3	3	100
<i>Porphyroglottis</i>	1	1	1	100	1	1	100
<i>Porroglossum</i>	53	43	43	100	43	43	100
<i>Porrhachis</i>	2	2	2	100	2	2	100
<i>Prescottia</i>	2	2	28	7.1	2	26	7.7
<i>Promenaea</i>	19	18	18	100	18	18	100
<i>Prosthechea</i>	116	116	119	97.5	116	118	98.3
<i>Pseuderia</i>	20	20	20	100	20	20	100
<i>Pseudocentrum</i>	2	2	7	28.6	2	7	28.6
<i>Pseudolaelia</i>	16	16	19	84.2	16	19	84.2
<i>Psychilis</i>	15	15	17	88.2	15	17	88.2
<i>Psychopsis</i>	5	5	5	100	5	5	100
<i>Pteroceras</i>	27	27	27	100	27	27	100
<i>Pterostemma</i>	4	3	3	100	3	3	100
<i>Pygmaeorchis</i>	2	2	2	100	2	2	100
<i>Quekettia</i>	6	5	5	100	5	5	100
<i>Quisqueya</i>	4	4	4	100	4	4	100
<i>Rangaeris</i>	7	7	7	100	7	7	100
<i>Rauhiella</i>	3	3	3	100	3	3	100
<i>Renanthera</i>	21	20	22	90.9	20	20	100
<i>Restrepia</i>	58	53	55	96.4	53	53	100
<i>Restrepiella</i>	4	2	2	100	2	2	100
<i>Rhaesteria</i>	1	1	1	100	1	1	100

<i>Rhetinantha</i>	14	14	14	100	14	14	100
<i>Rhinerrhiza</i>	1	1	1	100	1	1	100
<i>Rhinerrhizopsis</i>	3	3	3	100	3	3	100
<i>Rhipidoglossum</i>	37	35	35	100	35	35	100
<i>Rhynchogyna</i>	3	3	3	100	3	3	100
<i>Rhyncholaelia</i>	2	2	2	100	2	2	100
<i>Rhynchostele</i>	17	17	20	85.0	17	20	85.0
<i>Rhynchostylis</i>	4	3	4	75.0	3	3	100
<i>Ridleyella</i>	1	1	1	100	1	1	100
<i>Robiquetia</i>	82	45	45	100	45	45	100
<i>Rodriguezia</i>	50	50	55	90.9	50	50	100
<i>Rossioglossum</i>	9	9	9	100	9	9	100
<i>Rudolfiella</i>	6	6	6	100	6	6	100
<i>Saccoglossum</i>	5	5	5	100	5	5	100
<i>Saccolabiopsis</i>	14	14	15	93.3	14	15	93.3
<i>Saccolabium</i>	5	5	18	27.8	5	5	100
<i>Samarorchis</i>	1	1	1	100	1	1	100
<i>Sanderella</i>	2	2	2	100	2	2	100
<i>Sansonia</i>	1	1			1		
<i>Santanderella</i>	1	1	1	100	1	1	100
<i>Santotomasia</i>	1	1	1	100	1	1	100
<i>Sarcanthopsis</i>	6	5	5	100	5	5	100
<i>Sarcochilus</i>	25	25	29	86.2	25	25	100
<i>Sarcoglyphis</i>	15	2	12	16.7	2	12	16.7
<i>Sarcophyton</i>	3	3	3	100	3	3	100
<i>Sarcostoma</i>	5	5	6	83.3	5	5	100
<i>Saundersia</i>	2	2	2	100	2	2	100
<i>Sauvetrea</i>	11	11	11	100	11	11	100
<i>Scaphosepalum</i>	51	46	47	97.9	46	47	97.9
<i>Scaphyglottis</i>	77	69	79	87.3	69	69	100

<i>Schistotylus</i>	1	1	1	100	1	1	100
<i>Schlimia</i>	7	7	7	100	7	7	100
<i>Schoenorchis</i>	28	24	25	96.0	24	24	100
<i>Schomburgkia</i>	1	1	4	25.0	1	2	50.0
<i>Schunkea</i>	1	1	1	100	1	1	100
<i>Scuticaria</i>	12	10	11	90.9	10	10	100
<i>Sedirea</i>	2	2	2	100	2	2	100
<i>Seegeriella</i>	3	2	2	100	2	2	100
<i>Seidenfadenia</i>	1	1	1	100	1	1	100
<i>Seidenfadeniella</i>	3	1	2	50.0	1	2	50.0
<i>Sertifera</i>	2	2	7	28.6	2	7	28.6
<i>Sievekingia</i>	16	16	17	94.1	16	16	100
<i>Singchia</i>	1	1	1	100	1	1	100
<i>Sirhookera</i>	2	2	2	100	2	2	100
<i>Smithsonia</i>	3	3	3	100	3	3	100
<i>Smitinandia</i>	3	3	3	100	3	3	100
<i>Sobennikoffia</i>	2	2	4	50.0	2	4	50.0
<i>Sobralia</i>	31	30	161	18.6	30	150	20.0
<i>Solenangis</i>	8	8	8	100	8	8	100
<i>Solenidium</i>	3	3	4	75.0	3	3	100
<i>Solenocentrum</i>	1	1	4	25.0	1	4	25.0
<i>Soterosanthus</i>	1	1	1	100	1	1	100
<i>Spathoglottis</i>	4	4	51	7.8	4	49	8.2
<i>Specklinia</i>	158	135	139	97.1	135	135	100
<i>Sphyrarhynchus</i>	1	1	1	100	1	1	100
<i>Spiranthes</i>	1	1	62	1.6	1	39	2.6
<i>Spongiola</i>	1	1	1	100	1	1	100
<i>Stalkya</i>	1	1	1	100	1	1	100
<i>Stanhopea</i>	76	62	88	70.5	62	68	91.2
<i>Staurochilus</i>	14	14	14	100	14	14	100

<i>Stelis</i>	1130	886	912	97.1	884	887	99.7
<i>Stenia</i>	22	22	22	100	22	22	100
<i>Stenorrhynchos</i>	1	1	15	6.7	1	7	14.3
<i>Stenotyla</i>	9	9	9	100	9	9	100
<i>Stereochilus</i>	7	7	7	100	7	7	100
<i>Stichorkis</i>	18	8	9	88.9	8	8	100
<i>Stolzia</i>	16	16	16	100	15	15	100
<i>Suarezia</i>	1	1	1	100	1	1	100
<i>Sudamerlycaste</i>	33	33	45	73.3	33	45	73.3
<i>Summerhayesia</i>	2	2	2	100	2	2	100
<i>Sunipia</i>	24	24	24	100	24	24	100
<i>Sutrina</i>	2	2	2	100	2	2	100
<i>Systeloglossum</i>	5	5	5	100	5	5	100
<i>Taeniophyllum</i>	233	185	188	98.4	185	185	100
<i>Taeniorrhiza</i>	1	1	1	100	1	1	100
<i>Tainia</i>	6	6	34	17.6	6	32	18.8
<i>Teagueia</i>	18	13	13	100	13	13	100
<i>Telipogon</i>	229	207	215	96.3	206	206	100
<i>Tetramicra</i>	1	1	15	6.7	1	14	7.1
<i>Teuscheria</i>	10	7	7	100	7	7	100
<i>Theana</i>	1	1			1		
<i>Thecopus</i>	2	2	2	100	2	2	100
<i>Thecostele</i>	1	1	1	100	1	1	100
<i>Thelasis</i>	28	26	26	100	26	26	100
<i>Thrixspermum</i>	186	160	163	98.2	160	161	99.4
<i>Thunia</i>	3	3	7	42.9	3	5	60.0
<i>Thysanoglossa</i>	2	2	3	66.7	2	3	66.7
<i>Tipularia</i>	1	1	8	12.5	1	7	14.3
<i>Tolumnia</i>	25	25	31	80.6	25	31	80.6
<i>Tomzanonia</i>	1	1	1	100	1	1	100

<i>Trachoma</i>	8	8			8		
<i>Trevoria</i>	5	5	5	100	5	5	100
<i>Trias</i>	13	13	14	92.9	13	14	92.9
<i>Triceratorhynchus</i>	1	1	1	100	1	1	100
<i>Trichocentrum</i>	92	64	73	87.7	63	65	96.9
<i>Trichoceros</i>	10	10	11	90.9	10	10	100
<i>Trichoglottis</i>	77	69	72	95.8	69	69	100
<i>Trichopilia</i>	45	45	49	91.8	45	45	100
<i>Trichosalpinx</i>	124	113	117	96.6	112	112	100
<i>Trichotosia</i>	82	78	78	100	78	78	100
<i>Tridactyle</i>	50	47	47	100	47	47	100
<i>Trigonidium</i>	13	13	14	92.9	13	13	100
<i>Trisetella</i>	24	23	23	100	23	23	100
<i>Trizeuxis</i>	1	1	1	100	1	1	100
<i>Tuberolabium</i>	18	18	18	100	18	18	100
<i>Uncifera</i>	6	6	6	100	6	6	100
<i>Vanda</i>	81	56	66	84.8	56	60	93.3
<i>Vandopsis</i>	3	3	6	50.0	3	4	75.0
<i>Vanilla</i>	2	2	122	1.6	2	106	1.9
<i>Vasqueziella</i>	1	1	1	100	1	1	100
<i>Ventricularia</i>	2	2	2	100	2	2	100
<i>Vitekorchis</i>	4	4	4	100	4	4	100
<i>Warczewiczella</i>	11	11	13	84.6	11	11	100
<i>Warmingia</i>	4	4	4	100	4	4	100
<i>Warrea</i>	2	2	3	66.7	2	3	66.7
<i>Warreella</i>	2	2	2	100	2	2	100
<i>Warreopsis</i>	4	4	4	100	4	4	100
<i>Xenikophyton</i>	2	2	2	100	2	2	100
<i>Xylobium</i>	38	33	33	100	31	31	100
<i>Ypsilopus</i>	5	5	5	100	5	5	100

	<i>Ypsilorchis</i>	1	1	1	100	1	1	100	
	<i>Zelenkoa</i>	1	1	1	100	1	1	100	
	<i>Zootrophion</i>	28	21	22	95.5	21	21	100	
	<i>Zygopetalum</i>	11	11	27	40.7	11	15	73.3	
	<i>Zygosepalum</i>	8	8	8	100	8	8	100	
	<i>Zygostates</i>	24	22	22	100	22	22	100	
Pandanaceae		7	7	1086	0.6	7	1062	0.7	
	<i>Benstonea</i>	4	4	0		4	0		
	<i>Freycinetia</i>	1	1	310	0.3	1	308	0.3	
	<i>Pandanus</i>	2	2	767	0.3	2	746	0.3	
Paulowniaceae		2	2	18	11.1	1	9	11.1	H
	<i>Wightia</i>	2	2	6	33.3	1	1	100	H
Phyllonomaceae		1	1	5	20.0	1	5	20.0	
	<i>Phyllonoma</i>	1	1	5	20.0	1	5	20.0	
Piperaceae		661	658	4942	13.3	509	2656	19.2	
	<i>Manekia</i>	1	1	6	16.7	1	5	20.0	
	<i>Peperomia</i>	659	656	1868	35.1	507	1160	43.7	
	<i>Piperanthera</i>	1	1	1	100	1	1	100	
Pittosporaceae		3	3	472	0.6	2	170	1.2	
	<i>Pittosporum</i>	3	3	345	0.9	2	110	1.8	
Poaceae		1		14789	<0.1		11548	<0.1	
	<i>Tripogon</i>	1		48	<0.1		46	<0.1	
Primulaceae		8	8	4947	0.2	8	2787	0.3	
	<i>Cybianthus</i>	7	7	173	4.0	7	160	4.4	
	<i>Myrsine</i>	1	1	301	0.3	1	131	0.8	
Rapateaceae		6	5	98	5.1	5	95	5.3	
	<i>Epidryos</i>	4	3	3	100	3	3	100	
	<i>Stegolepis</i>	2	2	33	6.1	2	33	6.1	
Rosaceae		6	5	19262	<0.1	4	4820	0.1	H
	<i>Sorbus</i>	6	5	537	0.7	4	243	1.2	H

Rubiaceae	263	253	15103	1.7	251	13668	1.8	
<i>Acranthera</i>	1	1	40	2.5	1	38	2.6	
<i>Aidia</i>	1	1	55	1.8	1	55	1.8	
<i>Anthorrhiza</i>	9	9	9	100	9	9	100	
<i>Argostemma</i>	2	2	169	1.2	2	163	1.2	
<i>Balmea</i>	1	1	1	100	1	1	100	
<i>Chassalia</i>	3	3	115	2.6	3	112	2.7	
<i>Coprosma</i>	2	2	125	1.6	2	118	1.7	
<i>Cosmibuena</i>	3	3	5	60.0	3	4	75.0	H
<i>Didymochlamys</i>	2	2	3	66.7	2	2	100	
<i>Gardenia</i>	1	1	163	0.6	1	134	0.7	
<i>Hillia</i>	22	22	30	73.3	21	24	87.5	
<i>Hoffmannia</i>	2	2	139	1.4	2	131	1.5	
<i>Hydnophytum</i>	96	94	94	100	94	94	100	
<i>Hymenodictyon</i>	4	4	23	17.4	4	23	17.4	
<i>Ladenbergia</i>	1	1	41	2.4	1	35	2.9	
<i>Lecananthus</i>	3	3	3	100	3	3	100	
<i>Leucocodon</i>	1	1	1	100	1	1	100	
<i>Manettia</i>	1	1	132	0.8	1	123	0.8	
<i>Mussaenda</i>	1	1	198	0.5	1	184	0.5	
<i>Mycetia</i>	1	1	46	2.2	1	45	2.2	
<i>Myrmecodia</i>	27	27	27	100	27	27	100	
<i>Myrmephytum</i>	5	5	5	100	5	5	100	
<i>Neohymenopogon</i>	1	1	3	33.3	1	3	33.3	
<i>Notopleura</i>	13	13	102	12.7	13	102	12.7	
<i>Pentapterygium</i>	1	1	2	50.0		0		
<i>Psychotria</i>	2	2	1989	<0.1	2	1865	0.1	
<i>Rudgea</i>	1	1	141	0.7	1	133	0.8	
<i>Schradera</i>	13	13	56	23.2	13	55	23.6	
<i>Squamellaria</i>	12	4	4	100	4	4	100	

	<i>Timonius</i>	31	31	182	17.0	31	182	17.0	H
Schlegeliaceae		16	16	41	39.0	16	37	43.2	
	<i>Gibsoniothamnus</i>	12	12	12	100	12	12	100	
	<i>Schlegelia</i>	4	4	27	14.8	4	23	17.4	
Scrophulariaceae		1	1	3815	<0.1	1	1545	0.1	
	<i>Dermatobotrys</i>	1	1	1	100	1	1	100	
Solanaceae		52	52	2467	2.1	49	2433	2.0	
	<i>Dyssochroma</i>	1	1	1	100	1	1	100	H
	<i>Juanulloa</i>	10	10	11	90.9	10	11	90.9	
	<i>Lycianthes</i>	5	5	98	5.1	4	98	4.1	H
	<i>Markea</i>	12	12	22	54.5	12	22	54.5	H
	<i>Merinthopodium</i>	3	3	3	100	3	3	100	
	<i>Salpichroa</i>	1	1	14	7.1		14	<0.1	
	<i>Schultesianthus</i>	7	7	8	87.5	7	8	87.5	
	<i>Solandra</i>	4	4	12	33.3	4	10	40.0	
	<i>Solanum</i>	4	4	1203	0.3	4	1203	0.3	
	<i>Trianaea</i>	5	5	5	100	4	5	80.0	
Urticaceae		83	83	3353	2.5	73	1465	5.0	
	<i>Coussapoa</i>	39	39	51	76.5	39	48	81.3	H
	<i>Elatostema</i>	12	12	597	2.0	11	525	2.1	
	<i>Pilea</i>	13	13	683	1.9	13	279	4.7	
	<i>Poikilospermum</i>	6	6	41	14.6	4	4	100	H
	<i>Procris</i>	9	9	77	11.7	2	2	100	
	<i>Urera</i>	4	4	97	4.1	4	43	9.3	
Zingiberaceae		33	32	1689	1.9	32	1587	2.0	
	<i>Alpinia</i>	1		253	<0.1		244	<0.1	
	<i>Amomum</i>	4	4	196	2.0	4	179	2.2	
	<i>Burbridgea</i>	5	5	5	100	5	5	100	
	<i>Cautleya</i>	2	2	2	100	2	2	100	
	<i>Globba</i>	1	1	102	1.0	1	98	1.0	

<i>Hedychium</i>	15	15	99	15.2	15	93	16.1
<i>Rhynchanthus</i>	4	4	5	80.0	4	4	100
<i>Riedelia</i>	1	1	78	1.3	1	75	1.3
“gymnosperms”	2	2	1288	0.2	2	1103	0.2
Zamiaceae	2	2	230	0.9	2	215	0.9
<i>Zamia</i>	2	2	73	2.7	2	65	3.1

Table 2. Systematic distribution of epiphytic ferns and lycopods. For each class, family, and genus, the numbers of epiphytic/hemiepiphytic species (“E/H”) and the total number of species (“Total”) are given. “%Total” gives the proportion of epiphytic species per taxon in %. An “H” marks genera, which have many or only hemiepiphytes, with “H?” indicating doubtful cases. Totals are from Hassler and Schmitt (2019).

Class/Family	Genus	E / H	Total	% Total
Lycopodiopsida		142	1360	10
Lycopodiaceae		114	480	24
	<i>Huperzia</i>	2	25	8
	<i>Palinhaea</i>	1	25	4
	<i>Phlegmariurus</i>	111	320	35
Selaginellaceae		28	700	4
	<i>Selaginella</i>	28	700	4
Polypodiopsida		2942	11000	27
Psilotaceae		18	18	100
	<i>Psilotum</i>	2	2	100
	<i>Tmesipteris</i>	16	16	100
Ophioglossaceae		5	110	5
	<i>Cheiroglossa</i>	2	2	100
	<i>Ophioderma</i>	2	5	40
	<i>Ophioglossum</i>	1	50	2
Hymenophyllaceae		433	600	72
	<i>Abrodictyum</i>	13	35	37
	<i>Callistopteris</i>	2	5	40
	<i>Cephalomanes</i>	1	5	20
	<i>Crepidomanes</i>	33	48	71
	<i>Didymoglossum</i>	48	52	92
	<i>Hymenophyllum</i>	262	300	89
	<i>Polyphlebium</i>	15	17	88
	<i>Trichomanes</i>	37	65	57
	<i>Vandenboschia</i>	22	24	92 H
Schizaeaceae		6	33	18
	<i>Schizaea</i>	2	18	11
	<i>Actinostachys</i>	4	15	27
Cyatheaceae		2	650	<1
	<i>Cyathea</i>	2	300	<1
Lindsaeaceae		30	230	13
	<i>Lindsaea</i>	28	170	16
	<i>Odontosoria</i>	1	30	3
	<i>Sphenomeris</i>	1	3	33
Pteridaceae		126	1300	10
	<i>Ananthacorus</i>	1	1	100
	<i>Antrophyopsis</i>	4	4	100

	<i>Antrophyum</i>	32	32	100	
	<i>Coniogramme</i>	1	25	4	
	<i>Haplopteris</i>	33	33	100	
	<i>Hecistopteris</i>	3	3	100	
	<i>Paragymnopteris</i>	1	5	20	
	<i>Polytaenium</i>	13	13	100	
	<i>Radiovittaria</i>	7	7	100	
	<i>Rheopteris</i>	1	1	100	
	<i>Scoliosorus</i>	1	1	100	
	<i>Vaginularia</i>	8	8	100	
	<i>Vittaria</i>	21	21	100	
Dennstaedtiaceae		4	230	2	
	<i>Dennstaedtia</i>	1	55	2	
	<i>Microlepia</i>	3	50	6	
Aspleniaceae		259	745	35	
	<i>Asplenium</i>	246	690	36	
	<i>Hymenasplenium</i>	13	55	24	
Blechnaceae		20	245	8	
	<i>Austroblechnum</i>	1	35	3	
	<i>Blechnum</i>	1	22	5	H
	<i>Icarus</i>	1	1	100	
	<i>Lomaridium</i>	15	15	100	H
	<i>Lomariocycas</i>	1	18	6	
	<i>Parablechnum</i>	1	60	2	
Hypodematiaceae		2	17	12	
	<i>Leucostegia</i>	2	2	100	
Dryopteridaceae		424	2100	20	
	<i>Arachniodes</i>	2	80	3	
	<i>Ctenitis</i>	1	140	1	
	<i>Dryopteris</i>	1	320	<1	
	<i>Elaphoglossum</i>	415	730	57	
	<i>Rumohra</i>	4	9	44	
	<i>Stigmatopteris</i>	1	25	4	
Nephrolepidaceae		19	25	76	
	<i>Nephrolepis</i>	19	25	76	
Lomariopsidaceae		50	55	91	
	<i>Lomariopsis</i>	50	50	100	H?
Tectariaceae		12	310	4	
	<i>Arthropteris</i>	12	15	80	H
Oleandraceae		22	25	88	
	<i>Oleandra</i>	22	25	88	
Davalliaceae		61	61	100	
	<i>Davallia</i>	61	61	100	
Polypodiaceae		1450	1629	89	
	<i>Acrosorus</i>	6	8	75	
	<i>Adenophorus</i>	9	10	90	
	<i>Adetogramma</i>	1	1	100	

<i>Aglaomorpha</i>	32	33	97
<i>Alansmia</i>	25	25	100
<i>Archigrammitis</i>	6	6	100
<i>Ascogrammitis</i>	18	18	100
<i>Bosmania</i>	1	3	33
<i>Calymmodon</i>	24	25	96
<i>Campyloneurum</i>	53	59	90
<i>Ceradenia</i>	68	78	87
<i>Chrysogrammitis</i>	3	3	100
<i>Cochlidium</i>	16	18	89
<i>Ctenopterella</i>	15	18	83
<i>Dasygrammitis</i>	6	6	100
<i>Dendroconche</i>	8	8	100
<i>Dictymia</i>	2	2	100
<i>Ellipinema</i>	1	1	100
<i>Enterosora</i>	18	18	100
<i>Galactodenia</i>	5	5	100
<i>Goniophlebium</i>	14	14	100
<i>Grammitis</i>	39	40	98
<i>Gymnogrammitis</i>	1	1	100
<i>Hovenkampia</i>	3	3	100
<i>Lecanopteris</i>	13	13	100
<i>Lellingeria</i>	45	54	83
<i>Lemmaphyllum</i>	6	7	86
<i>Lepisorus</i>	60	68	88
<i>Leptochilus</i>	18	33	55
<i>Leucotrichum</i>	6	6	100
<i>Lomaphlebia</i>	2	2	100
<i>Loxogramme</i>	30	36	83
<i>Luisma</i>	1	1	100
<i>Melpomene</i>	25	30	83
<i>Microgramma</i>	31	31	100
<i>Micropolypodium</i>	3	3	100
<i>Microsorium</i>	24	30	80
<i>Moranopteris</i>	28	30	93
<i>Mycopteris</i>	15	18	83
<i>Neocheiropteris</i>	2	5	40
<i>Neolepisorus</i>	3	3	100
<i>Niphidium</i>	8	11	73
<i>Notogrammitis</i>	9	12	75
<i>Oreogrammitis</i>	142	176	81
<i>Parrisia</i>	1	1	100
<i>Pecluma</i>	42	51	82
<i>Phlebodium</i>	3	3	100
<i>Phymatosorus</i>	13	17	76
<i>Pichisermollodes</i>	14	15	93
<i>Platyserium</i>	16	18	89

H

<i>Pleopeltis</i>	76	90	84
<i>Pleurosoriopsis</i>	1	1	100
<i>Polypodiodes</i>	20	21	95
<i>Polypodium</i>	37	40	92
<i>Prosaptia</i>	55	55	100
<i>Pyrrosia</i>	62	65	95
<i>Scleroglossum</i>	10	10	100
<i>Selliguea</i>	108	115	95
<i>Serpocaulon</i>	47	47	100
<i>Stenogrammitis</i>	19	25	76
<i>Synammia</i>	3	3	100
<i>Terpsichore</i>	15	15	100
<i>Thylacopteris</i>	1	1	100
<i>Tomophyllum</i>	28	28	100
<i>Tricholepidium</i>	5	5	100
<i>Xiphopterella</i>	9	11	82
<i>Zealandia</i>	4	4	100
<i>Zygophlebia</i>	16	16	100
